

Response of recruitment to light availability across a tropical lowland rainforest community

Nadja Rüger^{1*}, Andreas Huth¹, Stephen P. Hubbell² and Richard Condit^{3,4}

¹Department of Ecological Modelling, Helmholtz Centre for Environmental Research – UFZ, Permoserstr. 15, 04318 Leipzig, Germany; ²Department of Ecology and Evolutionary Biology, University of California Los Angeles, Los Angeles, CA 90095, USA; ³Center for Tropical Forest Science, Smithsonian Tropical Research Institute, Unit 0948, APO AA 34002-0948, USA; and ⁴National Center for Ecological Analysis and Synthesis, University of California, Santa Barbara, CA 93101, USA

Summary

1. Many hypotheses about species coexistence involve differential resource use and trade-offs in species' life-history traits. Quantifying resource use across most species in diverse communities, although, has seldom been attempted.
2. We use a hierarchical Bayesian approach to quantify the light dependence of recruitment in 263 woody species in a 50-ha long-term forest census plot in Panama. Data on sapling recruitment were obtained using the 1985–1990 and 1990–1995 census intervals. Available light was estimated for each recruit from yearly censuses of canopy density.
3. We use a power function (linear log–log relationship) to model the light effect on recruitment. Different responses of recruitment to light are expressed by the light effect parameter b . The distribution of b had a central mode at 0.8, suggesting that recruitment of many species responds nearly linearly to increasing light.
4. Nearly every species showed increases in recruitment with increasing light. Just nine species (3%) had recruitment declining with light, while 198 species (75%) showed increasing recruitment in both census intervals. Most of the increases in recruitment were decelerating, i.e. the increase was less at higher light ($b < 1$). In the remaining species, the response to light varied between census intervals (24 species) or species did not have recruits in both intervals (41 species).
5. *Synthesis.* Nearly all species regenerate better in higher light, and recruitment responses to light are spread along a continuum ranging from modest increase with light to a rather strict requirement for high light. These results support the hypothesis that spatio-temporal variation in light availability may contribute to the diversity of tropical tree species by providing opportunities for niche differentiation with respect to light requirements for regeneration.

Key-words: Barro Colorado Island, hierarchical Bayesian model, life-history traits, light requirements, niche partitioning, Panama, regeneration niche, shade tolerance, tropical rain forest

Introduction

Many hypotheses about species coexistence involve differential resource use and trade-offs in species' life-history traits (e.g. Tilman 1982, 2004; Rees *et al.* 2001; Silvertown 2004). Understorey light availability is one of the most limiting resources in tropical rain forests (Whitmore 1996), and its spatio-temporal

variation is an important component of the regeneration niche and gap dynamics hypotheses for species coexistence (Grubb 1977; Denslow 1987). Treefall gaps increase heterogeneity in light conditions and may contribute to the diversity of tropical tree species by providing opportunities for niche differentiation with respect to light requirements (Ricklefs 1977; Brokaw 1985).

Empirical studies quantifying niche differentiation in regeneration have yielded inconsistent conclusions. Several studies suggest that species differentially respond to light levels, even if variations are subtle, and that tight species packing along the small range of light environments may

*Correspondence author. Department of Forest Biometrics/Systems Analysis, Institute of Forest Growth and Forest Computer Sciences, Dresden University of Technology, Piennert Str. 8, 01737 Tharandt, Germany. E-mail: nadja.rueger@tu-dresden.de

contribute to coexistence (Denslow 1980; Kobe 1999; Svenning 2000; Poorter & Arets 2003). However, other studies conclude that many species in tropical rain forests share similar light requirements. While Hartshorn (1980) estimated that half of the tree species of the tropical lowland rain forest at La Selva, Costa Rica, required gaps for regeneration; other studies found that >80% of the tree species at Barro Colorado Island (BCI), Panama, regenerate at low light levels (Hubbell & Foster 1986; Welden *et al.* 1991; Condit, Hubbell & Foster 1996); and Wright *et al.* (2003) concluded that most of the same BCI species have intermediate light requirements.

Quantifying resource use in highly diverse communities is challenging because it theoretically involves measuring the continuous environmental variation of the resource and continuous variation in species responses (Lieberman, Lieberman & Peralta 1989; Wright *et al.* 2003). This is often not feasible, and previous studies have usually simplified the problem by either focusing on a small number of species (e.g. Brown & Whitmore 1992; Clark & Clark 1992; Van der Meer, Sterck & Bongers 1998; Svenning 2000; Poorter & Arets 2003; Uriarte *et al.* 2005; Queenborough *et al.* 2007), grouping species into broad growth form classes (Dupuy & Chazdon 2006), or using few categories of environmental conditions (e.g. gap/non-gap; Welden *et al.* 1991; Condit, Hubbell & Foster 1996; Barker, Press & Brown 1997).

Moreover, if forest inventories form the basis of the analyses, the samples usually contain only few individuals of the many rare species (Pitman *et al.* 1999). As a consequence, there are few studies quantifying the light response of recruitment across entire diverse tropical rain forest communities (but see Welden *et al.* 1991 for an analysis of 155 species).

In this study, we aim to quantify the response of recruitment to light availability across more than 80% of the 300 tree and shrub species occurring at BCI, Panama, and our approach explicitly addresses all three problems. Forest census data from the 50-ha Forest Dynamics Plot (FDP) at BCI provided information on the spatial location of more than 50 000 recruits of 263 species. Yearly canopy census data allowed an estimate of light availability across the 50-ha FDP with a spatial resolution of 5 m. We used a hierarchical Bayesian approach to quantify the response of recruitment to light availability for all species. The hierarchical Bayesian approach combines the probability models for the spatial distribution of recruits within species and the variation of the light response across species, including rare ones (e.g. Clark 2005; Condit *et al.* 2006). Moreover, the posterior distribution of parameter estimates provides direct information about the uncertainty associated with the parameter estimates (Ellison 2004).

We explore how the response of recruitment to light is distributed across the tree community and how it is related to the abundance of the species. Furthermore, we calculate a comparable measure of light requirements for recruitment, the light level at which recruitment falls to 50% of the maximum under high light. Our estimates of the sensitivity of recruitment to light availability can serve in a classification of tree species into functional groups (Swaine & Whitmore 1988).

Materials and methods

STUDY SITE

We analysed data from a 50-ha forest census plot on BCI, Panama (9°9'N, 79°51'W). Barro Colorado Island is a 1567-ha island in the Panama Canal, covered with tropical lowland moist forest. The plot consists of 48 ha of undisturbed old-growth forest and 2 ha of secondary forest about 100 years old (Foster & Brokaw 1982). The climate on BCI is warm throughout the year, but rainfall is seasonal with most of the 2500 mm falling from April to November (Windsor 1990; Windsor, Rand & Rand 1990). Elevation of the plot is 120–155 m a.s.l. (Hubbell & Foster 1983). Detailed descriptions of flora, fauna, geology and climate can be found in Croat (1978), Leigh, Rand & Windsor (1982) and Leigh (1999).

RECRUITMENT DATA

All freestanding woody individuals ≥ 1 cm diameter at breast height (d.b.h.) were mapped, identified to species and measured in 1981–1983, 1985 and every 5 years thereafter (<http://www.ctfs.si.edu>; Hubbell & Foster 1983; Condit 1998). In this study, we use the census intervals from 1985 to 1990 and 1990 to 1995 and define every individual absent in the first census of each interval but present in the second as a recruit. To avoid edge effects of the light availability calculation, we excluded all recruits within 20 m of any edge of the plot. In the 1990 census, 33 069 recruits of 252 species were recorded, and in the 1995 census, 17 809 recruits of 233 species. We divided the plot (excluding the 20 m perimeter) into 17 664 5 × 5 m grid cells and computed the number of recruits of each species in each grid cell.

ESTIMATION OF LIGHT AVAILABILITY

Yearly canopy censuses were conducted from 1983 to 1996, except for 1994. We use the canopy census data to produce an index of the amount of light reaching any point in the forest. The canopy census includes a record of the height at which vegetation intersected vertical lines above points on a square, 5-m grid across the 50 ha. Presence versus absence of vegetation was recorded in six height intervals, 0–2, 2–5, 5–10, 10–20, 20–30 and ≥ 30 m, with the assistance of an ocular range finder. A single census of all 20 301 points on the grid was done annually.

If vegetation was present in a height interval, we assume it casts shade below exactly as would a flat circle of diameter 5 m placed at the vertical midpoint of the cell (for the topmost height category, we assumed the circle was at 35 m). For any point below, the vegetation obscures a section of the sky. Using trigonometry, we calculate the angle of sky so obscured by vegetation at position C , denoted α_c (see Appendix S1 in Supporting Information). The total amount of sky obscured at any point P is the sum of α_c over all cells C that are near enough to cast shade; we assume that points with a horizontal distance of ≤ 20 m from C are near enough. These assumptions mean that each cell of vegetation reduces light, even if there is another cell above it: the top layer removes a certain quantity of light, and the next layer below removes more. This approach seems reasonable, in that several layers of leaves directly above a point reduce light to a level lower than a single layer would. The assumption that the presence of vegetation can be modelled as a horizontal circle of diameter 5 m is arbitrary, and we acknowledge that this is only a crude way of converting canopy density above a point to light reaching the point. By using many surrounding points and many vertical layers (cf. Connell, Lowman & Noble 1997), we suggest that this method may be some-

what better at estimating light than methods based on only the tallest nearby trees (Lieberman *et al.* 1995) or the top layer of vegetation (Welden *et al.* 1991).

We calculated the shading index for each point of the plot's 5-m grid at height $z = 2$ m because this is the approximate height of recruits at 1 cm d.b.h. For each grid cell, we averaged the shading index s at its four corners and for all years of a census interval, i.e. 6 years for the first interval (1985–1990) and 5 years for the second (1990–1995; no canopy census was done in 1994).

To convert the estimates of mean shading to an estimate of light availability at each grid cell and in each census interval, we made use of 396 direct measurements of relative irradiance by Wirth, Weber & Ryel (2001), who provide a probability distribution of irradiance l at 1 m height in December 1993 at BCI. We matched their distribution of irradiance with our probability distribution of canopy shading, s . We assumed the 5th, 25th, 50th, 75th and 95th percentiles of the two distributions match, allowing a simple conversion of each value s to l by using the regression line through the percentiles (Appendix S1). Using this conversion, we found that average light availability in the understorey was 2% in both census intervals.

HIERARCHICAL BAYESIAN MODEL

We used a hierarchical Bayesian framework to assess the light response of recruitment across the tree and shrub species at BCI. Our model involves a data model, a process model and parameter models (Fig. 1).

Data enter our model as observed recruit numbers O_{ij} of each species (i) in each 5×5 m grid cell (j). We assume that O_{ij} varies around the predicted R_{ij} following a negative binomial distribution (data model or likelihood)

$$O_{ij} \sim \text{NegBinomial}(R_{ij}, c_i).$$

Recruit distributions are not only affected by the light conditions but also by the distribution of seed trees, spatially heterogeneous seed dispersal by animals or spatially heterogeneous soil conditions (e.g. Uriarte *et al.* 2005; Comita, Condit & Hubbell 2007). The negative binomial incorporates the resulting spatial autocorrelation of recruit numbers at < 5 m into the clumping parameter c_i . Autocorrelation of recruitment at larger scales (> 5 m) was negligible: autocorrelation for adjacent grid cells was < 0.2 for 252 of the 263 species we studied,

and there was no correlation in recruitment across cells separated by 20 m or more.

We compared seven functional relationships to describe the dependence of the number of recruits on light availability: constant, linear, semi-log, log–log, sigmoid and Michaelis–Menten. For each model, we judged the fit to data using the Akaike Information Criterion (AIC; Akaike 1974). When all 263 species were tested, the log–log relationship was selected as the best model for the highest proportion of species (excluding the constant model). For the species, for which the log–log relationship was not the best model, the best model was significantly better than the log–log relationship (i.e. AIC difference > 2) only in 36 cases.

Thus, we use a power function (log–log relationship) to model the light effect on recruitment (process model). The log of the predicted number of recruits R_{ij} of each species (i) in each 5×5 m grid cell (j) given light availability l_j is

$$\log_{10}(R_{ij}) = a_i + b_i \log_{10}(l_j).$$

In this model, a_i represents a measure of the mean log of the number of recruits to be expected, and b_i measures the strength of the light response of species i . For values of $b < 0$, the number of recruits decreases with increasing light. For $0 < b < 1$, the number of recruits increases in a decelerating way with increasing light; for $b = 1$, it is a linear relationship; and for $b > 1$, the number of recruits increases in an accelerating way with light (Fig. 2).

We then describe the variation of a and b across the community (parameter model). We used normal distributions for both parameters,

$$a_i \sim \text{Normal}(\mu_a, \sigma_a),$$

$$b_i \sim \text{Normal}(\mu_b, \sigma_b),$$

with hyperparameters μ_a and μ_b describing the community-wide means of a and b , and σ_a and σ_b measuring the between-species variation of both characteristics. Model runs without hyperdistributions confirmed that both variables were normally distributed across the community. These runs also revealed that for species with > 100 recruits, parameter estimates were largely determined by the data, whereas for species with < 100 recruits, parameter estimates for b were 'pulled' towards the community-wide mean.

As we did not have prior knowledge of the parameters, we used non-informative flat priors for all parameters:




Fig. 1. Graphical representation of the hierarchical model.




Fig. 2. Illustration of the three types of light response of recruitment: negative ($b < 0$), decelerating ($0 < b < 1$) and accelerating ($b > 1$). The number of recruits at average light conditions is 0.001 in all three cases ($a = -3$).

$$\begin{aligned} c_i &\sim \text{Uniform}(0, 100), \\ \mu_a &\sim \text{Uniform}(-10, 1), \\ \mu_b &\sim \text{Uniform}(-5, 5), \\ \sigma_a, \sigma_b &\sim \text{Uniform}(0, 2). \end{aligned}$$

We restricted c to values < 100 because for $c > 10$, the negative binomial already approximates a spatially homogeneous Poisson distribution (Bolker 2008). Altogether, the joint posterior distribution for the full model is:

$$\begin{aligned} &P(R, a, b, c, \mu_a, \sigma_a, \mu_b, \sigma_b | O, l) \\ &\propto \prod_{i=1}^{spp} \prod_{j=1}^{17644} \text{NegBinomial}(O_{ij} | R_{ij}, c_i) \\ &\times \prod_{i=1}^{spp} \text{Normal}(a_i | \mu_a, \sigma_a) \times \prod_{i=1}^{spp} \text{Normal}(b_i | \mu_b, \sigma_b) \times \prod_{i=1}^{spp} \text{Unif}(c_i | 0, 100) \\ &\times \text{Unif}(\mu_a | -10, 1) \times \text{Unif}(\mu_b | -5, 5) \times \text{Unif}(\sigma_a | 0, 2) \times \text{Unif}(\sigma_b | 0, 2), \end{aligned}$$

with spp being the number of species with recruits in the respective census interval. The last three lines denote the likelihood, the priors and the hyperpriors, respectively.

MODEL IMPLEMENTATION AND DIAGNOSTICS

Posterior distributions of the parameters were obtained using a Markov chain Monte Carlo method that is a hybrid of the Metropolis–Hastings algorithm and the Gibbs sampler (Gelman *et al.* 1995; Condit *et al.* 2006). Parameter values are sequentially updated as in the Gibbs sampler, but acceptance depends on the likelihood ratios as in the Metropolis–Hastings algorithm (Muller-Landau *et al.* 2008). The proposal distribution is a normal distribution centred on the current value of the given parameter. The step width for each parameter, i.e. the standard deviation of the proposal distribution, is constantly adjusted during the burn-in period in such a way that the acceptance rate is kept around 0.25 (Gelman *et al.* 1995).

To speed up the convergence of the Gibbs sampler, we weakened the correlation of a and b by centring the light data at average \log_{10} light (\bar{l})

$$\log_{10}(R_{ij}) = a_i + b_i(\log_{10}(l_j) - \bar{l}).$$

Thus, a_i measures the mean log of the predicted number of recruits at average light conditions. Mean \log_{10} of l was -1.7 for the period 1985–1990 and -1.8 for 1990–1995. We used an average value of

$\bar{l} = -1.7$, corresponding to a relative irradiance of 2%, for the analysis of both periods to keep parameter estimates comparable.

We monitored convergence by running two chains with different initial values and used Gelman and Rubin's convergence diagnostics and a value of 1.1 to detect the convergence (Gelman & Rubin 1992; Gelman *et al.* 1995). Convergence required 500–5100 iterations. We used a burn-in period of 7000 iterations and additional 10 000 iterations were used for analysis. We computed the posterior parameter distributions for observed recruit numbers and light availability estimations for two intervals (1985–1990, 1990–1995). All analyses were carried out using the software package R version 2.5.0 (R Development Core Team 2007). The R-code is available in Appendix S2.

ANALYSIS OF RESULTS

From the posterior distributions, we computed the mean and 95% credible intervals (CI) of all parameters for the two periods (1985–1990, 1990–1995). We define maximum recruitment as recruitment at high light conditions, where high light is 20% relative irradiance, since 20% is close to the brightest the forest understorey ever gets. For each species, we calculated the light level at which the recruitment falls to 50% (*Light50*) of maximum recruitment. To compare model fits with data, we calculated the mean number of observed recruits in ten light intervals corresponding to 10th percentiles of light estimates across the 50-ha plot. Total abundance of each species across 50 ha in 1990 (i.e. number of individuals ≥ 1 cm d.b.h.) was used in comparing species. All BCI inventory data, including recruitment, can be downloaded from <http://www.ctfs.si.edu>.


Results

COMMUNITY-LEVEL PATTERN OF LIGHT RESPONSE

Mean posterior estimates of the light response of recruitment (b) ranged from -0.6 to 3.3 , with most species falling between 0 and 1, i.e. these species take advantage of higher light levels with a decelerating response of recruitment to light (Fig. 3; Table S1). When mean posterior estimates of b were compared between census intervals, 101 species (38%) showed a decelerating response of recruitment with increasing light availability ($0 < b < 1$) in both census intervals, 54 species (21%) showed an accelerating response ($b > 1$), and recruitment of nine species (3%) decreased with light availability ($b < 0$). For the remaining 99 species (38%), b indicated different types of light response for the two census intervals or species did not have recruits in both intervals.

For 97 species, the 95% CI of b did not include zero in either census interval. There were only three species with a negative b and a CI that did not include zero: *Xylopia macrantha* in both census intervals, *Trichilia tuberculata* only in the first interval, and *Drypetes standleyi* and *Licania platytypus* only in the second interval. There were 18 species with $b > 1$ and a 95% CI > 1 in both intervals, i.e. an accelerating response of recruitment to light availability. These included many species of the common pioneer genera, e.g. *Cecropia*, *Croton*, *Miconia*, *Palicourea*, *Spondias*, *Zanthoxylum*. Posterior estimates of the community-wide mean of b (μ_b) were 0.80 for 1985–1990 and 0.85 for 1990–1995, and the standard deviation (σ_b) was 0.76 and 0.72, for the two census intervals, respectively.

Fig. 3. Mean posterior estimates of the light dependence of recruitment (b) for the tree species at BCI, Panama, for the two census intervals (1985–1990 and 1990–1995).



OTHER PARAMETERS

Mean posterior estimates of the log of the mean number of recruits at average light conditions (a) ranged from -4.3 to -0.88 (i.e. 0.02–53 recruits per hectare; Fig. 4 and Table S1). The minimum roughly corresponded to one individual in the sampling area, while the maximum corresponded to about 0.1 individuals per 5×5 m grid cell. Posterior estimates of the community-wide mean of a (μ_a) were -3.017 and -3.162 , and posterior estimates of the standard deviation (σ_a) were 0.88 and 0.84, for the two census intervals, respectively. The distribution of a across the community was skewed to the right.

The distribution of the clustering parameter of the negative binomial was bimodal (Fig. 4; Table S1). Thirty species showed pronounced clustering ($c < 1$) in both census intervals, 32 only in the first and nine only in the second census

interval, while for the majority of the species either spatial clustering did not occur or could not be detected due to the small number of recruits ($c > 10$). The correlation between b and c is -0.13 in the first census interval and -0.18 in the second, indicating that recruits of more light-demanding species tend to be slightly more clustered than recruits of shade-tolerant species.

CONSISTENCY BETWEEN CENSUS INTERVALS

Comparing mean posterior estimates for the two census intervals showed that the log of the mean number of recruits (a) tended to be slightly higher in the first census interval 1985–1990 than in the second census interval 1990–1995 (Fig. 5a). However, parameter estimates were consistent over the two intervals. Likewise, the estimates of the strength of the light




Fig. 4. Mean posterior estimates of the \log_{10} of the mean number of recruits at average light conditions (a) and the clustering parameter of the negative binomial distribution (c) for the tree species at BCI, Panama, for the two census intervals (1985–1990 and 1990–1995).

response of recruitment (b) were consistent over the two analysed time intervals (Fig. 5b).

OBSERVED VERSUS PREDICTED RECRUITMENT RATES

Mean posterior parameter estimates predicted the mean number of recruits under different light conditions. Model fits are shown for two species with positive (*Trichilia pallida*) and negative (*Trichilia tuberculata*) response to light (Fig. 6). Model fits for the remaining species are available in Fig. S1.

UNCERTAINTY AND SAMPLE SIZE

The uncertainty of parameter estimates decreased rapidly with the number of recruits (Fig. 7). When recruit numbers were low, the 95% CI of a spanned up to an order of magnitude (Fig. 7a). Likewise, CIs for b ranged over more than one type of light response, i.e. $b < 0$, $0 < b < 1$, $b > 1$, when few recruits were available for parameter estimation (Fig. 7b). However, for 46% of the species with > 20 recruits, parameter estimates significantly determined the type of light response (95% CI of b neither including 0 nor 1), as opposed to 7% of the species with < 20 recruits. When the number of recruits increased to > 50 (100), this proportion increased to 59%

(66%), as opposed to 10% (15%) for species with < 50 (100) recruits. Below ten recruits, spatial aggregation could not be detected, but also for larger numbers of recruits, the clustering parameter c often did not contain information and varied over the entire range of allowed values (0–100) (Fig. 7c).

LIGHT RESPONSE AND ABUNDANCE

Across the community, the strength of the light response of recruitment (b) decreased slightly with abundance. In a linear regression, the effect of \log_{10} of abundance on b was significant ($P < 0.0001$, $n = 233$, $r^2 = 0.07$, intercept = 1.19, slope = -0.155). While b varied over a broad range for species with intermediate abundance, the most abundant species had a b close to zero, i.e. showed hardly any response of recruit numbers to light availability. For rare species (< 50 recruits), however, b was largely determined by the community-wide mean and the relationship is not informative for them.

LIGHT REQUIREMENTS FOR RECRUITMENT

The light intensity where recruitment fell to 50% of maximum recruitment ($Light_{50}$) was heavily skewed to the right (Fig. 8). Fifty-six species (22% of the total) in the first interval had




Fig. 5. Posterior means of parameters of the light dependence of recruitment for the tree species at BCI, Panama, for the two census intervals (1985–1990, 1990–1995). Parameters are (a) \log_{10} of the mean number of recruits at average light conditions a , and (b) the light effect b . The line represents the 1:1 relationship.




Fig. 6. Model fit for the predicted number of recruits at different light conditions vs. the observed number of recruits for two selected species. Mean number of recruits in intervals of tenth percentiles of light in the 5×5 m grid cells (large dots), observed number of recruits in the 5×5 m grid cells (small dots), predicted number of recruits using posterior means for the second interval 1990–1995 (line). Grid cells without recruits were assigned a small number ($1.5e^{-5}$).




Fig. 7. Uncertainty of estimates (width of the 95% credible interval) of (a) the \log_{10} of the mean number of recruits at average light conditions a , (b) the light dependence of recruitment b , and (c) the clustering parameter of the negative binomial distribution c as a function of the number of sampled recruits. Uncertainty estimates are given for 233 tree species at BCI, Panama, for the second census interval (1990–1995).

$Light_{50} < 2\%$, the understorey light average, i.e. they recruited well at low light. At the opposite end, 81 species (32% of the total) in the first interval had $Light_{50} > 10\%$ and thus only recruited well at high light. The second census interval had similar results, with 35 species recruiting well at average light and 84 requiring high light. The median $Light_{50}$ in the two census intervals was 7.7% and 8.5%.

Discussion

Our results broadly support the notion that light requirements for recruitment are distributed along a continuum (Denslow 1987; Wright *et al.* 2003) and that species belong to broad regeneration guilds rather than partitioning the light gradient at a fine scale (Hubbell & Foster 1986). The distribution of the light response of recruitment (b) had a strong central mode, with many species showing a nearly linear light response, suggesting that the majority of species share intermediate light requirements for recruitment (Wright *et al.* 2003). Using the hyperparameters of the posterior distribution of light dependence across the community, recruitment of 13% of the species is expected to decrease with increasing light availability, while recruitment of 87% of the species benefits from higher light,

46% in a decelerating manner and 41% in an accelerating manner.

Recruitment of only 20% of the species reached 50% or more of their maximum recruitment at light levels lower than the average in the forest understorey at BCI suggesting that relatively few species recruit well in the deepest forest shade. Thus, the ability to regenerate at low-light conditions may not be as widespread among tree species at BCI as we thought (Hubbell & Foster 1986; Welden *et al.* 1991; Condit, Hubbell & Foster 1996). Rather, nearly all species regenerate better in increasing light, and they are spread along a continuum from modest response to a rather strict requirement for high light. Species specialized to recruit in high-light conditions ($b > 1$) are more common (41%) than expected from the rarity of large gaps (Hubbell & Foster 1986; Hubbell *et al.* 1999).

Our results are consistent with the findings of a community-wide analysis of sapling distributions with respect to light in a temperate rain forest in Chile (Lusk, Chazdon & Hofmann 2006). In that study, half the species were randomly distributed with respect to canopy openness, whilst the remainder was non-randomly distributed. The majority of tree species were also found to have intermediate shade tolerance, and niche overlap was not smaller than expected.

There are several caveats we must consider. The basis of our analysis is the distribution of saplings with respect to the estimated light levels integrated over 6 years during which the recruit surpassed the 1 cm d.b.h. threshold. Based on mean growth at 1 cm d.b.h. ($< 1 \text{ mm year}^{-1}$), we estimate the average age of individuals reaching the 1 cm d.b.h. threshold to be 10–20 years. Therefore, the establishment of seedlings and subsequent growth and mortality may have occurred under light conditions different from those at the time of the study (Denslow 1987; Kobe 1999). However, we believe that our approach gives reasonable estimates of light requirements for recruitment not only for fast-growing species, but also for slower-growing species, due to the temporal correlation of light availability caused by the slow dynamics of the canopy cover (the correlation between the light estimates for both census intervals was 0.76).




Fig. 8. Estimates of the light level at which recruitment falls to 50% of recruitment under high light conditions (i.e. at 20% light, $Light_{50}$) for the tree species at BCI, Panama. Estimates are based on the means of posterior means of the species' parameters from both census intervals (1985–1990, 1990–1995).

Recruitment success in terms of our study integrates both growth and survival of plants during the 5 years of the census interval. Growth and mortality of seedlings and saplings are likely to vary with light availability (e.g. Montgomery & Chazdon 2002; Iriarte & Chazdon 2005; Aiba & Nakashizuka 2007). Since recruits only enter our census at 1 cm d.b.h., we cannot separate the impact of light on growth from the impact on survival.

Another concern is that our calculated light availability obviously is only a rough index of the light environment in the forest, unable to capture spatial variation at scales < 5 m horizontally or vertically. Measuring light directly at every recruit, however, would involve prohibitive time and labour. Until laser-mapping data of the forest are available, the method we propose offers an objective and straightforward measure of how much vegetation is blocking the sky above any tree of any height. The consistency of the derived light response between the two census intervals indicates that our light index contains information relevant to the species' performance. Interestingly, our light estimation procedure is similar to the index that best described gap light index obtained by hemispherical photographs in a temperate forest in Japan (Takashima, Kume & Yoshida 2006). The latter estimates also accounted for shading from topography, but we ignored this at BCI due to the relatively flat topography.

The power function does not capture the saturation of recruit numbers at very high light levels for some pioneer species with $b > 1$ (e.g. *Cecropia insignis*, *Croton billbergianus*, *Miconia argentea*, *Palicourea guianensis*, *Spondias radlkoferi*; Fig. S1). Thus for these species, we overestimate the number of recruits at very high light and predicted recruit numbers beyond light levels found in forest gaps (c. 20%) should be used with caution.

A final caveat is the low number of individuals per area for the many rare species in such a diverse forest (Pitman *et al.* 1999). Hierarchical Bayesian methods explicitly account for this problem by superimposing a form of variation across the community and by providing direct measures of uncertainty associated with parameter estimates (e.g. Clark 2005; Condit *et al.* 2006). We could significantly determine the type of light response (i.e. negative, decelerating or accelerating) for nearly half of the species (44%) with at least 20 recruits. For 167 (75%) of the 224 species with recruits in both census intervals, light response was consistent across the two intervals, i.e. mean posterior estimates of light dependence indicated the same type of light response (negative, decelerating or accelerating). However, applying the rather strong criterion, that the width of the 95% CI of b be < 0.5, about 100 recruits were necessary. This is certainly a large number for rare species. The main point of the Bayesian approach, however, is that the response of rare species need not be known to estimate the community-wide variation.

Conclusions

Light response of recruitment is only one aspect of species' performance under heterogeneous environmental conditions.

To arrive at a more complete understanding of species' life-history traits, differential response of growth and survival to light conditions have to be analysed for different ontogenetic stages (Clark & Clark 1992; Poorter *et al.* 2005). Such analyses will allow extending existing species classifications into functional groups on the basis of light (or other resource) dependence of all vital rates in several life stages (Condit, Hubbell & Foster 1996). Ultimately, only mechanistic models accounting for species differences will allow investigations of the consequences of species' similarities or differences for coexistence and to determine the relative importance of niche partitioning versus chance events (Grubb 1977; Brokaw & Busing 2000; Wright 2002).

Acknowledgements

We thank Liza Comita and Inés Ibáñez for advice on hierarchical Bayesian modelling, Helene Muller-Landau for providing the step width-adjusting routine of the Metropolis-Hastings algorithm, and Helene Muller-Landau and Robin Chazdon for valuable comments on the manuscript. N.R. was supported by a research grant from the Center for Tropical Forest Science, and R.C. by a fellowship from the National Center for Ecological Analysis and Synthesis (Santa Barbara, CA, USA). The BCI plot has been made possible through the support of the U.S. National Science Foundation, the John D. and Catherine D. McArthur Foundation and the Smithsonian Tropical Research Institute. We thank the dozens of field assistants and botanists who have collected data on trees and carried out the canopy censuses in the BCI plot over the past 25 years.

References

- Aiba, M. & Nakashizuka, T. (2007) Variation in juvenile survival and related physiological traits among dipterocarp species co-existing in a Bornean forest. *Journal of Vegetation Science*, **18**, 379–388.
- Akaike, H. (1974) A new look at the statistical model identification. *IEEE Transactions on Automatic Control*, **19**, 716–723.
- Barker, M.G., Press, M.C. & Brown, N.D. (1997) Photosynthetic characteristics of dipterocarp seedlings in three tropical rain forest light environments: a basis for niche partitioning? *Oecologia*, **112**, 453–463.
- Bolker, B. (2008) *Ecological Models and Data in R*. Princeton University Press, Princeton.
- Brokaw, N.V.L. (1985) Gap-phase regeneration in a tropical forest. *Ecology*, **66**, 682–687.
- Brokaw, N. & Busing, R.T. (2000) Niche versus chance and tree diversity in forest gaps. *Trends in Ecology & Evolution*, **15**, 183–188.
- Brown, N.D. & Whitmore, T.C. (1992) Do dipterocarp seedlings really partition tropical rain forest gaps? *Philosophical Transactions of the Royal Society London B*, **335**, 369–378.
- Clark, J.S. (2005) Why environmental scientists are becoming Bayesians. *Ecology Letters*, **8**, 2–14.
- Clark, D.A. & Clark, D.B. (1992) Life-history diversity of canopy and emergent trees in a neotropical rain forest. *Ecological Monographs*, **62**, 315–344.
- Comita, L.S., Condit, R. & Hubbell, S.P. (2007) Developmental changes in habitat associations of tropical trees. *Journal of Ecology*, **95**, 482–492.
- Condit, R. (1998) *Tropical Forest Census Plots*. Springer-Verlag, Berlin.
- Condit, R., Hubbell, S.P. & Foster, R.B. (1996) Assessing the response of plant functional types to climatic change in tropical forests. *Journal of Vegetation Science*, **7**, 405–416.
- Condit, R., Ashton, P.S., Bunyavejchewin, S., Dattaraja, H.S., Davies, S., Esufali, S. *et al.* (2006) The importance of demographic niches to tree diversity. *Science*, **313**, 98–101.
- Connell, J.H., Lowman, M.D. & Noble, I.R. (1997) Subcanopy gaps in temperate and tropical forests. *Australian Journal of Ecology*, **22**, 163–168.
- Croat, T.B. (1978) *Flora of Barro Colorado Island*. Stanford University Press, Stanford, CA.
- Denslow, J.S. (1980) Gap partitioning among tropical rainforest trees. *Biotropica*, **12**, 47–55.
- Denslow, J.S. (1987) Tropical rainforest gaps and tree species diversity. *Annual Review of Ecology and Systematics*, **18**, 431–451.

- Dupuy, J.M. & Chazdon, R.L. (2006) Effects of vegetation cover on seedling and sapling dynamics in secondary tropical wet forests in Costa Rica. *Journal of Tropical Ecology*, **22**, 65–76.
- Ellison, A.M. (2004) Bayesian inference in ecology. *Ecology Letters*, **7**, 509–520.
- Foster, R.B. & Brokaw, N.V. (1982) Structure and history of the vegetation of Barro Colorado Island. *The Ecology of a Tropical Forest: Seasonal Rhythms and Long-term Changes* (eds E.G. Leigh Jr, A.S. Rand & D.M. Windsor), pp. 67–81. Smithsonian Institution, Washington, DC.
- Gelman, A. & Rubin, D.B. (1992) Inference from iterative simulation using multiple sequences. *Statistical Science*, **7**, 457–511.
- Gelman, A., Carlin, J.B., Stern, H.S. & Rubin, D.B. (1995) *Bayesian Data Analysis*. Chapman & Hall, London.
- Grubb, P.J. (1977) The maintenance of species richness in plant communities: the importance of the regeneration niche. *Biological Reviews of the Cambridge Philosophical Society*, **52**, 107–145.
- Hartshorn, G.S. (1980) Neotropical forest dynamics. *Biotropica*, **12**, 23–30.
- Hubbell, S.P. & Foster, R.B. (1983) Diversity of canopy trees in a neotropical forest and implications for conservation. *Tropical Rain Forest: Ecology and Management* (eds S.L. Sutton, T.C. Whitmore & A.C. Chadwick), pp. 25–41. Blackwell Scientific, Oxford.
- Hubbell, S.P. & Foster, R.B. (1986) Biology, chance and history and the structure of tropical rain forest tree communities. *Community Ecology* (eds J.L. Diamond & T.J. Case), pp. 314–329. Harper & Row, New York.
- Hubbell, S.P., Foster, R.B., O'Brien, S.T., Harms, K.E., Condit, R., Wechsler, B., Wright, S.J. & de Lao, S.L. (1999) Light-gap disturbances, recruitment limitation, and tree diversity in a neotropical forest. *Science*, **283**, 554–557.
- Iriarte, V.B.S. & Chazdon, R.L. (2005) Light-dependent seedling survival and growth of four tree species in Costa Rican second-growth rain forests. *Journal of Tropical Ecology*, **21**, 383–395.
- Kobe, R.K. (1999) Light gradient partitioning among tropical tree species through differential seedling mortality and growth. *Ecology*, **80**, 187–201.
- Leigh, E.G.J. (1999) *Tropical Forest Ecology: A View From Barro Colorado Island*. Oxford University Press, Oxford.
- Leigh, E.G.J., Rand, S.A. & Windsor, D.M. (1982) *The Ecology of a Tropical Forest: Seasonal Rhythms and Long-term Changes*. Smithsonian Institution Press, Washington, DC.
- Lieberman, M., Lieberman, D. & Peralta, R. (1989) Forests are not just Swiss cheese: canopy stereogeometry of non-gaps in tropical forests. *Ecology*, **70**, 550–552.
- Lieberman, M., Lieberman, D., Peralta, R. & Hartshorn, G.S. (1995) Canopy closure and the distribution of tropical forest tree species at La Selva, Costa Rica. *Journal of Tropical Ecology*, **11**, 161–178.
- Lusk, C.H., Chazdon, R.L. & Hofmann, G. (2006) A bounded null model explains juvenile tree community structure along light availability gradients in a temperate rain forest. *Oikos*, **112**, 131–137.
- Montgomery, R.A. & Chazdon, R.L. (2002) Light gradient partitioning by tropical tree seedlings in the absence of canopy gaps. *Oecologia*, **131**, 165–174.
- Muller-Landau, H.C., Wright, S.J., Calderon, O., Condit, R. & Hubbell, S.P. (2008) Interspecific variation in primary seed dispersal in a tropical forest. *Journal of Ecology*, **96**, 653–667.
- Pitman, N.C.A., Terborgh, J., Silman, M.R. & Núñez, P. (1999) Tree species distributions in an upper Amazonian forest. *Ecology*, **80**, 2651–2662.
- Poorter, L. & Arets, E.J.M.M. (2003) Light environment and tree strategies in a Bolivian tropical moist forest: an evaluation of the light partitioning hypothesis. *Plant Ecology*, **166**, 295–306.
- Poorter, L., Bongers, F., Sterck, F.J. & Woll, H. (2005) Beyond the regeneration phase: differentiation of height-light trajectories among tropical tree species. *Journal of Ecology*, **93**, 256–267.
- Queenborough, S.A., Burslem, D.F.R.P., Garwood, N.C. & Valencia, R. (2007) Habitat niche partitioning by 16 species of Myristicaceae in Amazonian Ecuador. *Plant Ecology*, **192**, 193–207.
- R Development Core Team (2007) *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>.
- Rees, M., Condit, R., Crawley, M., Pacala, S. & Tilman, D. (2001) Long-term studies of vegetation dynamics. *Science*, **293**, 650–655.
- Ricklefs, R.E. (1977) Environmental heterogeneity and plant species diversity: a hypothesis. *American Naturalist*, **111**, 376–381.
- Silvertown, J. (2004) Plant coexistence and the niche. *Trends in Ecology & Evolution*, **19**, 605–611.
- Svenning, J.C. (2000) Small canopy gaps influence plant distributions in the rain forest understory. *Biotropica*, **32**, 252–261.
- Swaine, M.D. & Whitmore, T.C. (1988) On the definition of ecological species groups in tropical rain forests. *Vegetatio*, **75**, 81–86.
- Takashima, A., Kume, A. & Yoshida, S. (2006) Methods for estimating understory light conditions using crown projection maps and topographic data. *Ecological Research*, **21**, 560–569.
- Tilman, D. (1982) *Resource Competition and Community Structure*. Princeton University Press, Princeton.
- Tilman, D. (2004) Niche tradeoffs, neutrality, and community structure: a stochastic theory of resource competition, invasion, and community assembly. *Proceedings of the National Academy of Sciences of the United States of America*, **101**, 10854–10861.
- Uriarte, M., Canham, C.D., Thompson, J., Zimmerman, J.K. & Brokaw, N. (2005) Seedling recruitment in a hurricane-driven tropical forest: light limitation, density-dependence and the spatial distribution of parent trees. *Journal of Ecology*, **93**, 291–304.
- Van der Meer, P.J., Sterck, F.J. & Bongers, F. (1998) Tree seedling performance in canopy gaps in a tropical rain forest at Nouragues, French Guiana. *Journal of Tropical Ecology*, **14**, 119–137.
- Welden, C.W., Hewett, S.W., Hubbell, S.P. & Foster, R.B. (1991) Sapling survival, growth, and recruitment: relationship to canopy height in a neotropical forest. *Ecology*, **72**, 35–50.
- Whitmore, T.C. (1996) A review of some aspects of tropical rain forest seedling ecology with suggestions for further enquiry. *The Ecology of Tropical Forest Tree Seedlings*. Man and the Biosphere Series 17 (ed. M.D. Swaine), pp. 3–39. UNESCO, Paris.
- Windsor, D.M. (1990) *Climate and Moisture Variability in a Tropical Forest: Long-Term Records from Barro Colorado Island, Panama*. Smithsonian Institution Press, Washington, DC.
- Windsor, D.M., Rand, A.S. & Rand, W.M. (1990) Características de la Precipitación de la Isla de Barro Colorado. *Ecología de un Bosque Tropical: Ciclos Estacionales y Cambios a Largo Plazo* (eds A.S. Rand, D.M. Windsor & E.G. Leigh Jr), pp. 53–71. Smithsonian Tropical Research Institute, Balboa.
- Wirth, R., Weber, B. & Ryel, R.J. (2001) Spatial and temporal variability of canopy structure in a tropical moist forest. *Acta Oecologica*, **22**, 235–244.
- Wright, S.J. (2002) Plant diversity in tropical forests: a review of mechanisms of species coexistence. *Oecologia*, **130**, 1–14.
- Wright, S.J., Muller-Landau, H.C., Condit, R. & Hubbell, S.P. (2003) Gap-dependent recruitment, realized vital rates, and size distributions of tropical trees. *Ecology*, **84**, 3174–3185.

Received 22 January 2009; accepted 30 June 2009
 Handling Editor: Matthew Turnbull

Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1. Calculation of the shading index using canopy density and conversion of the shading index to relative irradiance.

Appendix S2. R code to generate posterior distributions of the parameters of the hierarchical Bayesian model.

Figure S1. Model fits for the predicted number of recruits at different light conditions versus the observed number of recruits for the woody species at Barro Colorado Island, Panama.

Table S1. Posterior means, lower and upper limits of the credible interval (CI⁻, CI⁺) of the parameters of the hierarchical Bayesian model and the light level at which recruitment falls to 50% (*Light*₅₀) of recruitment at 20% light for the woody species at Barro Colorado Island, Panama.

Please note: Wiley-Blackwell is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

Appendix S1

Calculation of the shading index using canopy density and conversion of the shading index to relative irradiance

1. Calculation of the shading index using canopy density

Consider an arbitrary point P whose three-dimensional coordinates in the forest are x, y, z , and consider a cell of vegetation C whose midpoint is at position x_c, y_c, z_c (Fig. 1). Call the horizontal distance between the two points d_c , calculated from x and y coordinates, and the vertical distance $h_c = z_c - z$. Assume the vegetation in the cell is a horizontal circle of radius r . Then the formula for the angle α of sky obscured by the layer of vegetation is

$$\alpha_c = \arctan\left(\frac{d_c + r}{h_c}\right) - \arctan\left(\frac{d_c - r}{h_c}\right) \text{ if } d_c \leq 20 \text{ m}$$


$$\alpha_c = 0 \text{ if } d_c > 20 \text{ m,}$$

as we assume vegetation further than 20 m away (horizontal distance) have no impact on the light reaching a point. We estimate the total shading at point P as

$$s_p = \sum_c \alpha_c,$$

where the summation means over all cells c in the 50-ha plot.

Figure 1: Diagram of the calculation converting vegetation density to shading. Point P at horizontal coordinates x, y , and height z , is where an estimate of shading is needed. The vegetation cell (Veg) has midpoint (vertical and horizontal) at point C . We assume the amount of light removed (*i.e.*, shading) by the vegetation cell is proportional to the angle α_c .



2. Conversion of the shading index to relative irradiance

We assumed a linear relationship between the log of relative irradiance l (data from Wirth *et al.* 2001) and the shading index s (for the period 1990–95 to be temporally closer to the light measurements which were carried out in 1993) and performed a quadratic regression through 5th, 25th, 50th, 75th and 95th percentiles of both distributions, assuming that the lowest estimate of s for any tree individual (a *Prioria copaifera* tree with a dbh > 1.2 m) in any census interval corresponds to 100% light (Fig. 2). Thus, $l = e^{-0.01351-0.08043 \cdot s-0.00315 \cdot s^2}$. Distributions of estimated light are given in Fig. 3. The spatial distribution of recruits closely follows the distribution of estimated light across the 50-ha plot (Fig. 4).




Figure 2: Conversion of shade estimates to relative irradiance.





Figure 3: Distribution of estimated light at 2 m in the center of 5 m × 5 m quadrats across the 50-ha plot on Barro Colorado Island, Panama.

Figure 4: Spatial distribution of recruits (1985–1990; red dots) with respect to estimated light at 2 m height across the 50-ha plot. Darker shades of grey indicate higher light availability.



References

Wirth, R., Weber, B. & Ryel, R.J. (2001) Spatial and temporal variability of canopy structure in a tropical moist forest. *Acta Oecologica*, **22**, 235–244.

Appendix S2

R code to generate posterior distributions of the parameters of the hierarchical Bayesian model

```
# Functions for fitting a power function to the number of recruits per species given available light, hyperdistributions of  
# the log of the mean number of recruits at average light availability (a) and light-dependence (b), and the clumping  
# parameter of the negative binomial distribution (k) using a hybrid of the Gibbs sampler and the Metropolis-Hastings  
# algorithm. These functions should source in R2.5.0, and given a table of data in the correct format, will produce the  
# posterior distribution of all parameters.
```

```
# The matrix of data has one row per species and columns with the number of recruits in each 5 m × 5 m quadrat.  
# normlight is a vector with centered light data: log10 of relative irradiance in each 5 m × 5 m quadrat – mean(log10 of  
# relative irradiance all 5 m × 5 m quadrats).  
# spp is a character vector with species' acronyms
```

```
# The four hyperparameters are the mean (MuA) and SD (SdA) of the log of the mean number of recruits at average  
# light availability (a) and mean (MuB) and SD (SdB) of the light-dependence (b). The argument start.param sets their  
# starting values.
```

```
# start.scale sets the initial step size for the proposal distribution of the Metropolis-Hastings algorithm, the step size is  
# adjusted scale.burnin steps (burn-in), so that acceptance rate is kept around 0.25. The number of cycles run by the  
# Gibbs sampler is the argument steps, and the current state will be printed to the screen every showstep steps. The  
# argument whichuse defines the post burn-in phase; estimates based on the Gibbs sampler are drawn from those steps.
```

```
library(coda)
```

```
metrop.normLight.Gibbs=function(data,start.param=c(1,0),start.scale=c(0.1,0.01),scale.burnin=7000,steps=17000,  
  whichuse=7000:17000,showstep=100)
```

```
{  
  MuA1=SdA1=MuB1=SdB1=MuA2=SdA2=MuB2=SdB2=matrix(nrow=steps,ncol=3)  
  MuA1[1,1]=MuA2[1,1]=log10(sum(data[,,])/((dim(data)[1]*dim(data)[2])))  
  SdA1[1,1]=SdA2[1,1]=start.param[1]  
  MuB1[1,1]=MuB2[1,1]=start.param[2]  
  SdB1[1,1]=SdB2[1,1]=start.param[1]  
  MuA1[1,3]=SdA1[1,3]=MuB1[1,3]=SdB1[1,3]=MuA2[1,3]=SdA2[1,3]=MuB2[1,3]=SdB2[1,3]=start.scale[1]
```

```
  nospp=length(spp)  
  spa1=spb1=spk1=drawsa1=drawsb1=drawsk1=scalea1=scaleb1=scalek1=matrix(ncol=steps,nrow=nospp)  
  spa2=spb2=spk2=drawsa2=drawsb2=drawsk2=scalea2=scaleb2=scalek2=matrix(ncol=steps,nrow=nospp)  
  rownames(spa1)=rownames(spb1)=rownames(drawsa1)=rownames(drawsb1)=rownames(spk1)=rownames(drawsk1)  
  =rownames(scalea1)=rownames(scaleb1)=rownames(scalek1)=spp  
  rownames(spa2)=rownames(spb2)=rownames(drawsa2)=rownames(drawsb2)=rownames(spk2)=rownames(drawsk2)  
  =rownames(scalea2)=rownames(scaleb2)=rownames(scalek2)=spp
```

```
# chain 1 is initialized with average values  
spa1[,1]=log10(sum(data[,,])/((dim(data)[1]*dim(data)[2])))  
spb1[,1]=0  
spk1[,1]=10
```

```
# chain 2 is initialized with more extreme values  
spa2[,1]=-3  
spb2[,1]=2  
spk2[,1]=1
```

```
convergence = matrix(data=NA, ncol=3, nrow=nospp)
```

```
scalea1[,1] = scaleb1[,1] = scalea2[,1] = scaleb2[,1] = start.scale[2]
```

```
scalek1[,1] = scalek2[,1] = start.scale[1]
```

```
for(i in 2:steps)
```

```
{
```

```
  adjust=ifelse((i <= scale.burnin), T, F)
```

```
  MuA1[i,] = metrop1step(func=mu.aGibbs,start.param=MuA1[i-1,1],scale.param=MuA1[i-1,3],SdA=SdA1[i-1,1],  
    spa=spa1[i-1],adjust.scale=adjust)
```

```
  SdA1[i,] = metrop1step(func=sd.aGibbs,start.param=SdA1[i-1,1],scale.param=SdA1[i-1,3],MuA=MuA1[i,1],  
    spa=spa1[i-1], adjust.scale=adjust)
```

```
  MuB1[i,] = metrop1step(func=mu.bGibbs,start.param=MuB1[i-1,1],scale.param=MuB1[i-1,3],SdB=SdB1[i-1,1],  
    spb=spb1[i-1],adjust.scale=adjust)
```

```
  SdB1[i,] = metrop1step(func=sd.bGibbs,start.param=SdB1[i-1,1],scale.param=SdB1[i-1,3],MuB=MuB1[i,1],  
    spb=spb1[i-1],adjust.scale=adjust)
```

```
  MuA2[i,] = metrop1step(func=mu.aGibbs,start.param=MuA2[i-1,1],scale.param=MuA2[i-1,3],SdA=SdA2[i-1,1],  
    spa=spa2[i-1],adjust.scale=adjust)
```

```
  SdA2[i,]=metrop1step(func=sd.aGibbs,start.param=SdA2[i-1,1],scale.param=SdA2[i-1,3],MuA=MuA2[i,1],  
    spa=spa2[i-1],adjust.scale=adjust)
```

```
  MuB2[i,]=metrop1step(func=mu.bGibbs,start.param=MuB2[i-1,1],scale.param=MuB2[i-1,3],SdB=SdB2[i-1,1],  
    spb=spb2[i-1],adjust.scale=adjust)
```

```
  SdB2[i,]=metrop1step(func=sd.bGibbs,start.param=SdB2[i-1,1],scale.param=SdB2[i-1,3],MuB=MuB2[i,1],  
    spb=spb2[i-1],adjust.scale=adjust)
```

```
for(j in 1:nospp)
```

```
{
```

```
  nexta = metrop1step(func=spa.Gibbs, start.param=spa1[j,i-1], scale.param=scalea1[j,i-1],N=data[j,], spb=spb1[j,i-1],  
    spk=spk1[j,i-1],MuA=MuA1[i,1], SdA=SdA1[i,1], MuB=MuB1[i,1], SdB=SdB1[i,1],adjust.scale=adjust)
```

```
  spa1[j,i] = nexta[1]
```

```
  draws1[j,i] = nexta[2]
```

```
  scalea1[j,i] = nexta[3]
```

```
  nextb = metrop1step(func=spb.Gibbs, start.param=spb1[j,i-1], scale.param=scaleb1[j,i-1], N=data[j,], spa=spa1[j,i],  
    spk=spk1[j,i-1], MuA=MuA1[i,1], SdA=SdA1[i,1], MuB=MuB1[i,1], SdB=SdB1[i,1], adjust.scale=adjust)
```

```
  spb1[j,i] = nextb[1]
```

```
  draws1[j,i] = nextb[2]
```

```
  scaleb1[j,i] = nextb[3]
```

```
  nextk = metrop1step(func=spk.Gibbs, start.param=spk1[j,i-1], scale.param=scalek1[j,i-1], N=data[j,], spa=spa1[j,i],  
    spb=spb1[j,i], adjust.scale=adjust)
```

```
  spk1[j,i] = nextk[1]
```

```
  draws1[j,i] = nextk[2]
```

```
  scalek1[j,i] = nextk[3]
```

```
  nexta = metrop1step(func=spa.Gibbs, start.param=spa2[j,i-1],scale.param=scalea2[j,i-1],N=data[j,], spb=spb2[j,i-1],  
    spk=spk2[j,i-1],MuA=MuA2[i,1], SdA=SdA2[i,1], MuB=MuB2[i,1], SdB=SdB2[i,1], adjust.scale=adjust)
```

```
  spa2[j,i] = nexta[1]
```

```
  draws2[j,i] = nexta[2]
```

```
  scalea2[j,i] = nexta[3]
```

```
  nextb = metrop1step(func=spb.Gibbs, start.param=spb2[j,i-1], scale.param=scaleb2[j,i-1], N=data[j,], spa=spa2[j,i],  
    spk=spk2[j,i-1], MuA=MuA2[i,1],SdA=SdA2[i,1], MuB=MuB2[i,1], SdB=SdB2[i,1], adjust.scale=adjust)
```

```

spb2[j,i] = nextb[1]
drawsb2[j,i] = nextb[2]
scaleb2[j,i] = nextb[3]

nextk = metrop1step(func=spk.Gibbs, start.param=spk2[j,i-1], scale.param=scalek2[j,i-1], N=data[j,], spa=spa2[j,i],
  spb=spb2[j,i], adjust.scale=adjust)
spk2[j,i] = nextk[1]
drawsk2[j,i] = nextk[2]
scalek2[j,i] = nextk[3]

if(i%showstep==0 & i>=500)
{
  g = gelman.diag(mcmc.list(mcmc(spa1[j,((i-500)+1):i]), mcmc(spa2[j,((i-500)+1):i])), confidence = 0.95,
    transform=T, autoburnin=F)
  if (g[[1]][1] < 1.1 & !converged[j,1]) convergence[j,1] = i

  g = gelman.diag(mcmc.list(mcmc(spb1[j,((i-500)+1):i]), mcmc(spb2[j,((i-500)+1):i])), confidence = 0.95,
    transform=T, autoburnin=F)
  if (g[[1]][1] < 1.1 & !converged[j,2]) convergence[j,2] = i

  g = gelman.diag(mcmc.list(mcmc(sp1[j,((i-500)+1):i]), mcmc(sp2[j,((i-500)+1):i])), confidence = 0.95,
    transform=T, autoburnin=F)
  if (g[[1]][1] < 1.1 & !converged[j,3]) convergence[j,3] = i

  write.table(convergence, file="convergence.txt", sep="\t")
} # convergence test

} # species loop

write.table(t(round(spa1[,i], digits=4)), file="spa1.txt", sep="\t", append=T, col.names=F)
write.table(t(round(spb1[,i], digits=4)), file="spb1.txt", sep="\t", append=T, col.names=F)
write.table(t(round(sp1[,i], digits=4)), file="sp1.txt", sep="\t", append=T, col.names=F)
write.table(t(round(spa2[,i], digits=4)), file="spa2.txt", sep="\t", append=T, col.names=F)
write.table(t(round(spb2[,i], digits=4)), file="spb2.txt", sep="\t", append=T, col.names=F)
write.table(t(round(sp2[,i], digits=4)), file="sp2.txt", sep="\t", append=T, col.names=F)

write.table(t(drawsa1[,i]), file="drawsa1.txt", sep="\t", append=T, col.names=F)
write.table(t(drawsb1[,i]), file="drawsb1.txt", sep="\t", append=T, col.names=F)
write.table(t(drawsk1[,i]), file="drawsk1.txt", sep="\t", append=T, col.names=F)
write.table(t(drawsa2[,i]), file="drawsa2.txt", sep="\t", append=T, col.names=F)
write.table(t(drawsb2[,i]), file="drawsb2.txt", sep="\t", append=T, col.names=F)
write.table(t(drawsk2[,i]), file="drawsk2.txt", sep="\t", append=T, col.names=F)

write.table(t(MuA1[i,]), file="mua1.txt", sep="\t", append=T, col.names=F)
write.table(t(SdA1[i,]), file="sda1.txt", sep="\t", append=T, col.names=F)
write.table(t(MuA2[i,]), file="mua2.txt", sep="\t", append=T, col.names=F)
write.table(t(SdA2[i,]), file="sda2.txt", sep="\t", append=T, col.names=F)
write.table(t(MuB1[i,]), file="mub1.txt", sep="\t", append=T, col.names=F)
write.table(t(SdB1[i,]), file="sdb1.txt", sep="\t", append=T, col.names=F)
write.table(t(MuB2[i,]), file="mub2.txt", sep="\t", append=T, col.names=F)
write.table(t(SdB2[i,]), file="sdb2.txt", sep="\t", append=T, col.names=F)

write.table(t(scalea1[,i]), file="scalea1.txt", sep="\t", append=T, col.names=F)
write.table(t(scaleb1[,i]), file="scaleb1.txt", sep="\t", append=T, col.names=F)
write.table(t(scalek1[,i]), file="scalek1.txt", sep="\t", append=T, col.names=F)

```

```

cat(i, "\n")

} # steps loop
}

# likelihood of the mean of the normal hyperdistribution for a
mu.aGibbs=function(MuA,spa,SdA)
{
if(SdA<=0) return(-Inf)
llike = dnorm(spa, mean=MuA, sd=SdA, log=T)
return(sum(llike) + dunif(MuA, min=-10, max=1, log=T))
}

# likelihood of the standard deviation of the normal hyperdistribution for a
sd.aGibbs=function(SdA,spa,MuA)
{
if(SdA<=0) return(-Inf)
llike = dnorm(spa, mean=MuA, sd=SdA, log=T)
return(sum(llike) + dunif(SdA, min=0, max=2, log=T))
}

# likelihood of the mean of the normal hyperdistribution for b
mu.bGibbs=function(MuB,spb,SdB)
{
if(SdB<=0) return(-Inf)
llike = dnorm(spb, mean=MuB, sd=SdB, log=T)
return(sum(llike) + dunif(MuB, min=-5, max=5, log=T))
}

# likelihood of the standard deviation of the normal hyperdistribution for b
sd.bGibbs=function(SdB,spb,MuB)
{
if(SdB<=0) return(-Inf)
llike = dnorm(spb, mean=MuB, sd=SdB, log=T)
return(sum(llike) + dunif(SdB, min=0, max=2, log=T))
}

# likelihood of the constant of the linear relationship between log light and log
# recruitment
spa.Gibbs=function(spa,N,spb,spk,MuA,SdA,MuB,SdB)
{
if(spk<0) return(-Inf)
pred.nr = 10^(spa+spb*normlight)
if(any(pred.nr<0)) return(-Inf)
llike=sum(dnbinom(N,mu=pred.nr,size=spk,log=T))+dnorm(spa,mean=MuA,sd=SdA,log=T)
return(llike)
}

# likelihood of the coefficient of light in the linear relationship between log
# light and log recruitment
spb.Gibbs=function(spb,N,spa,spk,MuA,SdA,MuB,SdB)
{
if(spk<0) return(-Inf)
pred.nr = 10^(spa+spb*normlight)
if(any(pred.nr<0)) return(-Inf)

```



```

llike=sum(dnbinom(N,mu=pred.nr,size=spk,log=T))+dnorm(spb,mean=MuB,sd=SdB,log=T)
return(llike)
}

```

```

# likelihood of the cluster parameter of the negative binomial distribution
spk.Gibbs=function(spk,spa,spb,N)
{
if(spk<0) return(-Inf)
pred.nr = 10^(spa+spb*normlight)
if(any(pred.nr<0)) return(-Inf)
llike=sum(dnbinom(N,mu=pred.nr,size=spk,log=T))+dunif(spk,min=0,max=100,log=T)
return(llike)
}

```

```

# A generic routine for taking a single Metropolis step given any probability function, func, an initial parameter,
# start.param, and a scaling parameter for # the step size, scale.param. It returns the next parameter value as well as an
# acceptance indicator (0 if the value was accepted, 1 if rejected), and the new scale parameter. If adjust.scale=T, the
# scale parameter is adjusted to keep the acceptance rate at around 0.25, adjust.scale should be T only for
# steps<which.use (only during burnin period). Based on programs by Richard Condit and Helene Muller-Landau.
metrop1step=function(func,start.param,scale.param,adjust.scale=T,...)

```

```

{
new.scale = scale.param
probaccept = 0.25
multreject = 0.975
multaccept = multreject^(-(1-probaccept)/probaccept)
origlike = func(start.param,...)
newval = rnorm(1,mean=start.param,sd=scale.param)
newlike = func(newval,...)

```

```

if(!is.finite(newlike))
accept = F
else
{
if(newlike>=origlike) accept = T
else
{
likeratio=exp(newlike-origlike)
if(runif(1)<likeratio) accept = T
else accept = F
}
}
}

```

```

if(accept)
{
if(adjust.scale) new.scale=multaccept*scale.param
return(c(newval,0,new.scale))
}
else
{
if(adjust.scale) new.scale=multreject*scale.param
return(c(start.param,1,new.scale))
}
}

```













Response of recruitment to light availability across a tropical lowland rainforest community













Nadja Rüger, Andreas Huth, Stephen P. Hubbell, and Richard Condit













Supporting Information













Figure S1:













Model fit for the predicted number of recruits at different light conditions vs. the observed number of recruits for the woody species at Barro Colorado Island, Panama. Observed mean number of recruits (1985–1990) in intervals of 10-percentiles of light in the 5×5 m quadrats (dots, only means different from zero are plotted), and predicted number of recruits using posterior means for the second interval 1985–1990 (line).













Acacia melanoceras**Acalypha diversifolia****Acalypha macrostachya****Adelia triloba****Aegiphila panamensis****Alchornea costaricensis****Alchornea latifolia****Alibertia edulis****Allophylus psilospermus****Alseis blackiana****Anaxagorea panamensis****Andira inermis**













Annona acuminata**Annona spraguei****Apeiba membranacea****Apeiba tibourbou****Apelandra sinclairiana****Appunia seibertii****Ardisia fendleri****Ardisia guianensis****Aspidosperma spruceanum****Astrocaryum standleyanum****Astronium graveolens****Bactris barronis**













Bactris coloradonis**Bactris major****Banara guianensis****Beilschmiedia pendula****Brosimum alicastrum****Calophyllum longifolium****Capparis frondosa****Casearia aculeata****Casearia arborea****Casearia guianensis****Casearia sylvestris****Cassipourea elliptica**













Cavanillesia platanifolia**Cecropia insignis****Cecropia obtusifolia****Cedrela odorata****Ceiba pentandra****Celtis schippii****Cespedesia spathulata****Cestrum megalophyllum****Chamaedorea tepejilote****Chamguava schippii****Chrysochlamys eclipses****Chrysophyllum argenteum**













Chrysophyllum cainito**Clidemia dentata****Clidemia octona****Coccoloba coronata****Coccoloba manzinellensis****Conostegia bracteata****Conostegia cinnamomea****Cordia alliodora****Cordia bicolor****Cordia lasiocalyx****Coussarea curvigemma****Croton billbergianus**













Cupania cinerea**Cupania latifolia****Cupania rufescens****Cupania seemanii****Solanum circinatum****Dendropanax arboreus****Desmopsis panamensis****Diospyros artanthifolia****Dipteryx oleifera****Drypetes standleyi****Elaeis oleifera****Erythrina costaricensis**













Erythroxyllum macrophyllum**Erythroxyllum panamense****Eugenia coloradoensis****Eugenia galonensis****Eugenia nesiotica****Eugenia oerstediana****Faramea occidentalis****Ficus citrifolia****Ficus insipida****Ficus maxima****Ficus obtusifolia****Ficus pertusa**













Ficus popenoei**Garcinia intermedia****Garcinia madruno****Genipa americana****Guapira standleyana****Guarea grandifolia****Guarea guidonia****Guarea fuzzy****Guatteria dumetorum****Guazuma ulmifolia****Guettarda foliacea****Gustavia superba**













Hamelia axillaris**Hampea appendiculata****Hasseltia floribunda****Heisteria acuminata****Heisteria concinna****Herrania purpurea****Hirtella americana****Hirtella triandra****Hybanthus prunifolius****Hieronyma alchorneoides****Inga cocleensis****Inga laurina**













Inga goldmanii**Inga marginata****Inga oerstediana****Inga pezizifera****Inga punctata****Inga nobilis****Inga ruiziana****Inga acuminata****Inga sapindoides****Inga spectabilis****Inga thibaudiana****Inga umbellifera**













Jacaranda copaia**Koanophyllon wetmorei****Lacistema aggregatum****Lacmellea panamensis****Laetia procera****Laetia thamnina****Leandra dichotoma****Licania hypoleuca****Licania platypus****Lindackeria laurina****Lonchocarpus heptaphyllus****Luehea seemannii**













Macrocnemum roseum**Mosannonna garwoodii****Malpighia romeroana****Maquira guianensis****Margaritaria nobilis****Maytenus schippii****Miconia affinis****Miconia argentea****Miconia elata****Miconia hondurensis****Miconia impetolaris****Miconia nervosa**













Miconia prasina**Mouriri myrtilloides****Myrcia gatunensis****Nectandra cissiflora****Nectandra lineata****Nectandra purpurea****Neea amplifolia****Ochroma pyramidale****Ocotea cernua****Ocotea oblonga****Ocotea puberula****Ocotea whitei**













Oenocarpus mapora**Trophis caucana****Ormosia coccinea****Ormosia macrocalyx****Ouratea lucens****Palicourea guianensis****Pentagonia macrophylla****Perebea xanthochyma****Cinnamomum triplinerve****Picramnia latifolia****Piper arboreum****Piper aequale**













Piper cordulatum**Piper colonense****Piper perlasense****Piper reticulatum****Platymiscium pinnatum****Platypodium elegans****Pachira quinata****Posoqueria latifolia****Poulsenia armata****Pourouma bicolor****Pouteria reticulata****Pouteria stipitata**

Prioria copaifera**Protium costaricense****Protium panamense****Protium confusum****Protium tenuifolium****Pseudobombax septenatum****Psidium friedrichsthalianum****Psychotria acuminata****Psychotria deflexa****Psychotria hoffmannseggiana****Psychotria graciliflora****Psychotria grandis**

Psychotria horizontalis**Psychotria limonensis****Psychotria marginata****Psychotria pittieri****Psychotria racemosa****Pterocarpus rohrii****Quararibea asterolepis****Quassia amara****Randia armata****Rinorea sylvatica****Sapium glandulosum****Attalea butyracea**

Schizolobium parahyba**Senna dariensis****Simarouba amara****Siparuna guianensis****Siparuna pauciflora****Sloanea terniflora****Socratea exorrhiza****Solanum asperum****Solanum hayesii****Sorocea affinis****Spondias mombin****Spondias radlkoferi**

Sterculia apetala**Stylogyne turbacensis****Swartzia simplex_var.grandiflora****Symphonia globulifera****Tabebuia guayacan****Tabebuia rosea****Tabernaemontana arborea****Tachigali versicolor****Talisia nervosa****Talisia princeps****Terminalia amazonia****Terminalia oblonga**

Tetragastris panamensis**Thevetia ahouai****Trattinnickia aspera****Trema micrantha****Trichanthera gigantea****Trichilia pallida****Trichilia tuberculata****Trichospermum galeottii****Triplaris cumingiana****Trophis racemosa****Turpinia occidentalis****Unonopsis pittieri**













Urera baccifera**Virola sebifera****Virola multiflora****Virola surinamensis****Vismia baccifera****Vochysia ferruginea****Xylopia macrantha****Xylosma oligandra****Zanthoxylum ekmanii****Zanthoxylum panamense****Zanthoxylum acuminatum****Zuelania guidonia**

Table S1: Posterior means, lower and upper limits of the credible interval (CI -, CI +) of the log of the mean number of recruits (a), the light effect (b), the clustering parameter of the binomial distribution (c), and the light level at which recruitment falls to 50% of recruitment at 20% light (Light50) for two census intervals (1985-1990, 1990-1995) and tree species in the 50-ha Forest Dynamics Plot at Barro Colorado Island, Panama. Dashes indicate no recruitment in the respective census interval.

Species	Parameters											
	a						b					
	1985-1990			1990-1995			1985-1990			1990-1995		
	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +
<i>Acacia melanoceras</i>	-3.69	-4.12	-3.3	-4.18	-4.91	-3.58	-0.34	-1.37	0.79	0.47	-0.74	1.74
<i>Acalypha diversifolia</i>	-2.08	-2.16	-2	-2.23	-2.32	-2.13	1.17	0.88	1.43	1.1	0.76	1.44
<i>Acalypha macrostachya</i>	-3.38	-3.67	-3.08	-3.28	-3.66	-2.95	1.23	0.46	2.06	1.45	0.39	2.42
<i>Adelia triloba</i>	-3.03	-3.28	-2.79	-3.69	-4.13	-3.26	1.87	1.17	2.66	1.16	0.19	2.24
<i>Aegiphila panamensis</i>	-3.74	-4.22	-3.35	-3.75	-4.18	-3.33	1.05	-0.01	2.01	1.52	0.56	2.5
<i>Alchornea costaricensis</i>	-3.2	-3.46	-2.97	-3.65	-4.1	-3.23	1.85	1.29	2.42	2.24	1.23	3.23
<i>Alchornea latifolia</i>	-4.17	-4.86	-3.54	-	-	-	0.32	-0.88	1.53	-	-	-
<i>Alibertia edulis</i>	-2.49	-2.62	-2.37	-2.76	-2.93	-2.61	0.41	-0.02	0.82	0.85	0.35	1.37
<i>Allophylus psilospermus</i>	-3.09	-3.32	-2.87	-3.5	-3.86	-3.17	0.65	-0.04	1.35	0.83	-0.08	1.76
<i>Alseis blackiana</i>	-1.24	-1.27	-1.21	-1.54	-1.58	-1.49	0.87	0.76	0.99	1.11	0.97	1.27
<i>Anaxagorea panamensis</i>	-2.61	-2.82	-2.38	-2.3	-2.5	-2.11	3.11	2.22	3.97	1.84	1.06	2.71
<i>Andira inermis</i>	-2.95	-3.15	-2.76	-3.69	-4.13	-3.25	0.4	-0.25	1.06	-0.02	-1.16	1.01
<i>Annona acuminata</i>	-2.44	-2.57	-2.33	-2.54	-2.67	-2.42	1.64	1.32	1.97	1.1	0.7	1.49
<i>Annona spraguei</i>	-2.59	-2.73	-2.45	-2.6	-2.75	-2.47	1.91	1.45	2.38	1.82	1.42	2.25
<i>Apeiba membranacea</i>	-3.55	-3.93	-3.19	-3.83	-4.32	-3.38	1.82	1.01	2.61	1.27	0.22	2.28
<i>Apeiba tibourbou</i>	-3.91	-4.46	-3.41	-3.97	-4.55	-3.45	1.35	0.27	2.52	1.07	0	2.18
<i>Aphelandra sinclairiana</i>	-3.99	-4.52	-3.4	-3.98	-4.53	-3.41	0.86	-0.29	1.97	1.2	0.01	2.3
<i>Appunia seibertii</i>	-3.98	-4.52	-3.44	-	-	-	0.76	-0.42	1.97	-	-	-
<i>Ardisia bartlettii</i>	-	-	-	-4.17	-4.94	-3.57	-	-	-	0.79	-0.49	1.9
<i>Ardisia fendleri</i>	-3.51	-3.84	-3.16	-3.37	-3.69	-3.06	-0.19	-1.08	0.77	-0.72	-1.58	0.14
<i>Ardisia guianensis</i>	-4.13	-4.66	-3.61	-3.73	-4.18	-3.33	2.68	1.85	3.63	1.37	0.28	2.27
<i>Aspidosperma spruceanum</i>	-2.66	-2.79	-2.53	-2.85	-3.02	-2.67	0.36	-0.1	0.87	0.34	-0.24	0.93
<i>Astrocaryum standleyanum</i>	-3.37	-3.69	-3.09	-	-	-	0.24	-0.68	1.17	-	-	-
<i>Astronium graveolens</i>	-3.5	-3.91	-3.13	-3.3	-3.6	-3.05	1.42	0.53	2.57	1.18	0.42	1.96
<i>Attalea butyracea</i>	-4.2	-4.89	-3.57	-	-	-	0.36	-0.83	1.69	-	-	-
<i>Bactris barronis</i>	-3.95	-4.5	-3.42	-	-	-	0	-1.33	1.07	-	-	-
<i>Bactris coloradonis</i>	-4.21	-4.89	-3.52	-	-	-	0.46	-0.81	1.69	-	-	-
<i>Bactris major</i>	-2.88	-3.09	-2.69	-3.33	-3.63	-3.03	1.24	0.6	1.86	-0.06	-0.94	0.78
<i>Banara guianensis</i>	-3.86	-4.34	-3.37	-4.18	-4.85	-3.52	1.04	0.04	2.25	0.7	-0.6	1.84
<i>Beilschmiedia pendula</i>	-1.67	-1.72	-1.61	-1.96	-2.03	-1.9	0.07	-0.1	0.27	-0.07	-0.31	0.15
<i>Brosimum alicastrum</i>	-2.54	-2.66	-2.43	-2.85	-3.05	-2.68	0.11	-0.3	0.56	-0.3	-0.87	0.29
<i>Calophyllum longifolium</i>	-1.96	-2.04	-1.89	-2	-2.07	-1.92	0.29	0.01	0.59	0.37	0.13	0.65
<i>Capparis frondosa</i>	-1.92	-1.98	-1.86	-2.31	-2.41	-2.22	0.24	0.02	0.45	-0.01	-0.34	0.35
<i>Casearia aculeata</i>	-2.36	-2.46	-2.27	-2.62	-2.75	-2.48	0.76	0.43	1.07	0.89	0.41	1.29
<i>Casearia arborea</i>	-3.34	-3.66	-3.07	-3.36	-3.66	-3.06	1.48	0.76	2.19	1.28	0.47	2.11
<i>Casearia guianensis</i>	-3.95	-4.55	-3.43	-3.95	-4.47	-3.44	0.03	-1.14	1.16	0.71	-0.4	1.97
<i>Casearia sylvestris</i>	-3.01	-3.23	-2.81	-3.18	-3.42	-2.93	1.05	0.42	1.7	0.3	-0.47	1.07
<i>Cassipourea elliptica</i>	-2.14	-2.22	-2.06	-2.34	-2.44	-2.25	0.57	0.28	0.85	0.5	0.16	0.87
<i>Cavanillesia platanifolia</i>	-4.2	-4.91	-3.57	-4.17	-4.9	-3.58	0.56	-0.79	1.71	0.72	-0.56	1.86
<i>Cecropia insignis</i>	-2.85	-3.04	-2.66	-2.49	-2.64	-2.36	2.94	2.4	3.46	2.16	1.65	2.69
<i>Cecropia longipes</i>	-	-	-	-4.21	-4.97	-3.64	-	-	-	1.31	0.11	2.52
<i>Cecropia obtusifolia</i>	-3.13	-3.36	-2.87	-3.03	-3.25	-2.82	2.11	1.48	2.87	1.79	1.26	2.35
<i>Cedrela odorata</i>	-3.44	-3.79	-3.13	-	-	-	0.46	-0.45	1.32	-	-	-
<i>Ceiba pentandra</i>	-3.53	-3.91	-3.2	-4.01	-4.6	-3.49	0.59	-0.42	1.56	1.42	0.21	2.5
<i>Celtis schippii</i>	-3.69	-4.07	-3.24	-4.18	-4.93	-3.54	0.07	-1.04	1.08	0.78	-0.43	1.97
<i>Cespedesia spathulata</i>	-4.18	-4.9	-3.6	-	-	-	0.14	-1.11	1.45	-	-	-
<i>Cestrum megalophyllum</i>	-2.74	-2.9	-2.59	-3.36	-3.7	-3.06	0.77	0.27	1.28	0.76	-0.05	1.61
<i>Chamaedorea tepejilote</i>	-3.69	-4.11	-3.26	-3.8	-4.29	-3.32	0.45	-0.5	1.53	0.86	-0.07	1.98
<i>Changuava schippii</i>	-2.63	-2.77	-2.48	-2.59	-2.74	-2.45	1.77	1.32	2.2	1.51	1.04	2
<i>Chrysochlamys eclipses</i>	-3.07	-3.28	-2.87	-3.11	-3.34	-2.89	-0.12	-0.84	0.59	0.72	0	1.44
<i>Chrysophyllum argenteum</i>	-1.99	-2.06	-1.92	-2.37	-2.48	-2.28	0.84	0.6	1.07	1.03	0.67	1.35
<i>Chrysophyllum cainito</i>	-2.83	-3	-2.66	-2.85	-3.02	-2.68	1.23	0.71	1.71	1.3	0.74	1.81

Species	Parameters											
	<i>a</i>						<i>b</i>					
	1985-1990			1990-1995			1985-1990			1990-1995		
mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	
<i>Cinnamomum triplinerve</i>	-2.98	-3.19	-2.78	-3.2	-3.44	-2.94	1.23	0.68	1.81	0.99	0.23	1.7
<i>Clidemia dentata</i>	-3.17	-3.44	-2.92	-3.63	-4.01	-3.25	1.3	0.62	1.96	1.35	0.47	2.28
<i>Clidemia octona</i>	-3.49	-3.82	-3.15	-3.52	-3.85	-3.17	1.34	0.51	2.2	1.12	0.16	2.08
<i>Coccoloba coronata</i>	-2.89	-3.07	-2.73	-3.26	-3.52	-2.98	0.47	-0.09	1.09	0.51	-0.3	1.35
<i>Coccoloba manzinellensis</i>	-2.62	-2.75	-2.48	-2.8	-2.96	-2.63	0.85	0.36	1.25	1.02	0.44	1.56
<i>Colubrina glandulosa</i>	-	-	-	-4.18	-4.86	-3.53	-	-	-	1.15	0.05	2.35
<i>Conostegia bracteata</i>	-4.22	-4.94	-3.59	-4.18	-4.94	-3.57	0.74	-0.48	2.02	0.64	-0.51	1.92
<i>Conostegia cinnamomea</i>	-2.44	-2.55	-2.34	-2.63	-2.77	-2.49	0.41	0.02	0.86	0.79	0.29	1.21
<i>Cordia alliodora</i>	-3	-3.23	-2.8	-3.23	-3.48	-2.96	1.52	0.94	2.12	0.84	0.06	1.63
<i>Cordia bicolor</i>	-1.9	-1.97	-1.83	-2.13	-2.21	-2.05	1.83	1.56	2.08	1.67	1.36	1.95
<i>Cordia lasiocalyx</i>	-1.95	-2.02	-1.89	-2.09	-2.18	-2.02	0.59	0.34	0.81	0.52	0.25	0.79
<i>Coussarea curvigemma</i>	-1.72	-1.76	-1.66	-1.9	-1.96	-1.85	0.68	0.5	0.86	0.44	0.24	0.67
<i>Croton billbergianus</i>	-1.81	-1.87	-1.73	-2.28	-2.4	-2.19	2.34	2.07	2.59	2.05	1.63	2.4
<i>Cupania cinerea</i>	-3.65	-4.06	-3.25	-	-	-	1.02	0.06	2.08	-	-	-
<i>Cupania latifolia</i>	-3.32	-3.6	-3.05	-3.66	-4.09	-3.23	0.39	-0.48	1.29	0.33	-0.63	1.42
<i>Cupania rufescens</i>	-2.9	-3.09	-2.7	-2.94	-3.15	-2.74	1.64	1.01	2.2	1.34	0.76	1.9
<i>Cupania seemanii</i>	-2.09	-2.16	-2.01	-2.23	-2.32	-2.15	0.72	0.49	0.96	0.79	0.47	1.09
<i>Dendropanax arboreus</i>	-4.2	-4.89	-3.53	-	-	-	0.64	-0.75	1.84	-	-	-
<i>Desmopsis panamensis</i>	-1.01	-1.03	-0.99	-1.23	-1.26	-1.2	0.16	0.08	0.24	0.14	0.04	0.24
<i>Diospyros artanthifolia</i>	-3.21	-3.49	-2.99	-3.31	-3.59	-3.03	1.34	0.64	2	0.39	-0.43	1.22
<i>Dipteryx oleifera</i>	-3.97	-4.61	-3.49	-4.19	-4.86	-3.58	0.47	-0.68	1.7	0.95	-0.21	2.13
<i>Drypetes standleyi</i>	-2.07	-2.14	-2.01	-2.35	-2.46	-2.25	-0.04	-0.29	0.23	-0.44	-0.79	-0.1
<i>Elaeis oleifera</i>	-3.84	-4.3	-3.33	-	-	-	0.89	-0.22	2.01	-	-	-
<i>Erythrina costaricensis</i>	-4.19	-4.9	-3.54	-4.16	-4.85	-3.54	0.29	-0.98	1.63	0.74	-0.51	1.89
<i>Erythroxylum macrophyllum</i>	-2.52	-2.63	-2.4	-2.75	-2.9	-2.59	0.72	0.31	1.1	0.34	-0.19	0.88
<i>Erythroxylum panamense</i>	-3.08	-3.31	-2.87	-3.3	-3.6	-3.03	0.58	-0.14	1.25	0.56	-0.24	1.39
<i>Eugenia coloradoensis</i>	-2.26	-2.35	-2.17	-2.51	-2.63	-2.4	0.72	0.43	1.01	0.37	-0.02	0.78
<i>Eugenia galalonensis</i>	-1.81	-1.86	-1.76	-1.87	-1.94	-1.82	0.37	0.18	0.56	0.73	0.54	0.96
<i>Eugenia nesiotica</i>	-2.53	-2.65	-2.42	-2.67	-2.81	-2.53	0.42	-0.01	0.83	0.66	0.15	1.18
<i>Eugenia oerstediana</i>	-1.63	-1.68	-1.59	-1.91	-1.97	-1.84	0.37	0.19	0.53	0.48	0.24	0.72
<i>Faramea occidentalis</i>	-0.76	-0.78	-0.75	-0.99	-1.01	-0.97	0.26	0.19	0.32	0.22	0.14	0.3
<i>Ficus bullenei</i>	-	-	-	-4.19	-4.89	-3.55	-	-	-	1.09	-0.07	2.26
<i>Ficus citrifolia</i>	-4.02	-4.66	-3.51	-	-	-	1.12	-0.05	2.2	-	-	-
<i>Ficus costaricana</i>	-	-	-	-4.2	-4.93	-3.55	-	-	-	0.99	-0.12	2.27
<i>Ficus insipida</i>	-3.99	-4.55	-3.44	-3.85	-4.34	-3.43	0.7	-0.38	1.86	1.42	0.34	2.41
<i>Ficus maxima</i>	-4.01	-4.61	-3.48	-	-	-	1.08	-0.1	2.17	-	-	-
<i>Ficus obtusifolia</i>	-4.2	-4.89	-3.55	-3.95	-4.5	-3.41	0.41	-0.86	1.7	0.56	-0.53	1.81
<i>Ficus pertusa</i>	-4.21	-4.89	-3.58	-4.18	-4.84	-3.55	0.76	-0.47	2.04	0.95	-0.23	2.17
<i>Ficus popenoei</i>	-4.26	-4.94	-3.62	-	-	-	1.2	-0.01	2.44	-	-	-
<i>Ficus trigonata</i>	-	-	-	-4.18	-4.9	-3.6	-	-	-	0.91	-0.35	2.05
<i>Garcinia intermedia</i>	-1.58	-1.62	-1.54	-1.81	-1.87	-1.76	0.16	0	0.31	0.16	-0.04	0.36
<i>Garcinia madruno</i>	-2.49	-2.6	-2.38	-2.87	-3.07	-2.71	-0.1	-0.52	0.28	-0.04	-0.61	0.56
<i>Genipa americana</i>	-3.97	-4.52	-3.4	-	-	-	0.05	-1.07	1.3	-	-	-
<i>Guapira standleyana</i>	-3.58	-3.97	-3.24	-4.17	-4.88	-3.54	0.35	-0.67	1.39	0.38	-0.81	1.56
<i>Guarea fuzzy</i>	-2.1	-2.17	-2.03	-2.37	-2.48	-2.28	0.47	0.19	0.72	0.54	0.18	0.89
<i>Guarea grandifolia</i>	-3.51	-3.86	-3.2	-3.44	-3.76	-3.13	-0.05	-1.09	0.81	1.03	0.14	1.83
<i>Guarea guidonia</i>	-1.89	-1.95	-1.83	-2.07	-2.14	-2	0.49	0.26	0.69	0.5	0.24	0.75
<i>Guatteria dumetorum</i>	-2.17	-2.25	-2.09	-2.47	-2.59	-2.36	0.59	0.31	0.85	0.81	0.43	1.2
<i>Guazuma ulmifolia</i>	-3.75	-4.17	-3.36	-3.88	-4.43	-3.4	2.14	1.36	2.93	1.61	0.53	2.61
<i>Guettarda foliacea</i>	-2.96	-3.17	-2.78	-3.78	-4.27	-3.37	0.55	-0.13	1.17	0.34	-0.81	1.45
<i>Gustavia superba</i>	-2.79	-2.96	-2.62	-3.22	-3.48	-2.93	1.37	0.88	1.81	1.92	1.33	2.62
<i>Hamelia axillaris</i>	-2.71	-2.86	-2.55	-2.73	-2.89	-2.57	1.69	1.28	2.07	1.64	1.18	2.15
<i>Hampea appendiculata</i>	-3.35	-3.65	-3.05	-3.53	-3.88	-3.17	1.52	0.82	2.28	1.23	0.38	2.14
<i>Hasseltia floribunda</i>	-2.84	-3.01	-2.66	-3.21	-3.49	-2.98	0.76	0.19	1.44	1.46	0.74	2.17
<i>Heisteria acuminata</i>	-3.74	-4.17	-3.3	-3.67	-4.08	-3.27	1.01	-0.06	1.95	0.27	-0.76	1.31
<i>Heisteria concinna</i>	-2.36	-2.45	-2.27	-2.93	-3.11	-2.73	0.05	-0.32	0.39	0.58	-0.06	1.2

Species	Parameters											
	a						b					
	1985-1990			1990-1995			1985-1990			1990-1995		
mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	
<i>Herrania purpurea</i>	-2.48	-2.58	-2.36	-2.79	-2.97	-2.64	0.48	0.08	0.87	1.19	0.68	1.72
<i>Hieronyma alchorneoides</i>	-3.35	-3.65	-3.05	-3.97	-4.53	-3.47	0.79	-0.1	1.56	1.11	0.04	2.26
<i>Hirtella americana</i>	-3.72	-4.18	-3.34	-4.17	-4.86	-3.57	0.86	-0.24	1.88	0.77	-0.42	2.04
<i>Hirtella triandra</i>	-1.5	-1.54	-1.47	-1.77	-1.82	-1.72	0.13	-0.02	0.27	0.24	0.05	0.43
<i>Hybanthus prunifolius</i>	-0.55	-0.56	-0.53	-0.88	-0.9	-0.86	0.13	0.07	0.19	0.19	0.12	0.27
<i>Inga acuminata</i>	-2.42	-2.53	-2.33	-2.52	-2.64	-2.4	0.64	0.27	0.99	0.25	-0.16	0.66
<i>Inga cocleensis</i>	-3.34	-3.67	-3.07	-3.51	-3.89	-3.18	1.73	1.08	2.4	1.05	0.13	1.96
<i>Inga goldmanii</i>	-2.47	-2.57	-2.36	-3.16	-3.4	-2.91	0.61	0.23	0.98	1.02	0.3	1.74
<i>Inga laurina</i>	-3.01	-3.22	-2.81	-3.31	-3.58	-3.01	1.07	0.47	1.72	0.68	-0.2	1.49
<i>Inga marginata</i>	-1.97	-2.04	-1.9	-2.33	-2.42	-2.22	0.88	0.64	1.13	1.22	0.91	1.55
<i>Inga nobilis</i>	-2.41	-2.51	-2.3	-2.86	-3.03	-2.68	0.27	-0.09	0.65	0.24	-0.31	0.83
<i>Inga oerstediana</i>	-4.02	-4.6	-3.51	-4.17	-4.88	-3.51	1.12	-0.01	2.27	0.54	-0.7	1.75
<i>Inga pezizifera</i>	-2.85	-3.02	-2.69	-3.36	-3.67	-3.07	-0.18	-0.78	0.4	0.68	-0.21	1.52
<i>Inga punctata</i>	-3.5	-3.86	-3.17	-3.95	-4.52	-3.43	1.39	0.51	2.24	1.11	-0.08	2.15
<i>Inga ruiziana</i>	-4.22	-4.96	-3.6	-	-	-	0.88	-0.26	2.22	-	-	-
<i>Inga sapindoides</i>	-2.82	-2.99	-2.64	-3.32	-3.6	-3.02	1.14	0.64	1.68	1.61	0.9	2.34
<i>Inga spectabilis</i>	-3.99	-4.54	-3.48	-	-	-	0.84	-0.38	2.01	-	-	-
<i>Inga thibaudiana</i>	-3.02	-3.24	-2.81	-2.89	-3.08	-2.71	1.65	1.03	2.12	1.67	1.05	2.32
<i>Inga umbellifera</i>	-2.18	-2.26	-2.1	-2.62	-2.74	-2.47	0.88	0.59	1.18	0.96	0.53	1.39
<i>Jacaranda copaia</i>	-3.48	-3.8	-3.14	-3.34	-3.61	-3.03	1.87	1.12	2.6	1.7	0.96	2.44
<i>Koanophyllon wetmorei</i>	-3.74	-4.17	-3.31	-4.19	-4.92	-3.56	1.04	-0.1	2.03	1.13	-0.16	2.28
<i>Lacistema aggregatum</i>	-1.81	-1.86	-1.76	-1.96	-2.02	-1.9	0.13	-0.05	0.33	0.47	0.25	0.7
<i>Lacmellea panamensis</i>	-3.46	-3.81	-3.16	-3.67	-4.09	-3.26	0.55	-0.46	1.46	0.4	-0.77	1.39
<i>Laetia procerca</i>	-3.6	-4.01	-3.23	-3.84	-4.35	-3.39	2	1.11	3.05	1.24	0.22	2.35
<i>Laetia thamnia</i>	-2.34	-2.44	-2.25	-2.49	-2.61	-2.37	0.96	0.66	1.27	0.8	0.4	1.19
<i>Leandra dichotoma</i>	-4.26	-5.03	-3.64	-	-	-	1.17	-0.16	2.32	-	-	-
<i>Licania hypoleuca</i>	-3.07	-3.28	-2.84	-3.3	-3.6	-3	0.03	-0.7	0.67	0.39	-0.47	1.16
<i>Licania platypus</i>	-2.59	-2.71	-2.45	-2.97	-3.18	-2.79	-0.39	-0.83	0.03	-0.67	-1.29	-0.06
<i>Lindackeria laurina</i>	-3.73	-4.21	-3.32	-3.96	-4.54	-3.44	0.92	-0.12	1.93	0.88	-0.23	2
<i>Lonchocarpus heptaphyllus</i>	-2.16	-2.24	-2.08	-2.43	-2.55	-2.32	0.93	0.67	1.2	1.35	0.95	1.71
<i>Luehea seemannii</i>	-2.77	-2.94	-2.61	-3	-3.22	-2.79	1.94	1.54	2.34	2.06	1.53	2.52
<i>Macrocnemum roseum</i>	-3.27	-3.6	-2.96	-4.18	-4.86	-3.59	-0.25	-1.2	0.6	0.94	-0.25	2.12
<i>Malpighia romeroana</i>	-3.44	-3.79	-3.16	-3.94	-4.52	-3.43	0.54	-0.41	1.45	0.69	-0.46	1.84
<i>Maquira guianensis</i>	-2.06	-2.12	-1.99	-2.37	-2.47	-2.26	-0.07	-0.32	0.19	-0.12	-0.5	0.22
<i>Margaritaria nobilis</i>	-4.14	-4.74	-3.58	-	-	-	1.71	0.55	2.8	-	-	-
<i>Maytenus schippii</i>	-3.38	-3.69	-3.09	-3.66	-4.08	-3.27	0.29	-0.61	1.2	0.36	-0.64	1.44
<i>Miconia affinis</i>	-2.34	-2.46	-2.22	-2.51	-2.63	-2.38	1.62	1.22	2.03	1.45	1	1.88
<i>Miconia argentea</i>	-1.78	-1.85	-1.72	-1.87	-1.94	-1.79	1.76	1.55	1.99	1.97	1.7	2.27
<i>Miconia elata</i>	-3.51	-4.01	-3.07	-	-	-	0.93	-0.19	1.97	-	-	-
<i>Miconia hondurensis</i>	-3.18	-3.45	-2.94	-3.31	-3.59	-3.03	0.73	0.03	1.48	0.44	-0.41	1.25
<i>Miconia impetolaris</i>	-3.67	-4.1	-3.29	-4.18	-4.85	-3.56	0.2	-0.86	1.27	0.76	-0.42	1.98
<i>Miconia nervosa</i>	-2.1	-2.19	-2.01	-2.27	-2.36	-2.17	1.53	1.24	1.82	1.39	1.05	1.7
<i>Miconia prasina</i>	-4	-4.61	-3.46	-	-	-	0.88	-0.27	2.04	-	-	-
<i>Mosammona garwoodii</i>	-2.55	-2.66	-2.43	-2.59	-2.73	-2.47	0.3	-0.14	0.71	0.03	-0.42	0.48
<i>Mouriri myrtilloides</i>	-1.2	-1.22	-1.17	-1.38	-1.42	-1.35	0.57	0.48	0.68	0.61	0.48	0.74
<i>Myrcia gatunensis</i>	-3.37	-3.66	-3.08	-3.58	-4	-3.23	0.26	-0.66	1.08	0.49	-0.5	1.46
<i>Nectandra cissiflora</i>	-2.82	-2.98	-2.66	-3.08	-3.3	-2.85	1.06	0.55	1.57	1.12	0.47	1.77
<i>Nectandra lineata</i>	-2.64	-2.77	-2.52	-2.9	-3.08	-2.72	0.47	-0.01	0.94	0.9	0.32	1.45
<i>Nectandra purpurea</i>	-3.13	-3.36	-2.88	-3.38	-3.71	-3.11	0.79	0.09	1.48	1.02	0.17	1.84
<i>Neea amplifolia</i>	-3.05	-3.28	-2.85	-3.26	-3.53	-3.01	0.59	-0.04	1.32	1.24	0.49	1.92
<i>Ochroma pyramidale</i>	-4.25	-4.92	-3.56	-4.18	-4.9	-3.58	1.1	-0.14	2.37	0.87	-0.39	2
<i>Ocotea cernua</i>	-2.35	-2.45	-2.25	-2.7	-2.84	-2.55	0.61	0.25	0.94	0.87	0.39	1.37
<i>Ocotea oblonga</i>	-2.46	-2.56	-2.35	-2.65	-2.79	-2.52	0.73	0.37	1.12	0.86	0.41	1.31
<i>Ocotea puberula</i>	-2.63	-2.77	-2.5	-2.91	-3.09	-2.72	0.9	0.48	1.35	0.76	0.17	1.39
<i>Ocotea whitei</i>	-2.56	-2.68	-2.43	-3.25	-3.53	-2.98	0	-0.45	0.43	-0.43	-1.26	0.31
<i>Denocarpus mapora</i>	-2.09	-2.16	-2.02	-2.25	-2.34	-2.16	0.45	0.19	0.71	0.15	-0.18	0.45

Species	Parameters											
	<i>a</i>						<i>b</i>					
	1985-1990			1990-1995			1985-1990			1990-1995		
mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	
<i>Ormosia coccinea</i>	-3.16	-3.41	-2.93	-3.29	-3.57	-3.03	0.13	-0.63	0.89	-0.13	-0.96	0.67
<i>Ormosia macrocalyx</i>	-2.98	-3.17	-2.79	-3.32	-3.62	-3.05	0.47	-0.18	1.11	0.76	-0.08	1.6
<i>Ouratea lucens</i>	-2.19	-2.27	-2.11	-2.44	-2.55	-2.34	0.67	0.38	0.94	0.76	0.39	1.16
<i>Pachira quinata</i>	-4.22	-4.99	-3.6	-	-	-	0.77	-0.39	2.02	-	-	-
<i>Pachira sessilis</i>	-	-	-	-4.19	-4.91	-3.56	-	-	-	0.95	-0.24	2.18
<i>Palicourea guianensis</i>	-1.45	-1.49	-1.4	-1.63	-1.69	-1.57	1.95	1.76	2.12	2.25	2.02	2.5
<i>Pentagonia macrophylla</i>	-2.67	-2.8	-2.53	-3.3	-3.59	-3.02	0.65	0.16	1.11	0.39	-0.4	1.26
<i>Perebea xanthochyma</i>	-2.69	-2.83	-2.56	-3.26	-3.53	-3.02	0.51	0.01	0.98	0.48	-0.4	1.29
<i>Picramnia latifolia</i>	-2.08	-2.14	-2	-2.27	-2.35	-2.17	0.25	-0.02	0.5	0.26	-0.08	0.59
<i>Piper aequale</i>	-3.32	-3.63	-3.05	-3.67	-4.12	-3.29	0.17	-0.71	0.95	0.87	-0.17	1.86
<i>Piper arboreum</i>	-3.81	-4.31	-3.39	-3.86	-4.42	-3.36	0.68	-0.45	1.88	1.45	0.3	2.49
<i>Piper colonense</i>	-3.59	-4	-3.23	-4.15	-4.82	-3.54	0.43	-0.69	1.39	0.56	-0.75	1.79
<i>Piper cordulatum</i>	-1.94	-2	-1.87	-2.82	-2.98	-2.65	0.63	0.38	0.89	1.02	0.43	1.51
<i>Piper perlasense</i>	-3.31	-3.59	-3.02	-3.94	-4.47	-3.41	-0.58	-1.39	0.21	0.6	-0.48	1.77
<i>Piper reticulatum</i>	-3.36	-3.66	-3.06	-3.94	-4.52	-3.44	0.8	-0.11	1.6	0.49	-0.74	1.64
<i>Platymiscium pinnatum</i>	-2.9	-3.08	-2.71	-2.79	-2.97	-2.63	1.18	0.59	1.7	1.7	1.26	2.18
<i>Platyopodium elegans</i>	-2.92	-3.11	-2.75	-3.12	-3.34	-2.89	1.07	0.55	1.66	0.93	0.18	1.62
<i>Posoqueria latifolia</i>	-4.23	-4.95	-3.59	-3.5	-3.85	-3.16	0.88	-0.38	2.05	0.12	-0.89	1.01
<i>Poulsenia armata</i>	-2.76	-2.91	-2.6	-3.33	-3.65	-3.05	0.33	-0.21	0.83	-0.12	-0.99	0.76
<i>Pourouma bicolor</i>	-3.81	-4.3	-3.39	-3.44	-3.78	-3.11	0.65	-0.36	1.78	0.94	0.05	1.8
<i>Pouteria reticulata</i>	-1.83	-1.89	-1.78	-2.09	-2.16	-2.02	0.78	0.58	0.98	0.91	0.64	1.15
<i>Pouteria stipitata</i>	-3.71	-4.15	-3.3	-3.51	-3.88	-3.14	0.8	-0.27	1.87	1.48	0.56	2.37
<i>Prioria copaifera</i>	-2.45	-2.56	-2.34	-2.83	-2.99	-2.65	0.34	-0.05	0.73	0.14	-0.45	0.68
<i>Protium confusum</i>	-3.97	-4.54	-3.45	-	-	-	-0.15	-1.44	1.07	-	-	-
<i>Protium costaricense</i>	-2.21	-2.29	-2.13	-2.61	-2.76	-2.49	0.8	0.53	1.08	1.18	0.79	1.62
<i>Protium panamense</i>	-1.64	-1.69	-1.6	-1.86	-1.92	-1.81	0.82	0.64	0.98	0.52	0.3	0.74
<i>Protium tenuifolium</i>	-1.63	-1.67	-1.58	-1.96	-2.03	-1.9	0.16	-0.01	0.32	0.45	0.21	0.69
<i>Pseudobombax septenatum</i>	-3.66	-4.07	-3.27	-4.18	-4.88	-3.58	1.79	0.93	2.71	0.66	-0.62	1.82
<i>Psidium friedrichsthalianum</i>	-3.52	-3.86	-3.15	-3.49	-3.85	-3.16	0.48	-0.6	1.36	0.44	-0.55	1.33
<i>Psychotria acuminata</i>	-3.26	-3.57	-3.01	-3.49	-3.86	-3.15	1.43	0.75	2.16	0.66	-0.3	1.53
<i>Psychotria chagensis</i>	-	-	-	-3.93	-4.49	-3.4	-	-	-	0.39	-0.84	1.47
<i>Psychotria deflexa</i>	-2.85	-3.04	-2.68	-3.33	-3.62	-3.05	0.91	0.36	1.51	1.09	0.28	1.91
<i>Psychotria graciliflora</i>	-2.79	-2.99	-2.6	-2.79	-3.05	-2.55	1.24	0.57	1.98	0.57	-0.31	1.42
<i>Psychotria grandis</i>	-2.91	-3.1	-2.74	-3.41	-3.73	-3.1	0.54	-0.04	1.17	0.52	-0.41	1.37
<i>Psychotria hoffmannseggiana</i>	-3.98	-4.59	-3.48	-	-	-	0.71	-0.47	1.85	-	-	-
<i>Psychotria horizontalis</i>	-1.2	-1.23	-1.17	-1.48	-1.52	-1.44	0.03	-0.07	0.15	0.2	0.05	0.34
<i>Psychotria limonensis</i>	-3.23	-3.55	-2.95	-3.26	-3.54	-2.99	1.15	0.27	2.06	0.59	-0.29	1.37
<i>Psychotria marginata</i>	-1.93	-1.99	-1.86	-2.04	-2.12	-1.97	0.7	0.49	0.95	0.87	0.57	1.17
<i>Psychotria pittieri</i>	-4.23	-4.91	-3.61	-	-	-	0.98	-0.19	2.29	-	-	-
<i>Psychotria racemosa</i>	-3.96	-4.55	-3.44	-	-	-	0.41	-0.71	1.65	-	-	-
<i>Pterocarpus rohrii</i>	-1.8	-1.86	-1.75	-2.05	-2.13	-1.99	0.72	0.53	0.91	0.93	0.7	1.18
<i>Quararibea asterolepis</i>	-2.1	-2.18	-2.03	-2.3	-2.4	-2.21	0.1	-0.18	0.39	-0.02	-0.36	0.3
<i>Quassia amara</i>	-3.67	-4.04	-3.27	-3.96	-4.52	-3.42	0.4	-0.55	1.53	1.07	-0.06	2.19
<i>Randia armata</i>	-2.31	-2.4	-2.22	-2.55	-2.67	-2.42	0.56	0.22	0.87	0.56	0.06	0.95
<i>Rauvolfia littoralis</i>	-	-	-	-4.19	-4.88	-3.57	-	-	-	0.86	-0.3	2.13
<i>Rinorea sylvatica</i>	-1.86	-1.92	-1.8	-2.11	-2.2	-2.04	0.43	0.22	0.67	0.29	-0.03	0.6
<i>Sapium glandulosum</i>	-3.5	-3.83	-3.17	-3.85	-4.32	-3.37	1.93	1.22	2.65	1.96	1	2.9
<i>Schizolobium parahyba</i>	-3.85	-4.38	-3.39	-4.17	-4.89	-3.53	0.93	-0.22	1.95	0.7	-0.73	1.82
<i>Senna dariensis</i>	-2.76	-2.93	-2.61	-3.38	-3.71	-3.08	1.39	0.96	1.86	2.33	1.65	2.98
<i>Simarouba amara</i>	-1.76	-1.81	-1.71	-1.89	-1.96	-1.83	0.7	0.52	0.89	0.93	0.66	1.15
<i>Siparuna guianensis</i>	-4.19	-4.9	-3.57	-	-	-	0.4	-0.83	1.67	-	-	-
<i>Siparuna pauciflora</i>	-2.64	-2.77	-2.5	-2.97	-3.16	-2.77	0.38	-0.07	0.8	0.35	-0.33	0.95
<i>Sloanea terniflora</i>	-3.05	-3.26	-2.83	-3.67	-4.07	-3.23	0.78	0.09	1.47	0.73	-0.3	1.71
<i>Socratea exorrhiza</i>	-2.32	-2.41	-2.23	-2.44	-2.56	-2.33	-0.13	-0.47	0.22	-0.2	-0.57	0.21
<i>Solanum asperum</i>	-3.82	-4.33	-3.39	-3.84	-4.34	-3.4	1.49	0.51	2.55	1.95	0.98	2.85
<i>Solanum circinatum</i>	-3.97	-4.59	-3.47	-4.18	-4.84	-3.52	0.5	-0.74	1.64	0.94	-0.26	2.11

Species	Parameters											
	a						b					
	1985-1990			1990-1995			1985-1990			1990-1995		
mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	mean	CI -	CI +	
<i>Solanum hayesii</i>	-2.91	-3.12	-2.73	-3.15	-3.39	-2.9	1.79	1.21	2.36	1.74	1.15	2.42
<i>Solanum steyermarkii</i>	-	-	-	-4.19	-4.88	-3.56	-	-	-	1.03	-0.27	2.18
<i>Sorocea affinis</i>	-1.87	-1.92	-1.82	-2.17	-2.25	-2.09	0.18	-0.02	0.4	0.42	0.13	0.72
<i>Spondias mombin</i>	-3.48	-3.75	-3.23	-3.47	-3.81	-3.16	3.28	2.9	3.68	3.07	2.44	3.68
<i>Spondias radlkoferi</i>	-2.63	-2.79	-2.48	-2.74	-2.91	-2.57	1.7	1.26	2.19	1.98	1.6	2.4
<i>Sterculia apetala</i>	-3.37	-3.67	-3.06	-3.59	-3.97	-3.25	0.95	0.15	1.82	0.93	-0.01	1.92
<i>Stylogyne turbacensis</i>	-2.19	-2.27	-2.11	-2.24	-2.32	-2.16	0.98	0.73	1.26	0.63	0.31	0.95
<i>Swartzia simplex_var.grandiflora</i>	-1.62	-1.66	-1.58	-1.86	-1.92	-1.81	0.28	0.12	0.43	0.06	-0.15	0.27
<i>Symphonia globulifera</i>	-2.97	-3.17	-2.78	-3.06	-3.29	-2.85	-0.04	-0.67	0.57	0.06	-0.63	0.82
<i>Tabebuia guayacan</i>	-3.64	-4.01	-3.25	-3.83	-4.26	-3.32	0.95	-0.08	1.95	1.28	0.16	2.31
<i>Tabebuia rosea</i>	-2.6	-2.73	-2.46	-3.12	-3.33	-2.88	0.73	0.31	1.21	0.85	0.2	1.63
<i>Tabernaemontana arborea</i>	-2	-2.07	-1.94	-2.02	-2.09	-1.94	1.04	0.82	1.26	1.09	0.84	1.34
<i>Tachigali versicolor</i>	-1.51	-1.54	-1.46	-1.72	-1.77	-1.67	0.09	-0.05	0.23	0.54	0.34	0.74
<i>Talisia nervosa</i>	-2.91	-3.1	-2.73	-3.36	-3.65	-3.07	0.47	-0.18	1.1	0.33	-0.49	1.3
<i>Talisia princeps</i>	-2.3	-2.39	-2.22	-2.5	-2.61	-2.38	0.02	-0.3	0.37	0.55	0.16	0.96
<i>Terminalia amazonia</i>	-4.28	-5.01	-3.63	-4.16	-4.89	-3.56	1.32	0.15	2.58	0.43	-0.69	1.76
<i>Terminalia oblonga</i>	-3.52	-3.86	-3.18	-3.42	-3.76	-3.14	1.15	0.22	1.95	0.79	-0.17	1.61
<i>Tetragastris panamensis</i>	-1.44	-1.48	-1.41	-1.64	-1.69	-1.6	0.5	0.36	0.63	0.35	0.19	0.53
<i>Thevetia ahouai</i>	-3.22	-3.77	-2.69	-3.72	-4.18	-3.29	1.43	0.28	2.69	1.35	0.42	2.43
<i>Trattinnickia aspera</i>	-4.03	-4.6	-3.54	-3.84	-4.41	-3.4	1.18	-0.01	2.31	1.38	0.28	2.42
<i>Trema micrantha</i>	-3.66	-4.08	-3.28	-3.28	-3.58	-3.03	1.53	0.62	2.47	1.07	0.26	1.8
<i>Trichanthera gigantea</i>	-4.03	-4.62	-3.48	-3.62	-4.04	-3.25	1.11	-0.02	2.3	1.29	0.32	2.22
<i>Trichilia pallida</i>	-2.43	-2.53	-2.32	-2.63	-2.77	-2.5	0.8	0.44	1.11	0.89	0.42	1.32
<i>Trichilia tuberculata</i>	-1.01	-1.03	-0.98	-1.28	-1.32	-1.25	-0.12	-0.22	-0.03	-0.11	-0.24	0.01
<i>Trichospermum galeottii</i>	-4.23	-4.97	-3.62	-	-	-	1.05	-0.2	2.27	-	-	-
<i>Triplaris cumingiana</i>	-3.1	-3.32	-2.88	-3.3	-3.59	-3.02	0.38	-0.36	1.07	1.13	0.34	1.95
<i>Trophis caucana</i>	-3.32	-3.64	-3.07	-	-	-	0.09	-0.79	0.88	-	-	-
<i>Trophis racemosa</i>	-2.68	-2.83	-2.54	-2.92	-3.11	-2.73	0.18	-0.35	0.65	1.12	0.53	1.67
<i>Turpinia occidentalis</i>	-4.26	-5.02	-3.64	-3.57	-3.93	-3.22	1.22	0.04	2.51	0.73	-0.24	1.72
<i>Unonopsis pittieri</i>	-2.37	-2.47	-2.28	-2.75	-2.91	-2.61	0.08	-0.29	0.44	0.33	-0.23	0.89
<i>Urera baccifera</i>	-3.11	-3.4	-2.84	-3.49	-3.9	-3.13	0.35	-0.44	1.22	1.95	0.96	2.99
<i>Virola multiflora</i>	-3.97	-4.53	-3.41	-4.16	-4.8	-3.52	0.07	-1.1	1.28	0.69	-0.47	1.92
<i>Virola sebifera</i>	-2.58	-2.69	-2.45	-3.31	-3.59	-3.03	0.26	-0.21	0.7	0.67	-0.15	1.54
<i>Virola surinamensis</i>	-3.99	-4.49	-3.45	-	-	-	0.81	-0.27	1.94	-	-	-
<i>Vismia baccifera</i>	-3.05	-3.26	-2.83	-3.04	-3.3	-2.84	0.88	0.19	1.61	0.62	-0.13	1.34
<i>Vismia macrophylla</i>	-	-	-	-4.2	-4.92	-3.58	-	-	-	1.03	-0.09	2.4
<i>Vochysia ferruginea</i>	-3.73	-4.17	-3.36	-3.95	-4.52	-3.44	1.54	0.54	2.47	0.88	-0.24	1.99
<i>Xylopia macrantha</i>	-2.13	-2.21	-2.05	-2.27	-2.36	-2.17	-0.36	-0.66	-0.08	-0.39	-0.71	-0.07
<i>Xylosma oligandra</i>	-3.97	-4.53	-3.44	-4.17	-4.92	-3.55	0.66	-0.47	1.87	0.69	-0.45	1.93
<i>Zanthoxylum acuminatum</i>	-3.7	-4.08	-3.31	-3.53	-3.91	-3.2	1.92	1.11	2.81	1.12	0.19	2.05
<i>Zanthoxylum ekmanii</i>	-2.77	-2.94	-2.57	-2.77	-3	-2.54	2.2	1.68	2.76	1.7	0.88	2.5
<i>Zanthoxylum panamense</i>	-3.07	-3.3	-2.83	-3.54	-3.9	-3.16	2.34	1.71	2.96	2.41	1.7	3.2
<i>Zuelania guidonia</i>	-3.52	-3.89	-3.18	-3.27	-3.53	-2.99	1.44	0.54	2.27	0.81	0.01	1.6

Species	Parameters							
	<i>c</i>						<i>Light50</i>	
	1985-1990			1990-1995			1985-90	1990-95
mean	CI -	CI +	mean	CI -	CI +	mean	mean	
<i>Acacia melanoceras</i>	46	0.025	94	51	0.94	96	0	0.05
<i>Acalypha diversifolia</i>	0.054	0.032	0.083	0.03	0.015	0.047	0.11	0.11
<i>Acalypha macrostachya</i>	47	0.089	94	0.0019	1e-04	0.0052	0.11	0.12
<i>Adelia triloba</i>	0.0058	0.0016	0.011	50	4.3	99	0.14	0.11
<i>Aegiphila panamensis</i>	49	0.97	95	50	2.4	95	0.1	0.13
<i>Alchornea costaricensis</i>	49	3.3	98	0.009	4e-04	0.026	0.14	0.15
<i>Alchornea latifolia</i>	50	1.8	97	-	-	-	0.02	-
<i>Alibertia edulis</i>	0.063	0.0071	0.16	47	0.0088	95	0.04	0.09
<i>Allophylus psilospermus</i>	52	3.8	98	50	0.76	94	0.07	0.09
<i>Alseis blackiana</i>	0.18	0.15	0.21	0.17	0.12	0.22	0.09	0.11
<i>Anaxagorea panamensis</i>	0.0028	0.0018	0.0039	0.0018	0.0011	0.0026	0.16	0.14
<i>Andira inermis</i>	49	1.3	96	50	0.29	94	0.04	0
<i>Annona acuminata</i>	0.1	0.035	0.19	17	0.04	84	0.13	0.11
<i>Annona spraguei</i>	0.025	0.011	0.041	0.33	0.017	1.6	0.14	0.14
<i>Apeiba membranacea</i>	48	0.55	95	50	0.39	95	0.14	0.12
<i>Apeiba tibourbou</i>	50	5.1	100	50	5	100	0.12	0.1
<i>Aphelandra sinclairiana</i>	48	0.017	95	49	0.13	95	0.09	0.11
<i>Appunia seibertii</i>	48	0.14	94	-	-	-	0.08	-
<i>Ardisia bartlettii</i>	-	-	-	47	0.095	94	-	0.08
<i>Ardisia fendleri</i>	49	0.89	95	51	4.4	100	0	0
<i>Ardisia guianensis</i>	45	0.0044	94	48	1.7	96	0.15	0.12
<i>Aspidosperma spruceanum</i>	50	3.1	97	50	6	100	0.03	0.03
<i>Astrocaryum standleyanum</i>	49	0.058	95	-	-	-	0.01	-
<i>Astronium graveolens</i>	0.0079	1e-04	0.035	47	0.089	95	0.12	0.11
<i>Attalea butyracea</i>	48	4.8	99	-	-	-	0.03	-
<i>Bactris barronis</i>	49	0.17	94	-	-	-	0	-
<i>Bactris coloradonis</i>	50	0.12	94	-	-	-	0.04	-
<i>Bactris major</i>	0.0069	0.0014	0.015	48	0.021	95	0.11	0
<i>Banara guianensis</i>	49	0.93	96	51	5.5	100	0.1	0.07
<i>Beilschmiedia pendula</i>	0.065	0.046	0.084	0.12	0.05	0.21	0	0
<i>Brosimum alicastrum</i>	49	0.39	95	50	3.4	98	0	0
<i>Calophyllum longifolium</i>	0.021	0.014	0.028	0.051	0.028	0.08	0.02	0.03
<i>Capparis frondosa</i>	42	0.13	93	52	6.1	100	0.01	0
<i>Casearia aculeata</i>	49	1.9	96	47	0.13	95	0.08	0.09
<i>Casearia arborea</i>	48	0.098	94	50	5.2	99	0.13	0.12
<i>Casearia guianensis</i>	50	1.2	95	50	9e-04	94	0	0.08
<i>Casearia sylvestris</i>	50	0.27	94	50	1.2	95	0.1	0.02
<i>Cassipourea elliptica</i>	0.061	0.027	0.12	27	0.027	91	0.06	0.05
<i>Cavanillesia platanifolia</i>	50	2.5	97	50	5.4	100	0.06	0.08
<i>Cecropia insignis</i>	0.03	0.017	0.045	0.018	0.0096	0.028	0.16	0.14
<i>Cecropia longipes</i>	-	-	-	50	0.048	94	-	0.12
<i>Cecropia obtusifolia</i>	0.028	0.0041	0.072	51	4.8	99	0.14	0.14
<i>Cedrela odorata</i>	50	3.2	97	-	-	-	0.04	-
<i>Ceiba pentandra</i>	50	4.2	98	51	5.8	100	0.06	0.12
<i>Celtis schippii</i>	50	5	100	51	5.4	100	0	0.08
<i>Cespedesia spathulata</i>	51	4.7	99	-	-	-	0	-
<i>Cestrum megalophyllum</i>	47	0.041	95	51	5.2	100	0.08	0.08
<i>Chamaedorea tepejilote</i>	50	4.2	99	49	4.1	99	0.04	0.09
<i>Changuava schippii</i>	0.08	0.016	0.19	1.2	0.012	9.3	0.14	0.13
<i>Chrysochlamys eclipes</i>	50	2.6	97	48	0.22	95	0	0.08
<i>Chrysophyllum argenteum</i>	0.39	0.045	0.94	47	0.19	95	0.09	0.1
<i>Chrysophyllum cainito</i>	51	2.3	96	52	5.5	99	0.11	0.12
<i>Cinnamomum triplinerve</i>	52	3.4	97	50	1.8	96	0.11	0.1
<i>Clidemia dentata</i>	50	4.8	99	46	0.026	93	0.12	0.12
<i>Clidemia octona</i>	45	0.16	94	50	5	100	0.12	0.11
<i>Coccoloba coronata</i>	51	5.9	99	51	5.9	100	0.05	0.05

Species	Parameters							
	<i>c</i>						<i>Light50</i>	
	1985-1990			1990-1995			1985-90	1990-95
mean	CI -	CI +	mean	CI -	CI +	mean	mean	
<i>Coccoloba manzinellensis</i>	50	0.21	94	49	2	96	0.09	0.1
<i>Colubrina glandulosa</i>	-	-	-	50	1.5	96	-	0.11
<i>Conostegia bracteata</i>	49	0.034	94	50	0.17	94	0.08	0.07
<i>Conostegia cinnamomea</i>	0.029	0.008	0.061	27	0.039	90	0.04	0.08
<i>Cordia alliodora</i>	28	0.0067	92	51	2.4	97	0.13	0.09
<i>Cordia bicolor</i>	0.068	0.049	0.087	0.07	0.04	0.11	0.14	0.13
<i>Cordia lasiocalyx</i>	0.15	0.059	0.27	37	0.04	94	0.06	0.05
<i>Coussarea curvigemma</i>	0.26	0.13	0.43	40	0.19	93	0.07	0.04
<i>Croton billbergianus</i>	0.031	0.026	0.038	0.03	0.018	0.045	0.15	0.14
<i>Cupania cinerea</i>	49	0.068	94	-	-	-	0.1	-
<i>Cupania latifolia</i>	50	1.6	96	51	1.5	95	0.03	0.02
<i>Cupania rufescens</i>	0.052	0.0043	0.15	13	0.0029	31	0.13	0.12
<i>Cupania seemannii</i>	42	0.14	94	45	0.077	94	0.08	0.08
<i>Dendropanax arboreus</i>	50	4.8	99	-	-	-	0.07	-
<i>Desmopsis panamensis</i>	1.4	0.87	2	1.2	0.52	2	0	0
<i>Diospyros artanthifolia</i>	49	0.017	96	50	0.15	95	0.12	0.03
<i>Dipteryx oleifera</i>	49	1.5	95	50	0.24	95	0.05	0.1
<i>Drypetes standleyi</i>	47	0.085	94	52	6.4	99	0	0
<i>Elaeis oleifera</i>	50	4.8	99	-	-	-	0.09	-
<i>Erythrina costaricensis</i>	50	2.4	96	49	0.084	95	0.02	0.08
<i>Erythroxylum macrophyllum</i>	48	0.074	94	50	1.9	96	0.08	0.03
<i>Erythroxylum panamense</i>	50	5.9	100	52	6.2	100	0.06	0.06
<i>Eugenia coloradoensis</i>	42	0.064	94	50	3.2	96	0.08	0.03
<i>Eugenia galalonensis</i>	0.18	0.082	0.3	0.25	0.09	0.49	0.03	0.08
<i>Eugenia nesiotica</i>	50	5.9	100	0.14	0.0026	0.57	0.04	0.07
<i>Eugenia oerstediana</i>	0.15	0.095	0.2	0.065	0.036	0.1	0.03	0.05
<i>Faramea occidentalis</i>	0.69	0.59	0.8	0.6	0.46	0.74	0.01	0.01
<i>Ficus bullenei</i>	-	-	-	50	3.2	97	-	0.11
<i>Ficus citrifolia</i>	50	0.12	95	-	-	-	0.11	-
<i>Ficus costaricana</i>	-	-	-	50	1.8	96	-	0.1
<i>Ficus insipida</i>	50	4.4	100	50	2	97	0.07	0.12
<i>Ficus maxima</i>	50	3.1	97	-	-	-	0.11	-
<i>Ficus obtusifolia</i>	52	2.5	95	50	5.6	100	0.04	0.06
<i>Ficus pertusa</i>	50	0.58	95	51	1.2	96	0.08	0.1
<i>Ficus popenoei</i>	51	5.6	100	-	-	-	0.11	-
<i>Ficus trigonata</i>	-	-	-	52	5.2	99	-	0.09
<i>Garcinia intermedia</i>	0.58	0.22	1.1	33	0.16	91	0	0
<i>Garcinia madruno</i>	50	2.8	97	51	0.044	95	0	0
<i>Genipa americana</i>	49	4.6	99	-	-	-	0	-
<i>Guapira standleyana</i>	49	0.18	95	51	4.6	99	0.03	0.03
<i>Guarea fuzzy</i>	47	0.22	93	51	4.5	99	0.05	0.06
<i>Guarea grandifolia</i>	50	4.5	99	49	3.3	97	0	0.1
<i>Guarea guidonia</i>	20	0.14	86	48	0.26	95	0.05	0.05
<i>Guatteria dumetorum</i>	47	0.12	94	47	0.17	95	0.06	0.08
<i>Guazuma ulmifolia</i>	51	0.42	95	50	5.7	100	0.14	0.13
<i>Guettarda foliacea</i>	51	1.4	96	50	1.8	97	0.06	0.03
<i>Gustavia superba</i>	51	0.83	94	47	0.016	95	0.12	0.14
<i>Hamelia axillaris</i>	45	0.11	94	0.059	0.0096	0.15	0.13	0.13
<i>Hampea appendiculata</i>	51	5.4	100	51	5.6	100	0.13	0.11
<i>Hasseltia floribunda</i>	0.021	0.0022	0.063	46	0.097	95	0.08	0.12
<i>Heisteria acuminata</i>	50	0.92	95	51	5.5	100	0.1	0.02
<i>Heisteria concinna</i>	52	5.6	100	49	4.1	98	0	0.06
<i>Herrania purpurea</i>	43	0.062	94	27	0.018	90	0.05	0.11
<i>Hieronyma alchorneoides</i>	49	0.76	95	48	2.8	97	0.08	0.11
<i>Hirtella americana</i>	50	4.7	99	50	5.2	100	0.09	0.08
<i>Hirtella triandra</i>	0.32	0.19	0.47	16	0.12	66	0	0.01

Species	Parameters							
	<i>c</i>						<i>Light50</i>	
	1985-1990			1990-1995			1985-90	1990-95
mean	CI -	CI +	mean	CI -	CI +	mean	mean	
<i>Hybanthus prunifolius</i>	0.67	0.61	0.74	0.51	0.43	0.6	0	0.01
<i>Inga acuminata</i>	49	0.1	95	49	0.6	95	0.07	0.01
<i>Inga cocleensis</i>	50	0.42	95	50	5.1	100	0.13	0.1
<i>Inga goldmanii</i>	44	0.029	94	48	0.006	94	0.06	0.1
<i>Inga laurina</i>	50	3.7	98	50	4.4	98	0.1	0.07
<i>Inga marginata</i>	0.089	0.044	0.14	37	0.055	93	0.09	0.11
<i>Inga nobilis</i>	46	0.17	95	50	0.51	94	0.02	0.01
<i>Inga oerstediana</i>	49	4.3	99	49	0.027	95	0.11	0.06
<i>Inga pezizifera</i>	48	0.044	94	50	4	99	0	0.07
<i>Inga punctata</i>	50	0.67	95	51	1	96	0.12	0.11
<i>Inga ruiziana</i>	50	4.7	99	-	-	-	0.09	-
<i>Inga sapindoides</i>	48	0.11	96	50	0.34	96	0.11	0.13
<i>Inga spectabilis</i>	50	5.1	100	-	-	-	0.09	-
<i>Inga thibaudiana</i>	49	4.8	100	0.025	0.0031	0.065	0.13	0.13
<i>Inga umbellifera</i>	15	0.063	85	49	2.2	96	0.09	0.1
<i>Jacaranda copaia</i>	49	0.88	95	48	0.12	95	0.14	0.13
<i>Koanophyllon wetmorei</i>	48	0.18	93	50	2.8	97	0.1	0.11
<i>Lacistema aggregatum</i>	0.81	0.094	2.4	44	0.11	94	0	0.05
<i>Lacmellea panamensis</i>	50	1	96	51	2.7	97	0.06	0.04
<i>Laetia procera</i>	0.049	4e-04	0.21	49	0.072	95	0.14	0.11
<i>Laetia thannia</i>	48	0.051	95	41	0.075	93	0.1	0.08
<i>Leandra dichotoma</i>	49	2.9	98	-	-	-	0.11	-
<i>Licania hypoleuca</i>	49	0.057	95	51	4	99	0	0.03
<i>Licania platypus</i>	50	1.5	96	50	2.8	96	0	0
<i>Lindackeria laurina</i>	51	4.8	99	47	0.13	94	0.09	0.09
<i>Lonchocarpus heptaphyllus</i>	48	0.19	95	0.19	0.011	0.5	0.09	0.12
<i>Luehea seemanii</i>	27	0.072	91	49	4.8	99	0.14	0.14
<i>Macrocnemum roseum</i>	0.02	1e-04	0.15	50	0.74	95	0	0.1
<i>Malpighia romeroana</i>	50	0.28	95	49	5.1	98	0.06	0.07
<i>Maquira guianensis</i>	5.5	0.042	29	42	0.092	93	0	0
<i>Margaritaria nobilis</i>	50	5.2	99	-	-	-	0.13	-
<i>Maytenus schippii</i>	49	0.7	95	50	1.9	96	0.02	0.03
<i>Miconia affinis</i>	0.028	0.015	0.044	0.045	0.012	0.096	0.13	0.12
<i>Miconia argentea</i>	0.071	0.055	0.087	0.034	0.025	0.043	0.13	0.14
<i>Miconia elata</i>	0.0033	0	0.015	-	-	-	0.1	-
<i>Miconia hondurensis</i>	49	0.056	94	51	3.1	97	0.08	0.04
<i>Miconia impetolaris</i>	50	2.4	96	50	2.7	96	0.01	0.08
<i>Miconia nervosa</i>	0.062	0.039	0.092	0.11	0.029	0.23	0.13	0.12
<i>Miconia prasina</i>	51	6.3	100	-	-	-	0.09	-
<i>Mosannonna garwoodii</i>	51	4.4	99	50	3.9	99	0.02	0
<i>Mouriri myrtilloides</i>	0.86	0.54	1.2	0.29	0.21	0.4	0.06	0.06
<i>Myrcia gatunensis</i>	50	3.1	97	49	0.84	94	0.01	0.05
<i>Nectandra cissiflora</i>	41	0.054	93	51	5	100	0.1	0.11
<i>Nectandra lineata</i>	50	0.87	95	51	0.12	95	0.05	0.09
<i>Nectandra purpurea</i>	50	0.86	95	50	2.4	97	0.08	0.1
<i>Neea amplifolia</i>	51	4.5	99	49	0.088	94	0.06	0.11
<i>Ochroma pyramidale</i>	50	6	100	49	4.4	99	0.11	0.09
<i>Ocotea cernua</i>	47	0.043	96	47	0.024	95	0.06	0.09
<i>Ocotea oblonga</i>	49	0.14	95	48	0.023	95	0.08	0.09
<i>Ocotea puberula</i>	47	0.052	94	50	2.2	96	0.09	0.08
<i>Ocotea whitei</i>	2.5	0.0051	8.7	49	1	95	0	0
<i>Oenocarpus mapora</i>	38	0.061	95	48	0.13	94	0.04	0
<i>Ormosia coccinea</i>	49	1.6	96	50	2.3	97	0	0
<i>Ormosia macrocalyx</i>	49	1.2	95	49	0.083	94	0.05	0.08
<i>Ouatea lucens</i>	50	4.9	98	50	3.3	97	0.07	0.08
<i>Pachira quinata</i>	49	0.79	95	-	-	-	0.08	-

Species	Parameters							
	<i>c</i>						<i>Light50</i>	
	1985-1990			1990-1995			1985-90	1990-95
mean	CI -	CI +	mean	CI -	CI +	mean	mean	
<i>Pachira sessilis</i>	-	-	-	51	0.065	94	-	0.1
<i>Palicourea guianensis</i>	0.091	0.077	0.1	0.051	0.043	0.06	0.14	0.15
<i>Pentagonia macrophylla</i>	50	3.5	98	49	0.2	95	0.07	0.03
<i>Perebea xanthochyma</i>	48	0.019	95	51	5.4	100	0.05	0.05
<i>Picramnia latifolia</i>	39	0.1	94	43	0.13	94	0.01	0.01
<i>Piper aequale</i>	49	0.29	94	50	0.23	94	0	0.09
<i>Piper arboreum</i>	49	2.6	97	2.6	1e-04	9.8	0.07	0.12
<i>Piper colonense</i>	51	5.6	100	50	1.5	96	0.04	0.06
<i>Piper cordulatum</i>	0.056	0.031	0.083	48	0.096	95	0.07	0.1
<i>Piper perlasense</i>	51	4.6	99	50	0.71	95	0	0.06
<i>Piper reticulatum</i>	51	4.5	100	51	3.9	98	0.08	0.05
<i>Platymiscium pinnatum</i>	51	5.5	100	43	0.015	95	0.11	0.13
<i>Platyopodium elegans</i>	51	5	99	47	0.077	94	0.1	0.1
<i>Posoqueria latifolia</i>	48	0.28	95	51	5.4	100	0.09	0
<i>Poulsenia armata</i>	49	2.9	96	50	1.4	96	0.02	0
<i>Pourouma bicolor</i>	51	0.8	95	47	0.026	95	0.07	0.1
<i>Pouteria reticulata</i>	0.49	0.12	1.1	52	7.5	99	0.08	0.09
<i>Pouteria stipitata</i>	50	1.5	96	50	1.6	95	0.08	0.13
<i>Prioria copaifera</i>	0.25	0.011	1	49	0.96	95	0.03	0
<i>Protium confusum</i>	50	2.1	96	-	-	-	0	-
<i>Protium costaricense</i>	32	0.069	93	45	0.074	94	0.08	0.11
<i>Protium panamense</i>	0.23	0.14	0.34	0.24	0.075	0.48	0.09	0.05
<i>Protium tenuifolium</i>	0.16	0.1	0.23	0.12	0.047	0.22	0	0.04
<i>Pseudobombax septenatum</i>	50	0.55	95	49	3.1	98	0.14	0.07
<i>Psidium friedrichsthalianum</i>	50	2.8	98	50	1.2	96	0.05	0.04
<i>Psychotria acuminata</i>	46	0.024	95	50	5.9	100	0.12	0.07
<i>Psychotria chagensis</i>	-	-	-	51	4.7	100	-	0.03
<i>Psychotria deflexa</i>	47	0.012	94	51	3.4	99	0.09	0.11
<i>Psychotria graciliflora</i>	0.0043	0.0014	0.008	0.0017	5e-04	0.0035	0.11	0.06
<i>Psychotria grandis</i>	48	0.052	94	43	0.001	94	0.05	0.05
<i>Psychotria hoffmannseggiana</i>	50	5.4	100	-	-	-	0.07	-
<i>Psychotria horizontalis</i>	0.23	0.18	0.27	0.12	0.086	0.15	0	0.01
<i>Psychotria limonensis</i>	0.0055	3e-04	0.017	15	0.0061	84	0.11	0.06
<i>Psychotria marginata</i>	0.19	0.075	0.35	0.037	0.021	0.057	0.07	0.09
<i>Psychotria pittieri</i>	50	3.3	98	-	-	-	0.1	-
<i>Psychotria racemosa</i>	49	3.3	97	-	-	-	0.04	-
<i>Pterocarpus rohrii</i>	0.93	0.13	2.6	34	0.16	91	0.08	0.09
<i>Quararibea asterolepis</i>	0.19	0.035	0.48	47	0.1	95	0	0
<i>Quassia amara</i>	50	2.5	97	50	0.009	93	0.04	0.1
<i>Randia armata</i>	51	1.6	95	46	0.089	95	0.06	0.06
<i>Rauvolfia littoralis</i>	-	-	-	48	1.9	95	-	0.09
<i>Rinorea sylvatica</i>	0.06	0.036	0.087	0.038	0.017	0.062	0.04	0.02
<i>Sapium glandulosum</i>	51	4.7	99	48	0.0025	95	0.14	0.14
<i>Schizolobium parahyba</i>	52	2.9	98	50	4.8	100	0.1	0.07
<i>Senna dariensis</i>	50	5.2	100	49	0.15	94	0.12	0.15
<i>Simarouba amara</i>	0.16	0.096	0.25	0.088	0.049	0.14	0.07	0.09
<i>Siparuna guianensis</i>	50	0.057	94	-	-	-	0.04	-
<i>Siparuna pauciflora</i>	50	0.059	95	48	0.053	94	0.03	0.03
<i>Sloanea terniflora</i>	50	3.4	97	50	2.7	97	0.08	0.08
<i>Socratea exorrhiza</i>	18	0.02	67	35	0.057	93	0	0
<i>Solanum asperum</i>	52	4.1	99	51	3.2	97	0.13	0.14
<i>Solanum circinatum</i>	51	3.6	97	50	1.9	96	0.05	0.1
<i>Solanum hayesii</i>	0.079	0.0059	0.24	50	0.26	95	0.14	0.13
<i>Solanum steyermarkii</i>	-	-	-	51	4.3	99	-	0.1
<i>Sorocea affinis</i>	43	0.29	95	42	0.043	94	0	0.04
<i>Spondias mombin</i>	32	0.092	92	0.11	0.02	0.25	0.16	0.16

Species	Parameters							
	<i>c</i>						<i>Light50</i>	
	1985-1990			1990-1995			1985-90	1990-95
mean	CI -	CI +	mean	CI -	CI +	mean	mean	
<i>Spondias radlkoferi</i>	0.046	0.014	0.094	46	0.13	94	0.13	0.14
<i>Sterculia apetala</i>	50	0.81	95	50	1.7	97	0.1	0.1
<i>Stylogyne turbacensis</i>	47	0.14	95	49	2	97	0.1	0.07
<i>Swartzia simplex_var.grandiflora</i>	48	0.77	94	25	0.14	88	0.02	0
<i>Symphonia globulifera</i>	50	0.23	95	48	0.064	95	0	0
<i>Tabebuia guayacan</i>	49	2	96	50	5	99	0.1	0.12
<i>Tabebuia rosea</i>	48	0.059	95	51	5.3	99	0.08	0.09
<i>Tabernaemontana arborea</i>	48	0.34	94	0.23	0.066	0.51	0.1	0.11
<i>Tachigali versicolor</i>	0.19	0.12	0.25	0.085	0.054	0.12	0	0.06
<i>Talisia nervosa</i>	23	0.012	89	49	5.2	99	0.05	0.02
<i>Talisia princeps</i>	44	0.041	94	49	4.7	100	0	0.06
<i>Terminalia amazonia</i>	50	8e-04	94	50	4.7	99	0.12	0.04
<i>Terminalia oblonga</i>	51	3.1	97	49	4.4	98	0.11	0.08
<i>Tetragastris panamensis</i>	0.22	0.16	0.3	0.14	0.085	0.21	0.05	0.03
<i>Thevetia ahouai</i>	0.00036	1e-04	9e-04	46	0.026	95	0.12	0.12
<i>Trattinnickia aspera</i>	50	4.7	99	50	2.4	97	0.11	0.12
<i>Trema micrantha</i>	49	0.24	96	44	0.015	94	0.13	0.1
<i>Trichanthera gigantea</i>	50	0.91	95	45	0.051	95	0.11	0.12
<i>Trichilia pallida</i>	50	2.7	96	50	0.7	95	0.08	0.09
<i>Trichilia tuberculata</i>	0.21	0.19	0.25	0.17	0.14	0.21	0	0
<i>Trichospermum galeottii</i>	51	0.14	95	-	-	-	0.1	-
<i>Triplaris cumingiana</i>	51	5.5	98	51	3.4	98	0.03	0.11
<i>Trophis caucana</i>	50	0.86	95	-	-	-	0	-
<i>Trophis racemosa</i>	51	1.5	95	51	2.7	97	0	0.11
<i>Turpinia occidentalis</i>	48	0.46	93	47	0.0023	94	0.11	0.08
<i>Unonopsis pittieri</i>	50	0.073	95	50	4.8	99	0	0.03
<i>Urera baccifera</i>	0.004	2e-04	0.012	0.0044	1e-04	0.014	0.03	0.14
<i>Virola multiflora</i>	51	5.5	100	50	4.3	98	0	0.07
<i>Virola sebifera</i>	49	4	98	50	5.3	99	0.01	0.07
<i>Virola surinamensis</i>	51	1.7	96	-	-	-	0.09	-
<i>Vismia baccifera</i>	9.5	0.014	72	0.025	5e-04	0.11	0.09	0.07
<i>Vismia macrophylla</i>	-	-	-	50	4	98	-	0.1
<i>Vochysia ferruginea</i>	50	5.2	98	49	1.4	95	0.13	0.09
<i>Xylopia macrantha</i>	0.1	0.03	0.22	22	0.031	87	0	0
<i>Xylosma oligandra</i>	51	1.8	96	50	0.12	95	0.07	0.07
<i>Zanthoxylum acuminatum</i>	50	0.26	95	51	3.8	98	0.14	0.11
<i>Zanthoxylum ekmanii</i>	0.031	0.013	0.055	0.0022	9e-04	0.004	0.15	0.13
<i>Zanthoxylum panamense</i>	0.054	0.0091	0.13	47	0.027	95	0.15	0.15
<i>Zuelania guidonia</i>	52	5.4	99	50	4.9	100	0.12	0.08