

Continue

Família lamiaceae pdf

Harley RM, Atkins S, Budantsev AL, Cantino PD, Conn BJ, Grayer R, et al. Labiatae. In: Kadereit JW, editor.
The families and genera of vascular plants, vol. 7. Berlin: Springer Verlag; 2004. p. 167–275. Google Scholar Olmstead RG. A synoptical classification of the Lamiales, version 2.6.2. 2016. . Accessed 25 Feb 2020. (last updated 12 April, 2016). Google Scholar APG IV. An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Bot J Linn Soc. 2016;181:1–20.Article Google Scholar Bentham G. Verbenaceae and Labiatae. In: Bentham G, Hooker JD, editors. Genera plantarum. London: Reeve; 1876. p. 1131–223. Google Scholar Briquet J. Verbenaceae, Labiatae. In: HGA E, KAE P, editors. Die Natürlichen Pflanzenfamilien. Berlin: Engelmann, W; 1897. p. 132–375. Google Scholar Erdtman G. Pollen morphology and plant taxonomy. IV. Labiatae, Verbenaceae and Avicenniaceae. Svensk Bot Tidskr. 1945;39:277–85. Google Scholar Wunderlich R. Ein Vorschlag zu einer natürlichen Gliederung der Labiaten auf Grund der Pollenkörner, der Samenentwicklung und des reifen Samens. Oester Bot Zeit.

1967;114:383–483.Article Google Scholar Cantino PD, Sanders RW. Subfamilial classification of Labiatae.
Syst Bot. 1986;11:163–85.Article Google Scholar Cantino PD. Evidence for a polyphyletic origin of the Labiatae. Ann Mo Bot Gard. 1992;79:361–79.Article Google Scholar Cantino PD. Toward a phylogenetic classification of the Labiatae. In: Harley RM, Reynolds T, editors. Advances in Labiatae Science. London: Royal Botanic Gardens, Kew; 1992. p. 27–37. Google Scholar Cantino PD, Harley RM, Wagstaff SJ. Genera of Labiatae: status and classification. In: Harley R, Reynolds T, editors. Advances in Labiatae science. London: Royal Botanic Gardens, Kew; 1992. p.

511–22. Google Scholar Junell S. Zur Gynäzeummorphologie und Systematik der Verbenaceen und Labiaten, nebst Bemerungen über ihre Samenentwicklung. Symb Bot Upsal. 1934;4:1–129. Google Scholar Kadereit JW. Lamiales: introduction and conspectus. In: Kadereit JW, editor. The Families and Genera of Vascular Plants, Vol. 7. Berlin: Springer Verlag; 2004. p. 1–8. Google Scholar Wink M, Kaufmann M. Phylogenetic relationships between some members of the subfamily Lamioideae (Family Labiatae) inferred from nucleotide suquences of the rbcL gene. Bot Acta. 1996;109:139–48.Article CAS Google Scholar Wagstaff SJ, Olmstead RG. Phylogeny of Labiatae and Verbenaceae inferred from rbcL sequences. Syst Bot. 1997;22:165–79.Article Google Scholar Wagstaff SJ, Hickerson L, Spangler R, Reeves PA, Olmstead RG. Phylogeny in Labiatae s.l., inferred from cpDNA sequences. Plant Syst Evol. 1998;209:265–74.Article CAS Google Scholar Schäferhoff B, Fleischmann A, Fischer E, Albach DC, Borsch T, Heubl G, et al. Towards resolving Lamiales relationships: insights from rapidly evolving chloroplast sequences. BMC Evol Biol. 2010;10:352.Article PubMed PubMed Central CAS Google Scholar Refulio-Rodriguez NF, Olmstead RG. Phylogeny of Lamiidae. Am J Bot. 2014;101:287–99.Article PubMed Google Scholar Li B, Cantino PD, Olmstead RG, Bramley GL, Xiang C-L, Ma Z-H, et al. A large-scale chloroplast phylogeny of the Lamiaceae sheds new light on its subfamilial classification. Sci Rep. 2016;6:34343.Article CAS PubMed PubMed Central Google Scholar Mint Evolutionary Genomics Consortium. Phylogenomic mining of the mints reveals multiple mechanisms contributing to the evolution of chemical diversity in Lamiaceae. Mol Plant.

2018;11:1084–96.Article CAS Google Scholar Soltis DE, Smith SA, Cellinese N, Wurdack KJ, Tank DC, Brockington SF, et al. Angiosperm phylogeny: 17 genes, 640 taxa. Am J Bot. 2011;98:704–30.Article PubMed Google Scholar Liu B, Tan Y-H, Liu S, Olmstead R, Ming D-Z, Chen Z-D, et al. Phylogenetic relationships of Cyrtandromoea and Wightia revisited: a new tribe in Phrymaceae and a new family in Lamiales. J Syst Evol. 2020;58:1–17.Article CAS Google Scholar Paton AJ, Springate D, Suddee S, Otieno D, Grayer RJ, Harley MM, et al. Phylogeny and evolution of basils and allies (Ocimeae, Labiatae) based on three plastid DNA regions. Mol Phylogenet Evol. 2004;31:277–99.Article CAS PubMed Google Scholar Conn BJ, Streiber N, Brown EA, Heywood MJ, Olmstead RG. Infrageneric phylogeny of Chloantheae (Lamiaceae) based on chloroplast ndhF and nuclear ITS sequence data. Aust Syst Bot. 2009;22:243–56.Article Google Scholar Bräuchler C, Meimberg H, Heubl G. Molecular phylogeny of Menthinae (Lamiaceae, Nepetoideae, Mentheae) – taxonomy, biogeography and conflicts. Mol Phylogenet Evol. 2010;55:501–23.Article PubMed CAS Google Scholar Pastore JFB, Harley RM, Forest F, Paton AJ, Van den Berg C. Phylogeny of the subtribe Hyptidinae (Lamiaceae tribe Ocimeae) as inferred from nuclear and plastid DNA. Taxon. 2011;62:1217–329.
Google Scholar Drew BT, Sytsma KJ.
Phylogenetics, biogeography, and staminal evolution in the tribe Mentheae (Lamiaceae). Am J Bot. 2012;99:933–53.Article PubMed Google Scholar Salmaki Y, Zarre S, Ryding O, Lindqvist C, Scheunert A, Bräuchler C, et al. Phylogeny of the tribe Phlomideae (Lamioideae: Lamiaceae) with special focus on Eremostachys and Phlomoides: new insights from nuclear and chloroplast sequences. Taxon. 2012;65:161–79.Article Google Scholar Salmaki Y, Zarre S, Ryding O, Lindqvist C, Bräuchler C, Heubl G, et al. Molecular phylogeny of tribe Stachydeae (Lamiaceae subfamily Lamioideae). Mol Phylogenet Evol. 2013;69:535–51.Article PubMed Google Scholar Roy T, Chang T-H, Lan T, Lindqvist C. Phylogeny and biogeography of New World Stachydeae (Lamiaceae) with emphasis on the origin and diversification of Hawaiian and South American taxa. Mol Phylogenet Evol. 2013;69:218–38.Article CAS PubMed Google Scholar Chen Y-P, Drew BT, Li B, Soltis DE, Soltis PS, Xiang C-L. Resolving the phylogenetic position of Ombrocharis (Lamiaceae), with reference to the molecular phylogeny of tribe Elsholtzieae. Taxon.



2016;65:123–36.Article Google Scholar Paton AJ, Mwanyambo M, Culham A. Phylogenetic study of Plectranthus, Coleus and allies (Lamiaceae): taxonomy, distribution and medicinal use. Bot J Linn Soc.

2018;188:355–76. Google Scholar Siadati S, Salmaki Y, Mehrvarz SS, Heubl G, Weigned M. Untangling the generic boundaries in tribe Marrubieae (Lamiaceae: Lamioideae) using nuclear and plastid DNA sequences. Taxon. 2018;67:770–83.Article Google Scholar Steane DA, Scotland RW, Mabberley DJ, Wagstaff SJ, Reeves PA, Olmstead RG. Phylogenetic relationships of Clerodendrum s.l. (Lamiaceae) inferred from chloroplast DNA. Syst Bot. 1997;22:229–43.Article Google Scholar Steane DA, Scotland RW, Mabberley DJ, Olmstead RG. Molecular systematics of Clerodendrum (Lamiaceae): ITS sequences and total evidence. Am J Bot. 1999;86:98–107.Article CAS PubMed Google Scholar Steane DA, De Kok RPJ, Olmstead RG. Phylogenetic relationships between Clerodendrum (Lamiaceae) and other Ajugoid genera inferred from nuclear and chloroplast DNA sequences data. Mol Phylogenet Evol. 2004;32:39–45.Article CAS PubMed Google Scholar Barber JC, Francisco-Ortega J, Santos-Guerra A, Turner KG, Jansen RK. Origin of Macaronesian Sideritis L. (Lamioideae: Lamiaceae) inferred from nuclear and chloroplast sequence datasets. Mol Phylogent Evol. 2002;23:293–306.Article CAS Google Scholar Barber JC, Finch CC, Francisco-Ortega J, Santos-Guerra A, Jansen RK. Hybridization in Macaronesian Sideritis (Lamiaceae): Evidence from incongruence of multiple independent nuclear and chloroplast sequence datasets. Taxon. 2007;56:74–88. Google Scholar Lindqvist C, Albert VA. Origin of the Hawaiian endemic mints within the North American Stachys (Lamiaceae). Am J Bot. 2002;89:1709–24.Article PubMed Google Scholar Walker JB, Sytsma KJ, Treutlein J, Wink M. Salvia (Lamiaceae) is not monophyletic: implications for the systematics, radiation, and ecological specializations of Salvia and tribe Mentheae.
Am J Bot. 2004;91:1115–25.Article PubMed Google Scholar Walker JB, Drew BT, Sytsma KJ.
Unravelling species relationships and diversification within the iconic California Floristic Province sages (Salvia subgenus Audibertia, Lamiaceae). Syst Bot. 2015;40:826–44.Article Google Scholar Bramley GLC, Forest F, De Kok RPJ. Troublesome tropical mints: Re-examining generic limits of Vitex and relations (Lamiaceae) in South East Asia. Taxon. 2009;58:500–10.Article Google Scholar Xiang C-L, Zhang Q, Scheen A-C, Cantino PD, Funamoto T, Peng H. Molecular phylogenetics of Chelonopsis (Lamiaceae: Gomphostemmateae) as inferred from nuclear and plastid DNA and morphology. Taxon. 2013;62:375–86.Article Google Scholar Chen Y-P, Li B, Olmstead RG, Cantino PD, Xiang C-L, Liu E-D. Phylogenetic placement of the enigmatic genus Holocheila (Lamiaceae) inferred from plastid DNA sequences. Taxon. 2014;63:355–66.Article Google Scholar Drew BT, Cacho NI, Sytsma KJ.
The transfer of two rare monotypic genera, Neoeplingia and Chaunostoma, to Lepechinia (Lamiaceae), and notes on their conservation. Taxon.
2014;63:831–42.Article Google Scholar Drew BT, González-Gallegos JG, Xiang C-L, Kriebel R, Drummond CP, Walker JB, et al. Salvia united: The greatest good for the greatest number. Taxon. 2017;66:133–45.Article Google Scholar González-Gallegos JG, Drew B, Cuevas-Guzmán R. Novelties on the distribution of Lepechinia flammea (Lamiaceae), rediscovery of L. glomerata in Jalisco, Mexico, and their phylogenetic position within Lepechinia. Bot Sci. 2015;93:1–14.Article Google Scholar Salmaki Y, Kattari S, Heubl G, Bräuchler C. Phylogeny of non-monophyletic Teucrium (Lamiaceae: Ajugoideae): implications for character evolution and taxonomy. Taxon. 2016;65:802–22.Article Google Scholar Will M, Claßen-Bockhoff R. Time to split Salvia s.l. (Lamiaceae)–New insights from Old World Salvia phylogeny. Mol Phylogenet Evol.
2017;109:33–58.Article PubMed Google Scholar Hu G-X, Takano A, Drew BT, Liu E-D, Soltis DE, Soltis PS, et al. Phylogeny and staminal evolution of Salvia (Lamiaceae, Nepetoideae) in East Asia. Ann Bot. 2018;122:649–68.Article PubMed PubMed Central Google Scholar Li B, Olmstead RG. Two new subfamilies in Lamiaceae. Phytotaxa.
2017;313:222–6.Article Google Scholar Scheen A-C, Bendiksby M, Ryding O, Mathiesen C, Albert VA, Lindqvist C. Molecular phylogenetics, character evolution and suprageneric classification of Lamioideae (Lamiaceae). Ann Mo Bot Gard. 2010;97:191–219.Article Google Scholar Bendiksby M, Thorbek L, Scheen A-C, Lindqvist C, Ryding O. An updated phylogeny and classification of Lamiaceae subfamily Lamioideae. Taxon. 2011;60:471–84.Article Google Scholar Xiang C-L, Zhao F, Cantino PD, Drew BT, Li B, Liu E-D, et al. Molecular systematics of Caryopteris (Lamiaceae) and its allies with reference to the molecular phylogeny of subfamily Ajugoideae. Taxon. 2018;67:376–94.Article Google Scholar Moore MJ, Bell CD, Soltis PS, Soltis DE. Using plastid genome-scale data to resolve enigmatic relationships among basal angiosperms. Proc Natl Acad Sci U S A. 2007;104:19363–8.Article PubMed PubMed Central Google Scholar Olmstead RG, Bedoya AM. Whole genomes: the holy grail. A commentary on: 'molecular phylogenomics of the tribe Shoreeae (Dipterocarpaceae) using whole plastidgenomes'. Ann Bot.
2019;123:iv–v.Article CAS PubMed PubMed Central Google Scholar Lin C-P, Huang J-P, Wu C-S, Hsu C-Y, Chaw S-M. Comparative chloroplast genomics reveals the evolution of Pinaceae genera and subfamilies. Genome Biol Evol. 2010;2:504–17.Article PubMed PubMed Central CAS Google Scholar Moore MJ, Soltis PS, Bell CD, Burleigh JG, Soltis DE. Phylogenetic analysis of 83 plastid genes further resolves the early diversification of eudicots. Proc Natl Acad Sci U S A. 2010;107:4623–8.Article CAS PubMed PubMed Central Google Scholar Xi Z, Ruhfel BR, Schaefer H, Amorim AM, Sugumaran M, Wurdack KJ, et al. Phylogenomics and a posteriori data partitioning resolve the Cretaceous angiosperm radiation Malpighiales. Proc Natl Acad Sci U S A.
2012;109:17519–24.Article CAS PubMed PubMed Central Google Scholar Stull GW, De Stefano RD, Soltis DE, Soltis PS. Resolving basal lamiid phylogeny and the circumscription of Icacinaceae with a plastome-scale data set. Am J Bot. 2015;102:1794–813.Article CAS PubMed Google Scholar Zhang S-D, Jin J-J, Chen S-Y, Chase MW, Soltis DE, Li H-T, et al. Diversification of Rosaceae since the late Cretaceous based on plastid phylogenomics. New Phytol. 2017;214:1355–67.Article CAS PubMed Google Scholar Zhang R, Wang Y-H, Jin J-J, Stull GW, Bruneau A, Cardoso D, et al.
Exploration of plastid phylogenomic conflict yields new insights into the deep relationships of Leguminosae. Syst Biol.
2020;69:613–22.Article PubMed PubMed Central Google Scholar Yu X-Q, Gao L-M, Soltis DE, Soltis PS, Yang J-B, Fang L, et al. Insights into the historical assembly of East Asian subtropical evergreen broadleaved forests revealed by the temporal history of the tea family. New Phytol.
2017;215:1235–48.Article CAS PubMed Google Scholar Yang X-Y, Wang Z-F, Luo W-C, Guo X-Y, Zhang C-H, Liu J-Q, et al. Plastomes of Betulaceae and phylogenetic implication. J Syst Evol. 2019;57:508–18.Article Google Scholar Xiang C-L, Dong H-J, Landrein S, Zhao F, Yu W-B, Soltis DE, et al. Revisiting the phylogeny of Dipsacales: new insights from phylogenomic analyses of complete plastomic sequences. J Syst Evol. 2020;58:103–17.Article Google Scholar Zhao F, Li B, Drew BT, Chen Y-P, Wang Q, Yu W-B, et al. Leveraging plastomes of comparative analysis and phylogenomic inference within Scutellarioideae (Lamiaceae). Plos One.
2020;15:e0232602.Article CAS PubMed PubMed Central Google Scholar Zhao F, Drew BT, Chen Y-P, Hu G-X, Li B, Xiang C-L.
The chloroplast genome of Salvia species: genomic characterization and phylogenetic analysis. Int J Plant Sci.
2020;181:812–30.Article Google Scholar Jin J-J, Yu W-B, Yang J-B, Song Y, de Pamphilis CW, Yi T-S, et al. GetOrganelle: a fast and versatile toolkit for accurate de novo assembly of organelle genomes. Genome Biol. 2020;21:241.Article PubMed PubMed Central Google Scholar Roy T, Lindqvist C. New insights into evolutionary relationships within the subfamily Lamioideae (Lamiaceae) based on pentatricopeptide repeat (PPR) nuclear DNA sequences. Am J Bot. 2015;102:1721–35.Article CAS PubMed Google Scholar Guignard ML. Recherches sur la développement de la graine et en particulier du tégument seminal: Labiées. J Bot. 1893;7:241–50. Google Scholar Olmstead RG, Reeves PA, Lepschi BJ. Confirmation of a monophyletic Chloanthoideae (Lamiaceae) comprising tribes Chloantheae and Prostanthereae. Lamiales Newsletter. 1998;6:7–10. Google Scholar Godden GT, Kinser TJ, Soltis PS, Soltis DE. Phylotranscriptomic analyses reveal asymmetrical gene duplication dynamics and signatures of ancient polyploidy in mints. Genome Biol Evol. 2019;11:3393–408.CAS PubMed PubMed Central Google Scholar Conn BJ, Henwood MJ, Streiber N. Synopsis of the tribe Chloantheae and new nomenclatural combinations in Pityrodia s.lat. (Lamiaceae).
Aust Syst Bot. 2011;24:1–9.Article Google Scholar Munir AA. A taxonomic revision of Chloanthaceae trib. Physopsideae Brunonia. 1978;1:407–692. Google Scholar Brown R. Prodromus Florae Novae Hollandiae. London: J. Johnson & Co; 1810.
Google Scholar Reichenbach HGL. Conspectus regni vegetabilis per gradus naturales evoluti. Leipzig: Carolum Cnobloch; 1828.Book Google Scholar Von Mueller FJH. Nova genera et species aliquot rariores in plagis Australiae. Hooker's J Bot Kew Gard Misc. 1857;9:14–24. Google Scholar Hutchinson J. The Families of Flowering Plants. 2nd ed. London: Oxford University Press; 1959. Google Scholar Munir AA. A taxonomic revision of the genus Hemiphora (Chloanthaceae). J Adel Bo Gard. 1978;1:161–6. Google Scholar Eichler H. Supplement to J.M. Black's Flora of South Australia. 2nd edition. Adelaide: Govt. Printer; 1965. Google Scholar Munir AA. A taxonomic revision of the genus Chloanthes (Chloanthaceae). J Adel Bo Gard. 1977;1:83–106. Google Scholar Munir AA. A taxonomic revision of the genus Cyanostegia (Chloanthaceae). Brunonia. 1978;1:45–67. Google Scholar Munir AA.
A taxonomic revision of the genus Pityrodia (Chloanthaceae).
J Adel Bo Gard. 1979;2:1–138. Google Scholar Blackall WE, Grieve BJ. How to know Western Australian wildflowers: a key to the flora of the temperate regions of Western Australia. Nedlands: University of Western Australia Press; 1965. Google Scholar Cronquist A.
The evolution and classification of flowering plants. Aylesbury: Hazel Watson & Viney; 1968. Google Scholar Wilson TC, Radunz EA, Chen S, Conn BJ, Henwood MJ. A new genus and species for Chloantheae (Lamiaceae). Aust Syst Bot. 2020; In press.Guerin GR. Evidence for polyphyly in Hemigenia and Microcorys (Lamiaceae: Westringieae). Aust Syst Bot. 2008;21:313–25.Article Google Scholar Huck RB. Overview of pollination biology in the Lamiaceae.
In: Harley RM, Reynolds T, editors. Advances in Labiatae Science. London: Royal Botanic Gardens, Kew; 1992. p. 167–81. Google Scholar Wilson TC, Conn BJ, Henwood MJ. Great expectations: correlations between pollinator assemblages and floral characters in Lamiaceae.
Int J Plant Sci. 2017;178:170–87.Article Google Scholar Wilson TC, Conn BJ, Henwood MJ. Molecular phylogeny and systematics of Prostanthera (Lamiaceae). Aust Syst Bot. 2012;25:341–52.Article Google Scholar Bentham G. Labiatarum genera et species. London: J. Ridgeway & Sons; 1834. Google Scholar Guerin GR. Nutlet morphology in

Hemigenia R.Br. and Microcorys R.Br. (Lamiaceae). Plant Syst Evol. 2005;254:49–68.Article Google Scholar Hedge IC. Labiatae.
In: Ali SI, Nasir YJ, editors. Flora of Pakistan, vol. 192. Karachi: University of Karachi; 1990. p.
310. Google Scholar Kaufmann M, Wink M. Molecular systematics of the Nepetoideae (Lamiaceae): Phylogenetic implications from rbcL gene sequences. Z Naturforsch C. 1994;49:635–45.Article CAS PubMed Google Scholar Wagstaff SJ, Olmstead RG, Cantino PD. Parsimony analysis of cpDNA restriction site variation in subfamily Nepetoideae (Labiatae).
Am J Bot.
1995;82:886–92.Article CAS Google Scholar Martin AC. The comparative internal morphology of seeds. Ame Mid Natur. 1946;36:513–660.Article Google Scholar Ryding O. The distribution and evolution of myxocarpy in Lamiaceae. In: Harley RM, Reynolds T, editors. Advances in Labiatae Science. London: Royal Botanic Gardens, Kew; 1992. p. 85–96.
Google Scholar Li P, Qi Z-C, Liu L-X, Ohi-Toma T, Lee J, Hsieh TH, et al.
Molecular phylogenetics and biogeography of the mint tribe Elsholtzieae (Nepetoideae, Lamiaceae), with an emphasis on its diversification in East Asia. Sci Rep. 2017;7:2057.Article PubMed PubMed Central CAS Google Scholar Sanders RW, Cantino PD. Nomenclature of the subdivisions of the Lamiaceae. Taxon. 1984;33:64–72.Article Google Scholar Funamoto T, Xiang C-L, Ogawa M, Peng H. A comparative study of chromosome characters in four species of Elsholtzia Will.
(Lamiaceae) in Japan and China. Chromosome Bot. 2012;7:119–23.Article Google Scholar Mayta-Anco LF, Molinari-Novoa EA, Raju AJS.
A new genus within the Elsholtziae (Lamiaceae) from southeast Asia. Weberbauerella. 2016;1:1–7. Google Scholar Paton AJ, Ryding O. Hanceola, Siphocranion and Isodon and their position in the Ocimeae (Labiatae). Kew Bull. 1998;53:723–31.Article Google Scholar Harley RM, Pastore JFB. A generic revision and new combinations in the Hyptidinae (Lamiaceae), based on molecular and morphological evidence. Phytotaxa. 2012;58:1–55.Article Google Scholar Paton AJ, Mwanyambot M, Govaerts RHA, Smitha K, Suddee S, Phillipson PB, et al. Nomenclatural changes in Coleus and Plectranthus (Lamiaceae): a tale of more than two genera. PhytoKeys. 2019;129:1–158.Article PubMed PubMed Central Google Scholar Harley RM, Paton AJ, Ryding O.
New synonym and taxonomic changes in Laibatae. Kew Bull. 2003;58:485–9.Article Google Scholar Zhong J-S, Li J, Li L, Conran JG, Li H-W. Phylogeny of Isodon (Schrad. ex Benth.) Spach (Lamiaceae) and related genera inferred from nuclear ribosomal ITS, trnL-trnF region, and rps16 intron sequences and morphology. Syst Bot. 2010;35:207–19.Article Google Scholar Chen Y-P, Zhu X-X, Zhao F, Feng H-Z, Paton A, Xiang C-L. Molecular and morphological evidence for a new species of Siphocranion (Lamiaceae) from the Sino-Vietnamese border. Phytotaxa. 2019;425:1–18.Article Google Scholar Walker JB, Sytsma KJ. Staminal evolution in the genus Salvia (Lamiaceae): molecular phylogenetic evidence for multiple origins of the staminal lever. Ann Bot. 2007;100:375–91.Article CAS PubMed Google Scholar Drew BT, Sytsma KJ. Testing the monophyly and placement of Lepechinia in the tribe Mentheae (Lamiaceae). Syst Bot. 2011;36:1038–49.Article Google Scholar Drew BT. Phylogenetics and biogeography of Lepechinia (Lamiaceae), and evolutionary studies within the Mentheae Tribe. Ph.D. dissertation. Madison: University of Wisconsin; 2011. Google Scholar Rouy G. Compectus de la Flore de France. Paris: Rue de tournon; 1927. Google Scholar Moon HK, Vinckier S, Smets E, Huysmans S. Palynological evolutionary trends within the tribe Mentheae with special emphasis on subtribe Menthinae (Nepetoideae: Lamiaceae). Plant Syst Evol. 2008;275:93–108.Article Google Scholar Li J-C, Zhang J-W, Zhang D-G, Deng T, Volis S, Sun H, et al. Phylogenetic position of the Chinese endemic genus Heterolamium: a close relative of subtribe Nepetinae (Lamiaceae). J Jap Bot.
2017;92:12–9.CAS Google Scholar Takhtajan A. Flowering Plants. Netherlands: Springer; 2009.Book Google Scholar Thorne RF. Classification and geography of the flowering plants. Bot Rev. 1992;58:225–348.Article Google Scholar Cronquist A. The evolution and classification of flowering plants. New York: Columbia University Press; 1988.
Google Scholar Reveal JL. An outline of a classification scheme for extant flowering plants. Phytoneuron. 2012;37:1–211. Google Scholar Barrabé L, Karnadi-Abdelkader G, Ounémoa J, De Kok RPJ, Robert N, Gâteblé G. Recircumscription of Oxera (Lamiaceae: Ajugoideae) to include Faradaya based on molecular and anatomical data. Bot J Linn Soc.
2015;179:693–791.Article Google Scholar Barrabé L, Lavergne S, Karnadi-Abdelkader G, Drew BT, Birnbaum P, Gâteblé G. Changing ecological opportunities facilitated the explosive diversification of New Caledonian Oxera (Lamiaceae). Syst Biol. 2018;68:460–81.Article PubMed Central Google Scholar Thorne RF. The calssificaiton and geography of the flowering plants: Dicotyledons of the class Angiospermae.
Bot Rev. 2000;66:441–647.Article Google Scholar Abu-Asab MS, Cantino PD. Phylogenetic implications of pollen morphology in Tribe Ajugeae (Labiatae). Syst Bot. 1993;18:100–22.Article Google Scholar Dumortier BC. Analyse des Familles des Plantes. Tournay: J. Casterman; 1829. Google Scholar Steane DA, Mabberley DJ.
Rotheca (Lamiaceae) revived. Novon. 1998;8:204–6.Article Google Scholar Yuan Y-W, Mabberley DJ, Steane DA, Olmstead RG. Further disintegration and redefinition of Clerodendrum (Lamiaceae): implications for the understanding of the evolution of an intriguing breeding strategy. Taxon. 2010;59:125–33.Article Google Scholar Charles-Dominique T, Davies TJ, Hempson GP, Bezeng BZ, Daru BH, Kabongo RM, et al. Spiny plants, mammal browsers, and the origin of African savannas. Proc. Natl Acad Sci U S A. 2016;113:E5572–9.Article CAS Google Scholar Huang MJ, Crawford DJ, Freudenstein JV, Cantino PD. Systematics of Trichostema (Lamiaceae): evidence from ITS, ndhF, and morphology.
Syst Bot.
2008;33:437–46.Article Google Scholar Chen S-L, Guo R-L. Caryopteris. In: Chen S-L, Pei J, editors. Flora Reipublicae Popularis Sinicae, vol. 65(1). Beijing: Science Press; 1982. p. 194–208. Google Scholar Rimpler H, Winterhalter C, Falk U. Cladistic analysis of the subfamily Caryopteridoideae Briq. and related taxa of Verbenaceae and Lamiaceae using morphological and chemical characters. In: Harley RM, Reynolds T, editors. Advances in Labiatae Sciences. London: Royal Botanic Gardens, Kew; 1992. p. 39–54. Google Scholar Huang MJ. Systematics of Trichostema L. (Lamiaceae) and phylogenetic relationships with its disjunct taxa in Asia. Ph.D. Dissertation. The Ohio State University; 2002.Shi S-H, Du Y-Q, Boufford DE, Gong X, Huang Y-L, He H-H, et al. Phylogenetic position of Schnabelia, a genus endemic to China: Evidence from sequences of cpDNA matK gene and ITS regions.
Chin Sci Bull. 2003;48:1576–80.Article CAS Google Scholar Cantino PD, Wagstaff ST, Olmstead RG. Caryopteris (Lamiaceae) and the conflict between phylogenetic and pragmatic considerations in botanical nomenclature. Syst Bot. 1998;23:369–86.Article Google Scholar P'ei C, Chen S-L. Verbenaceae. In: Chen SL, Pei J, editors. Flora Reipublicae Popularis Sinicae, Vol. 65(1). Beijing: Science Press; 1982. p. 1–208. Google Scholar Chen S-L, Gilbert MG. Verbenaceae. In: Wu Z-Y, Raven PH, editors. Flora of China, Vol. 17. Beijing: Science Press; St. Louis: Missouri Botanical Garden Press; 1994. pp. 1–49.Paton AJ. A global taxonomic investigation of Scutellaria (Labiatae). Kew Bull. 1990;45:399–450.Article Google Scholar Paton AJ. The phytogeography of Scutellaria L.
Notes Roy Bot Gard Edinburgh. 1990;46:345–9. Google Scholar Li B, Xu W-X, Tu T-Y, Wang Z-S, Olmstead RG, Peng H, et al. Phylogenetic position of Wenchengia (Lamiaceae): a taxonomically enigmatic and critically endangered genus. Taxon. 2012;61:392–401.Article Google Scholar Ryding O. Amount of calyx fibres in Lamiaceae, relation to calyx structure, phylogeny and ecology. Plant Syst Evol. 2007;268:45–58.Article Google Scholar Chiang Y-C, Huang B-H, Liao P-C.
Diversification, biogeographic pattern, and demographic history of Taiwanese Scutellaria species inferred from nuclear and chloroplast DNA.
Plos One. 2012;7:e50844.Article CAS PubMed PubMed Central Google Scholar Zhao F, Liu E-D, Peng H, Xiang C-L. A new species of Scutellaria (Scutellarioideae, Lamiaceae) from Sichuan Province in southwest China. PeerJ. 2017;5:e3624.Article PubMed PubMed Central Google Scholar Safikhani K, Jamzad Z, Saeidi H. Phylogenetic relationships in Iranian Scutellaria (Lamiaceae) based on nuclear ribosomal ITS and chloroplast trnL-F DNA data. Plant Syst Evol. 2018;304:1077–89.Article Google Scholar Scheen A-C, Albert VA. Nomenclatural and taxonomic changes within the Leucas clade (Lamioideae; Lamiaceae). Syst Geogr Pl. 2007;77:229–38. Google Scholar Mathiesen C, Scheen A-C, Lindqvist C. Phylogeny and biogeography of the lamioid genus Phlomis (Lamiaceae). Kew Bull.
2011;66:83–99.Article Google Scholar Bendiksby M, Salmaki Y, Bräuchler C, Ryding O. The generic position of Stachys tibetica Vatke and amalgamation of the genera Eriophyton and Stachyopsis (Lamiaceae subfam. Lamioideae). Plant Syst Evol. 2014;300:961–71.Article Google Scholar Bean AR. A taxonomic revision of Anisomeles R.Br. (Lamiaceae). Austrobaileya. 2015;9:321–81. Google Scholar Wang Q. A monograph of the genus Microtoena (Lamiaceae). Beijing: Science Press; 2018. Google Scholar Bhatti GR, Ingrouilie M. Systematics of Pogostemon (Labiatae). Bull Nat Hist Mus Lond. (Bot.). 1997;27:77–147. Google Scholar Yao G, Deng Y-F, Ge X-J. A taxonomic revision of Pogostemon (Lamiaceae) from China. Phytotaxa. 2015;200:1–67.Article Google Scholar Yao G, Drew BT, Yi T-S, Yan H-F, Yuan Y-M, Ge X-J. Phylogenetic relationships, character evolution and biogeographic diversification of Pogostemon s.l. (Lamiaceae).
Mol Phylogenet Evol. 2016;98:184–200.Article PubMed Google Scholar Kudô Y. Labiatarum sino-japonicarum prodromus.
Mem Fac Taihoku Imp Univ. 1929;2:1–332.
Google Scholar Press JR. Taxonomic studies in the Labiatae tribe Pogostemoneae. Bull Nat Hist Mus Lond (Bot). 1982;10:1–89. Google Scholar Keng H. Labiatae. In: CGGJ VS, editor. Flora Malesiana ser. 1, 8(3). Leyden: Noordhoff International Publishing; 1978. p. 301–94. Google Scholar Hasskarl JC. Plantarum genera et species novae aut reformatae Javenses. Flora 25 Beibl. 1842;2(1–2):1–56.Ryding O. Pericarp structure and phylogeny of Lamiaceae subfamily Pogostemonoideae. Nord J Bot. 1994;14:59–63.Article Google Scholar Ryding O. Pericarp structure and phylogeny of the Lamiaceae-Verbenaceae-complex. Pl Syst Evol. 1995;198:101–41.Article Google Scholar Xiang C-L, Liu E-D, Peng H. A key to the genus Chelonopsis (Lamiaceae) and two new combinations: C. rosea var. siccanea and C. souliei var. cashmerica comb. nov. Nord J Bot. 2008;26:31–4. Google Scholar Xiang C-L, Dong Z-H, Peng H, Liu Z-W. Trichome micromorphology of the East Asiatic genus Chelonopsis (Lamiaceae) and its systematic implications. Flora. 2010;205:434–41.Article Google Scholar Xiang C-L, Funamoto T, Evangelista EV, Zhang Q, Peng H. Pollen morphology of the East Asiatic genus Chelonopsis (Lamioideae: Lamiaceae) and allied genera, with reference to taxonomic implications and potential pollination ecology. Plant Biosyst. 2013;147:620–8.Article Google Scholar Bongcheewin B, Grayer RJ, Porter EA, Kite GC, Simmonds MSJ, Ingrouille MJ, et al. A chemossystematic study of the genus Gomphostemma and related genera (Lamiaceae). Biochem Syst Ecol.
2014;57:305–16.Article CAS Google Scholar Abu-Asab MS, Cantino PD. Systematic implications of pollen morphology in subfamilies Lamioideae and Pogostemonoideae (Labiatae). Ann Mo Bot Gard.
1994;81:653–86.Article Google Scholar Ryding O. Pericarp structure in the subtribe Melittidinae (Lamiaceae-Lamioideae) and its systematic implications. Bot Jahrb Syst. 1994;115:547–55. Google Scholar Ryding O. Pericarp structure in the tribe Prasieae (Lamiaceae-Lamioideae) and its systematic implications. Bot Jahrb Syst. 1994;116:391–9. Google Scholar Hu G-X, Balangcod TD, Xiang C-L. Trichome micromorphology of the Chinese-Himalayan genus Colquhounia (Lamiaceae), with emphasis on taxonomic implications. Biologia.
2012;67:867–74.Article Google Scholar Scheen A-C, Lindqvist C, Fossdal CG, Albert VA. Molecular Phylogenetics of tribe Synandreae, a North American lineage of lamioid mints (Lamiaceae). Cladistics. 2008;32:299–314.Article Google Scholar Roy T, Catlin NS, Garner DMG, Cantino PD, Scheen A-C, Lindqvist C. Evolutionary relationships within the lamioid tribe Synandreae (Lamiaceae) based on multiple low-copy nuclear loci. PeerJ. 2016;4:e2220.Article PubMed PubMed Central Google Scholar WCSP. World Checklist of Selected Plant Families. Facilitated by the Royal Botanic Gardens, Kew. 2020. Accessed 16 Nov 2020.Linnaeus C. Species Plantarum, vol. 2.
Stockholmiae: Salvius; 1753.Dumortier BC. Labiatae. In: Dumortier BC, editor. Florula Belgica. Tournay: J. Casterman; 1827. p. 42–9. Google Scholar Bentham G. Labiatae. In: De Candolle A, editor. Prodromus systematis naturalis regni vegetabilis. Vol.
12. Paris: V. Masson; 1848. p. 27–603. Google Scholar Reichenbach L.
Labiatae. In: Reichenbach L, editor. Flora Germanica Excursoria. Leipzig: Carolum Cnobloch; 1830–1832. pp. 305–335.Cosson E, German E. Tribu V.
Lamioideae. In: Cosson E, German E, editors. Synopsis analytique de la Flore des environs de Paris ou description abregee des familles et des genres. Paris: Fortin, Masson et Cie; 1845. pp. 122–127.Boissier E. Flora Orientalis. Vol. 4(2). Genevae & Basilea: H. Georg; 1879. Google Scholar Bhattacharjee R. Taxonomic studies in Stachys: II. A new infrageneric classification of Stachys L. Notes Roy Bot Gard Edinburgh. 1980;38:65–96. Google Scholar Tomas-Bárberán FA, Gil MI, Ferreres F, Tomas-Lorente F. Flavonoid p-coumaroylglucosides and 8-hydroxyflavone allosylglucosides in some labiatae. Phytochemistry. 1992;31:3097–102.Article Google Scholar Giuliani C, Bini LM. Glandular trichomes as further differential characters between Stachys subgenus Betonica (L.) Bhattacharjee and Stachys subgenus Stachys. Plant Biosyst. 2012;146:1–8.Article Google Scholar Krestovskaya T. Synopsis of the genus Betonica L. (Lamiaceae). Novosti Sist Vyssh Rast. 2014;45:96–109. Google Scholar Lazkov GA, Sennikov AN. Taxonomic corrections and new records in vascular plants of Kyrgyzstan, 4. Mem Soc. Fauna Flora Fenn.
2015;91:67–83. Google Scholar Ball PW. Stachys L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, et al., editors. Flora Europaea. Cambridge: Cambridge University Press; 1972. p. 151–7. Google Scholar Visiani R. Labiatae Juss.
In: Visiani R, editor. Flora Dalmatica, Vol. 2. Lipsae: Apud Pribdbricum Hofmbistbr; 1847. p. 182–225.Townsend CC. Galeopsis L. In: Tutin TG, Heywood VH, Burges NA, Moore DM, Valentine DH, Walters SM, et al., editors. Flora Europaea, Vol. 3: Diapensiaceae to Myoporaceae. Cambridge: Cambridge University Press; 1972. p. 145–7. Google Scholar Meusel H, Jäger EJ, Rauschert SW, Weiner E. Vergleichende Chorologie der zentraleuropäischen Flora. Text u. Karten. Bd. 2. Jena: VEB Fischer; 1978. Google Scholar Goldblatt P, Johnson DE. Index to plant chromosome numbers 2001–2003. Monogr Syst Bot Missouri Bot Gard. 2006;106:1–242. Google Scholar Tomás-Bárberán FA, Gil MI, Ferreres F, Tomás-Lorente F. Correlations between flavonoid composition and infrageneric taxonomy of some European Galeopsis species.
Phytochemistry. 1991;30:3311–4.Article Google Scholar Müntzing A. Outlines to a genetic monograph of the genus Galeopsis with special reference to the nature and inheritance of partial sterility. Hereditas.
1930;13:185–341.Article Google Scholar Bendiksby M, Tribsch A, Borgen L, Trávníček P, Brysting A. Allopolyploid origins of the Galeopsis tetraploids – revisiting Müntzing's classical textbook example using molecular tools. New Phytol. 2011;191:1150–67.Article CAS PubMed Google Scholar Lindqvist C, Motley TJ, Jeffrey JJ, Albert VA.
Cladogenesis and reticulation in the Hawaiian endemic mints (Lamiaceae). Cladistics. 2003;19:480–95.Article PubMed Google Scholar Salmaki Y, Heubl G, Weigend M. Towards a new classification of tribe Stachydeae (Lamiaceae): naming clades using molecular evidence.
Bot J Linn Soc. 2019;190:345–59.Article Google Scholar Cantino PD, De Queiroz K. International code of phylogenetic nomenclature, version 4c.
2010. . Accessed 1 May 2020. Google Scholar Abu-Asab MS, Cantino PD. Pollen morphology in subfamily Lamioideae (Labiatae) and its phylogenetic implications. In: Harley R, Reynolds T, editors. Advances in Labiatae science. London: Royal Botanic Gardens, Kew; 1992. p. 361–79. Google Scholar Salmaki Y, Jamazad Z, Zarre S, Bräuchler C. Pollen morphology of Stachys (Lamiaceae) in Iran and its systematic implication. Flora. 2008;203:627–39.Article Google Scholar Salmaki Y, Zarre S, Jamzad Z.
Nutlet micromorphology and its systematic implication in Stachys L. (Lamiaceae) in Iran. Feddes Repert. 2008;119:607–21.Article Google Scholar Salmaki Y, Zarre S, Jamzad Z, Bräuchler C. Trichome micromorphology of Iranian Stachys (Lamiaceae) with emphasis on its systematic implication. Flora. 2009;204:371–81.Article Google Scholar Salmaki Y, Zarre S, Lindqvist C, Heubl G, Bräuchler C. Comparative leaf anatomy of Stachys (Lamiaceae: Lamioideae) in Iran with a discussion on its subgeneric classification. Plant Syst Evol.
2011;294:109–25.Article Google Scholar Li HW, Hedge IC. Lamiaceae. In: WU Z-Y, Raven PH, editors. Flora of China, Vol. 17. Beijing: Science Press; St. Louis: Missouri Botanical Garden Press; 1994. pp.
50–299.Ryding O. Pericarp structure and phylogeny of the Phlomis group (Lamiaceae subfam. Lamioideae). Bot Jahrb Syst. 2008;127:299–316.Article Google Scholar Seyedi Z, Salmaki Y. Trichome morphology and its significance in the systematics of Phlomoides (Lamiaceae; Lamioideae; Phlomideae). Flora. 2015;213:40–8.Article Google Scholar Patzak VA.
Revision der Gattung Ballota Section Ballota. Ann Naturhist Mus Wien. 1958;62:57–86. Google Scholar Sebald O. Die Gattung Otostegia Bentham (Labiatae) in Afrika und auf der arabischen Halbinsel. Stuttgart Beitr Naturk Ser A. 1973;263:1–84. Google Scholar Seybold VS. Revision der Persischen Marrubium-Arten (Labiatae). Stuttgart Beitr Naturk, Ser A. 1978;310:1–31. Google Scholar Scheen A-C, Albert VA. Molecular Phylogenetics of the Leucas Group (Lamioideae; Lamiaceae). Syst Bot. 2009;34:173–81.Article Google Scholar Kilian N, Hein P, Hubaishan MA. New and noteworthy records for the flora of Yemen, chiefly of Hadhramout and Al-Mahra. Willdenowia. 2002;32:239–69.Article Google Scholar Ryding O. Reconsideration of Wiedemannia and notes on the circumscription of Lamium (Lamiaceae). Bot Jahrb Syst. 2003;124:325–35.Article Google Scholar Bendiksby M, Brysting AK, Thorbek L, Gussarova G, Ryding O. Molecular phylogeny and taxonomy of the genus Lamium L. (Lamiaceae): Disentangling origins of presumed allotetraploids. Taxon. 2011;60:986–1000.Article Google Scholar Krawczyk K, Korniak T, Sawicki J. Taxonomic status of Galeobdolon luteum Huds. (Lamiaceae) from classical taxonomy and phylogenetics perspectives. Acta Biol Craco Ser B. 2013;55:18–28.
Google Scholar Krawczyk K, Szczecińska M, Sawicki J. Evaluation of 11 single-locus and seven multilocus DNA barcodes in Lamium L.
(Lamiaceae). Mol Ecol Resour.
2013;14:272–85.Article PubMed CAS Google Scholar Krestovskaya TV. Menitskia, a new genus of the Labiatae family.
Bot Zhurn. 2006;91:1892–4. Google Scholar Zhao F, Peng H. The complete chloroplast genome of Caryopteris incana (Lamiaceae) and phylogenetic analysis. Mitochondrial DNA B. 2020;5:1399–400.Article Google Scholar Zhao Y, TO T, Xiang C-L. The complete chloroplast genome of Phlomoides betonicoides (Lamiaceae), a traditional Tibetan medicinal herb. Mitochondrial DNA B. 2020;5:75–6.Article Google Scholar Xu W-Q, Losh J, Chen C, Li P, Wang R-H, Zhao Y-P, et al. Comparative genomics of figworts (Scrophularia, Scrophulariaceae), with implications for the evolution of Scrophularia and Lamiales. J Syst Evol. 2019;57:55–65.Article Google Scholar Doyle JJ, Doyle JL. A rapid DNA isolation procedure for small amounts of fresh leaf tissue. Phytochem Bull. 1987;19:11–5. Google Scholar Andrews S. Fast QC: A quality control tool for high throughput sequence data. v. 0.10.0. 2010. . Google Scholar Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: a new genome assembly algorithm and its applications to single cell sequencing. J Comput Biol. 2012;19:455–77.Article CAS PubMed PubMed Central Google Scholar Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, et al. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. Bioinformatics.
2012;28:1647–9.Article PubMed PubMed Central Google Scholar Chen H, Zhang J, Yuan G, Liu C. Complex Interplay among DNA modification, noncoding RNA expression and protein-coding RNA expression in Salvia miltiorrhiza chloroplast genome. Plos One.
2014;9:e99314.Article PubMed PubMed Central CAS Google Scholar Schattner P, Brooks AN, Lowe TM.
The tRNAscan-SE, snoscan and snoGPS web servers for the detection of tRNAs and snoRNAs. Nucleic Acids Res. 2005;33:686–9.Article CAS Google Scholar Lohse M, Drechsel O, Kahlau S, Bock R. Organellar Genome DRAW-a suite of tools for generating physical maps of plastid and mitochondrial genomes and visualizing expression datasets.
Nucleic Acids Res. 2013;41:W575–81.Article PubMed PubMed Central Google Scholar Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods. 2012;9:357–9.Article CAS PubMed PubMed Central Google Scholar Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7.0: improvements in performance and usability. Mol Biol Evol. 2013;30:772–80.Article CAS PubMed PubMed Central Google Scholar Müller K, Müller J, Quandt D. PhyDe: Phylogenetic data editor, version 0.9971. 2010. . Google Scholar Talavera G, Castresana J.
Improvement of phylogenies after removing divergent and ambiguously aligned blocks from protein sequence alignments. Syst Biol.
2007;56:564–77.Article CAS PubMed Google Scholar Darriba D, Taboada GL, Doallo R, Posada D. JModelTest 2: more models, new heuristics and parallel computing. Nat Methods. 2012;9:772.Article CAS PubMed PubMed Central Google Scholar Ronquist F, Teslenko M, Van der Mark P, Ayres DL, Darling A, Höhna S, et al. MrBayes 3.2: Efficient Bayesian phylogenetic inference and model choice across a large model space. Syst Biol. 2012;61:539–42.Article PubMed PubMed Central Google Scholar Miller MA, Pfeiffer W, Schwartz T. Creating the CIPRES Science Gateway for inference of large phylogenetic trees. Gateway Computing Environments Workshop (GCE). 2010. p. 1–8.Rambaut A, Suchard MA, Drummond AJ. Tracer v1.6. 2014. . Google Scholar Rambaut A. FigTree, v.1.4.2. 2014. . Accessed 16 Dec 2019.
Google Scholar Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 2014;30:1312–3.Article CAS PubMed PubMed Central Google Scholar Zhao F, Chen YP, Salmaki Y, Drew BT, Wilson TC, Scheen AC et al. An updated tribal phylogeny of Lamiaceae: evidence from complete plastome sequences. Treebase. S26639?x-access-code=bb02a4c5bc226f4604690ea0f21ccd41&format=html) Accessed 16 July 2020.Thiers B. Index Herbariorum: a global directory of public herbaria and associated staff. New York Botanical Garden's Virtual Herbarium. 2020.Available at . Accessed 16 July 2020. Google Scholar Page 2Phrymaceae (outgroup) Mimulus sp.The United States Botanic Garden (USBG), United States19,584,540478MT473772AjugoideaeAjugeaeCaryopteris forrestii DielsLijiang, Yunnan, China67,295,160485MT473742AjugoideaeTeucrieaeSchnabelia oligophylla Hand.-Mazz.Kunming, Yunnan, China67,359,376726MT473777AjugoideaeClerodendreaeClerodendrum japonicum (Thunb.) SweetKunming, Yunnan, China69,357,954854MT473745AjugoideaeClerodendreaeClerodendrum trichotomum Thunb.Huairou, Beijing, China69,621,568536MT473746AjugoideaeRotheceaeRotheca serrata (L.) Steane & Mabb.Kunming, Yunnan, China69,698,896328MT473776Callicarpioideae–Callicarpa americana L.Gainesville, Florida, United States69,222,992NA--Callicarpioideae–Callicarpa arborea Roxb.Kunming, Yunnan, China70,066,596341MT473738Callicarpioideae–Callicarpa brevipes (Benth.) HanceGuangzhou, Guangdong, China68,119,222383MT473739Callicarpioideae–Callicarpa macrophylla VahlKunming, Yunnan, China69,104,110499MT473740Callicarpioideae–Callicarpa peichieniana Chun & S.L. Chen ex H. Ma & W.B. YuGuangzhou, Guangdong, China68,759,068215MT473741Cymarioideae–Cymaria dichotoma Benth.Changjiang, Hainan, China68,070,4641189MT473753LamioideaeParaphlomideaeParaphlomis javanica (Blume) PrainKunming, Yunnan, China66,797,022239MT473773LamioideaeGomphostemmateaeGomphostemma lucidum Wall. ex Benth.Changjiang, Hainan, China66,781,246274MT473764LamioideaeGomphostemmateaeChelonopsis souliei (Bonati) Merr.Litang, Sichuan, China67,646,436572MT473743LamioideaeColquhouunieaeColquhounia coccinea Wall.Kunming, Yunnan, China66,842,836171MT473749LamioideaeColquhounieaeColquhounia seguinii VaniotKunming, Yunnan, China66,760,344337MT473750LamioideaeColquhounieaeColquhounia vestita Wall.Cuona, Xizang, China67,753,130192MT473751LamioideaeLamieaeLamium amplexicaule L.Zuogong, Xizang, China67,339,8143104MT473770LamioideaeSynandreaeMacbridea alba Chapm.The United States Botanic Garden (USBG), United States20,514,794474MT473771LamioideaeStachydeaeGaleopsis bifida Boenn.Deqin, Yunnan, China67,442,714500MT473759NepetoideaeElsholtzieaeElsholtzia densa Benth.Shangri-La, Yunnan, China18,273,001888MT473757NepetoideaeElsholtzieaeElsholtzia rugulosa Hemsl.Kunming, Yunnan, China67,318,028553MT473758NepetoideaeOcimeaeSiphocranion flavidum Y.P. Chen & C.L. XiangMalipo, Yunnan, China13,829,468436MT473778NepetoideaeOcimeaeSiphocranion macranthum (Hook. f.) C.Y. WuNanchuan, Congqing, China13,860,798241MT473779NepetoideaeOcimeaeHanceola exserta Y.Z. Sun ex C.Y. WuHezhou, Guangxi, China67,557,758203MT473765NepetoideaeOcimeaeIsodon amethystoides (Benth.) H. HaraLin'an, Zhejiang, China25,146,824696MT473767NepetoideaeOcimeaeIsodon lophanthoides (Buch.-Ham. ex D. Don) H. HaraKunming, Yunnan, China40,730,966316MT473768NepetoideaeOcimeaeIsodon ternifolius (D. Don) KudôLongling, Yunnan, China32,984,960542MT473769NepetoideaeOcimeaeColeus xanthanthus C.Y. Wu & Y.C. HuangMengla, Yunnan, China25,669,120821MT473748NepetoideaeMenheaeDracocephalum taliense ForrestHeqing, Yunnan, China68,863,176446MT473756NepetoideaeMenheaeClinopodium abyssinicum (Benth.) KuntzeKabarnet, Baringo, Kenya48,657,815833MT473747Peronematoideae–Garrettia siamensis H.R. FletcherMengla, Yunnan, China69,566,4861905MT473760Peronematoideae–Hymenopyramis cana CraibChangjiang, Hainan, China66,946,216298MT473766Premnoideae–Premna szemaoensis C. P'eiKunming, Yunnan, China69,409,616477MT473775Premnoideae–Premna vietnamensis Bo LiK'Bang, Gia Lai, Vietnam80,675,070460MT473774Premnoideae–Gmelina arborea Roxb. ex Sm.Mengla, Yunnan, China67,974,942493MT473761Premnoideae–Gmelina hainanensis Oliv.Kunming, Yunnan, China67,354,6401527MT473762Premnoideae–Gmelina philippensis Cham.Mengla, Yunnan, China69,953,046479MT473763ProstantheroideaeChloantheaeChloanthes coccinea Bartl.Australian National Botanic Gardens (ANBG), Australia81,265,290598MT473744ProstantheroideaeChloantheaeDasymalla teckiana (F. Muell.) B.J. Conn & HenwoodAustralian National Botanic Gardens (ANBG), Australia41,308,508519MT473754ProstantheroideaeChloantheaeDicrastylis parvifolia F. Muell.Australian National Botanic Gardens (ANBG), Australia81,081,410577MT473755Symphorematoideae–Congea tomentosa Roxb.Mengla, Yunnan, China40,494,132110MT473752Symphorematoideae–Sphenodesme mollis CraibMengla, Yunnan, China81,008,454529MT473780Tectonoideae–Tectona grandis L. f.Mengla, Yunnan, China40,169,710514MT473781Viticoideae–Vitex glabrata R. Br.Mengla, Yunnan, China70,126,282722MT473782Viticoideae–Vitex negundo var. cannabifolia (Siebold & Zucc.) Hand.-Mazz.Kunming, Yunnan, China67,083,4681387MT473783Viticoideae–Vitex quinata (Lour.) F.N. WilliamsMengla, Yunnan, China69,282,366828MT473784Viticoideae–Vitex tripinnata (Lour.) Merr.Guangzhou, Guangdong, China67,065,5141404MT473785Viticoideae–Vitex yunnanensis W.W. Sm.Luquan, Yunnan, China70,217,642395MT473786