

# Checklist of the Orchids of Australia Including its Island Territories



Gary N. Backhouse  
Robert J. Bates  
Andrew P. Brown  
Lachlan M. Copeland



# Contents

Introduction.....	page 1
Australian species.....	page 4
Non-native (exotic or weed) species.....	page 52
Australian hybrids.....	page 53
Australian Capital Territory species.....	page 63
Australian Capital Territory hybrids.....	page 66
New South Wales species.....	page 67
New South Wales hybrids.....	page 79
Northern Territory species.....	page 81
Queensland species.....	page 82
Queensland hybrids.....	page 92
South Australian species.....	page 93
South Australian hybrids.....	page 102
Tasmanian species.....	page 105
Tasmanian hybrids.....	page 109
Victorian species.....	page 110
Victorian hybrids.....	page 123
Western Australian species.....	page 124
Western Australian hybrids.....	page 135
Australia’s Island Territories species.....	page 139

version 1 March 2016

cover photograph: *Sarcochilus fitzgeraldii* Dorrigo, New South Wales

# A Checklist of the Orchids of Australia Including its Island Territories

## Introduction

This Checklist covers the 1,869 species, subspecies and varieties of wild orchids (comprising 1,794 species and 116 subspecies and varieties) in 113 genera found in Australia and its island territories. This total includes 395 species considered distinct but not yet named and described, and these species have been identified by a 'tag name'. An additional six species that are not native to Australia but have become established in the wild there are included at the end of the main Checklist. This checklist also includes 546 naturally occurring hybrids (including 48 named wild hybrids) in 15 genera plus 6 intergeneric crosses, that have been observed in the field. Many of the named hybrids were mistakenly named as species before their hybrid status was recognised, while others were deliberately named as hybrids. The inclusion of the many hundreds of recognised but un-named species and naturally occurring hybrids makes this the most comprehensive checklist of Australian orchids ever compiled. The Checklist has been prepared using information sourced from recent State or regional guides to orchids, herbarium checklists, photographs and the personal observations and experience of the authors. It has been prepared in two versions: an Excel spreadsheet that can be modified by users, and a PDF version that can be printed for a hard copy.

## Design of the Checklist

The Checklist is divided into a number of columns, headed 'Genus', 'Alternate Genus', 'Species', 'Subspecies/Variety', 'Taxonomic Authority', 'Total' (total number of species) and 'Seen'. The Checklist also includes separate sheets for the orchids (both species and hybrids) recorded in each State/Territory plus 'Extralimital' which includes Australia's island territories (e.g. Christmas Island, Cocos-Keeling Islands, Norfolk Island, islands in the Torres Strait etc.). Note that for the purposes of this Checklist Lord Howe Island is included under New South Wales and Macquarie Island is included under Tasmania, which are the relevant administrative authorities for those islands. Jervis Bay Territory, although administered by the Australian Government, is here included under New South Wales. The final column is 'Notes', which for undescribed species (those with a 'tag name') provides some information on which named species the taxon most closely resembles, how it differs from that taxon and distribution/habitat where this might be important information for identification. For a few validly described species some information has been provided where there is doubt as to the veracity of a described species. Genera are arranged alphabetically and then named species within each genus arranged alphabetically, followed by un-named species with tag names arranged alphabetically.

## Scientific Names

The taxonomy used in this checklist generally follows the genera used in the Australian Plant Census and that are broadly recognised by the Council of Heads of Australia Herbaria. Unfortunately there is not complete agreement amongst all the State and Territory herbaria on a standard list of orchid genera, so we have had to make some judgements on which genera to use in this Checklist. In many cases (such as for *Caladenia* and *Pterostylis*) an alternate genus is provided for users of the many segregate genera that have been described in recent years but that are mostly not recognised by State and Territory herbaria. In most cases where a species was described in a genus that is not currently widely recognised in Australia, a new name combination in the appropriate recognised genus has been made. However, there are several instances where species have recently been described in alternate genera but the appropriate new name combinations have not yet been made. Although the species in question have been included in the relevant recognised genera, the relevant cell has been highlighted in yellow. Where a species is recognised as distinct but has not yet been described, a unique 'tag name' is provided, and information on its affinities, a brief description and distribution included under 'Notes'.

## Acknowledgements

Dean Rouse provided details for many of the undescribed *Prasophyllum* species from south-eastern Australia. Colin Bower and David Jones generously provided advice on many of the undescribed species, especially in New South Wales. Jean Egan and Tony Wood provided comments for the Australian Capital Territory list and David James provided comments for the Queensland list. Our thanks to all contributors.

## Abbreviations

ACT=Australian Capital Territory  
AIT= Australia's Island Territories  
AVH=Australia's Virtual Herbarium (avh.ala.org.au)  
NSW=New South Wales  
NT=Northern Territory  
Qld=Queensland  
SA=South Australia  
Tas=Tasmania  
Vic=Victoria  
WA=Western Australia

## References

- Backhouse, G.N. and Jeanes, J.A. 1995. The Orchids of Victoria. The Meigunyah Press, Carlton.
- Backhouse, G., Kosky, B., Rouse, D. and Turner, J. 2016. Bush Gems: A Guide to the Wild Orchids of Victoria. Ebook, Melbourne.
- Bates, R.J. 2011. South Australia's Native Orchids. DVD-ROM. Native Orchid Society of South Australia Inc., Adelaide.
- Bates R.J. 2016. *Diuris calcicola* a new species of mallee donkey orchid from South Australia. *Australian Orchid Review* 80(5): 43-46
- Bishop, T. 2000. Field Guide to the Orchids of New South Wales and Victoria. University of New South Wales Press, Sydney.
- Bower, C. 2007. The Wasp, Ant and Bird Orchids of the *Chiloglottis* alliance - a general overview. *The Orchadian* 15(8): 402-415.
- Brown, A.P. and Brockman, G. 2007. *Caladenia petrensis* and *C. saxicola* (Orchidaceae), two new ironstone endemics from south-west Western Australia. *Nuytsia* 17: 73-79
- Brown, A.P. and Brockman, G. 2015. New taxa of *Caladenia* (Orchidaceae) from south-west Western Australia. *Nuytsia* 25: 45-123
- Brown, A., Dundas, P., Dixon, K. and Hopper, S. 2008. Orchids of Western Australia. University of Western Australia Press, Perth.
- Brown, A., Dixon, K., French, C. and Brockman, G. 2013. Field Guide to the orchids of Western Australia. Simon Nevill Publications, Perth.
- Copeland, L. 2005. Rare or threatened orchids of the Guyra-Ebor Road. *The Orchadian* 14(12): 554-559
- Govaerts, R. 2016. World Checklist of Orchidaceae. Royal Botanic Gardens, Kew
- Hoffman, N. and Brown, A.P. 2011. Orchids of South-Western Australia, 3rd edition. University of Western Australia Press, Perth.
- Jeanes, J. 2004. A revision of the *Thelymitra pauciflora* R.Br. (Orchidaceae) complex in Australia. *Muelleria* 19: 19-79.
- Jeanes, J. and Backhouse, G. 2006. Wild Orchids of Victoria, Australia. Aquatic Photogrtaphics, Seaford.
- Jones, D.L. 2006. A Complete Guide to Native Orchids of Australia. New Holland, Frenchs Forest.
- Jones, D.L. 2008. Field Guide to the Orchids of the Australian Capital Territory. National Parks Association, Australian Capital Territory.
- Jones, D.L. and Clements, M.A. 2006. New taxa of Australian Orchidaceae. *Australian Orchid Research* 5: 1-184

Jones, D., Wapstra, H., Tonelli, P. and Harris, S. 1999. The Orchids of Tasmania. The Meigunyah Press, Carlton.

Kuiter, R.H. 2010. Victoria's Small Caladenias Ebook. Aquatic Photographics, Seaford.

Rouse, D.T. 2002. Report on the taxonomic, ecological and conservation status of taxa in the genus *Prasophyllum* R. Br. in south-eastern Australia. Unpublished report to the Australian Orchid Foundation.

Stephenson, A. 2011. Orchid Species of the Shoalhaven NSW Australia. self-published

### Corrections and Updates

This Checklist of Australian Orchids will be periodically amended and updated, especially as new orchid species are described. Also, there will inevitably be errors or omissions, especially for new State/Territory records or novel hybrids not included in this list. Please send any comments to Gary by email to [gnb1803@gmail.com](mailto:gnb1803@gmail.com)

### Citation

When making reference to this checklist, cite as Backhouse, G.N., Bates, R.J., Brown, A.P. and Copeland, L.M. 2016. A Checklist of the Orchids of Australia Including its Island Territories.

**Table 1:** Orchid numbers nationally and by each State and Territory

State/Territory	Taxa*	Hybrids
National	1,869	546
Australian Capital Territory	132	11
New South Wales	558	60
Northern Territory	45	0
Queensland	494	23
South Australia	378	149
Tasmania	210	8
Victoria	454	132
Western Australia	488	218
Australia's Island Territories	27	0

\* taxa includes species, subspecies and varieties

## Checklist of the Orchids of Australia Including its Island Territories

<b>Acianthella</b> D.L. Jones & M.A.Clem			
<i>Acianthella amplexicaulis</i> (F.M.Bailey) D.L.Jones & M.A.Clem.	<i>Acianthus</i>	NSW Qld	
<i>Acianthella sublesta</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Acianthus</i>	Qld	
<b>Acianthus</b> R.Br.			
<i>Acianthus apprimus</i> D.L.Jones		NSW	
<i>Acianthus borealis</i> D.L.Jones		Qld	
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	SA Vic Tas NSW Qld	
<i>Acianthus caudatus</i> variety <i>pallidus</i> Rupp	<i>Nemacianthus</i>	SA Tas	doubtfully distinct variety described from Tasmania but recognised in South Australia (Bates 2011)
<i>Acianthus collinus</i> D.L.Jones		Vic NSW ACT	
<i>Acianthus exiguus</i> D.L.Jones		NSW	
<i>Acianthus exsertus</i> R.Br.		Vic NSW ACT Qld	
<i>Acianthus fornicatus</i> R.Br.		NSW Qld	
<i>Acianthus ledwardii</i> Rupp		Qld	probably extinct
<i>Acianthus pusillus</i> D.L.Jones		SA Vic Tas NSW Qld	
<i>Acianthus saxatilis</i> D.L.Jones & M.A.Clem.		NSW Qld	
<i>Acianthus</i> species 'Gibraltar Range'		NSW	autumn flowering aff. <i>caudatus</i> with smaller flowers; wet forest in the Gibraltar Range (L. Copeland unpublished)
<i>Acianthus</i> species 'NSW Northern Tablelands'		NSW	autumn flowering aff. <i>fornicatus</i> with short dark sepal clubs and no teeth or lobes on lateral sepals; grassy open forest on the New England Tableland (Bishop 2000); records of <i>A. apprimus</i> from northern NSW are probably this taxon
<i>Acianthus</i> species 'Tathra'		NSW	tiny aff. <i>fornicatus</i> with labellum <4 mm long; coastal scrubs and heathy woodland on South Coast headlands (Bishop 2000)
<b>Acriopsis</b> Reinw. ex Blume			
<i>Acriopsis emarginata</i> D.L.Jones & M.A.Clem.		Qld	
<b>Adenochilus</b> Hook.f.			
<i>Adenochilus nortonii</i> Fitzg.		NSW	
<b>Anoectochilus</b> Blume			
<i>Anoectochilus yatesiae</i> F.M.Bailey		Qld	
<b>Aphyllorchis</b> Blume			
<i>Aphyllorchis anomala</i> Dockrill		Qld	
<i>Aphyllorchis queenslandica</i> Dockrill		Qld	
<b>Apostasia</b> Blume			
<i>Apostasia wallichii</i> R.Br.		Qld	also called <i>A. stylioides</i> (F.Muell.) Rchb.f.
<b>Appendicula</b> Blume			
<i>Appendicula australiensis</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Podochilus</i>	Qld	
<b>Arthrochilus</b> F.Muell.			
<i>Arthrochilus apectus</i> D.L.Jones		Qld	
<i>Arthrochilus aquilus</i> D.L.Jones		Qld	
<i>Arthrochilus byrnesii</i> Blaxell	<i>Phoringopsis</i>	WA NT	
<i>Arthrochilus corinnae</i> D.L.Jones		Qld	
<i>Arthrochilus dockrillii</i> Lavarack	<i>Phoringopsis</i>	Qld	
<i>Arthrochilus irritabilis</i> F.Muell.		Qld	
<i>Arthrochilus latipes</i> D.L.Jones		NT	
<i>Arthrochilus lavarackianus</i> (D.L.Jones) Lavarack	<i>Phoringopsis</i>	Qld	
<i>Arthrochilus oreophilus</i> D.L.Jones		Qld	
<i>Arthrochilus prolixus</i> D.L.Jones		NSW Qld	
<i>Arthrochilus rosulatus</i> D.L.Jones		Qld	
<i>Arthrochilus sabulosus</i> D.L.Jones		Qld	

Australian Species	Alternate Genus	Distribution	Notes
<i>Arthrochilus stenophyllus</i> D.L.Jones		Qld	
<i>Arthrochilus</i> species 'Bradshaw Station'	<i>Phoringopsis</i>	NT	no information available
<i>Arthrochilus</i> species 'spinifex dweller'	<i>Phoringopsis</i>	NT	no information available; NT Herbarium record
<b>Brachypeza</b> Garay			
<i>Brachypeza archytas</i> (Ridl.) Garay		AIT	Christmas Island
<b>Bromheadia</b> Lindl.			
<i>Bromheadia pulchra</i> Schltr.		Qld	
<b>Bryobium</b> Lindl.			
<i>Bryobium dischorense</i> (Schltr.) M.A.Clem. & D.L.Jones	<i>Eria</i>	Qld	may be the same as <i>Bryobium intermedium</i>
<i>Bryobium eriaeoides</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Eria</i>	Qld	
<i>Bryobium intermedium</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Eria</i>	Qld	may be the same as <i>Bryobium dischorense</i>
<i>Bryobium irukandjianum</i> St.Cloud	<i>Eria</i>	Qld	
<i>Bryobium queenslandicum</i> T.E.Hunt	<i>Eria</i>	Qld	
<i>Bryobium retusum</i> (Blume) Y.P.Ng & P.J.Cribb	<i>Eria</i>	AIT	Christmas Island
<b>Bulbophyllum</b> Thouars			
<i>Bulbophyllum argyropus</i> (Endl.) Rchb.f.	<i>Adelopetalum</i>	NSW Qld AIT	also on Lord Howe Island, Norfolk Island
<i>Bulbophyllum baileyi</i> F.Muell.	<i>Carpamorchis</i>	Qld	
<i>Bulbophyllum boonjee</i> B.Gray & D.L.Jones	<i>Adelopetalum</i>	Qld	
<i>Bulbophyllum bowkettiae</i> F.M.Bailey	<i>Serpenticaulis</i>	Qld	
<i>Bulbophyllum bracteatum</i> Fitzg.	<i>Adelopetalum</i>	NSW Qld	
<i>Bulbophyllum elisae</i> F.Muell.	<i>Adelopetalum</i>	NSW Qld	
<i>Bulbophyllum evasum</i> T.E.Hunt & Rupp	<i>Kauorchis</i>	Qld	
<i>Bulbophyllum exiguum</i> F.Muell.	<i>Adelopetalum</i>	NSW Qld	
<i>Bulbophyllum gadgarrense</i> Rupp	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum globuliforme</i> Nicholls	<i>Oncophyllum</i>	NSW Qld	
<i>Bulbophyllum gracillimum</i> Rolfe	<i>Cirrhopetalum</i>	Qld	
<i>Bulbophyllum grandimesense</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum johnsonii</i> T.E.Hunt	<i>Serpenticaulis</i>	Qld	
<i>Bulbophyllum kirkwoodiae</i> T.E.Hunt	<i>Serpenticaulis</i>	Qld	usually included in <i>Bulbophyllum johnsonii</i>
<i>Bulbophyllum lageniforme</i> F.Muell.	<i>Adelopetalum</i>	Qld	
<i>Bulbophyllum lamingtonense</i> D.L.Jones	<i>Oxysepala</i>	NSW Qld	
<i>Bulbophyllum lewisense</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum lilianae</i> Rendle	<i>Adelopetalum</i>	Qld	
<i>Bulbophyllum longiflorum</i> Thouars	<i>Cirrhopetalum</i>	Qld	also called <i>B. clavigerum</i> (Fitzg.) F.M.Bailey
<i>Bulbophyllum macphersonii</i> Rupp	<i>Blepharochilum</i>	Qld	
<i>Bulbophyllum maxillare</i> (Lindl.) Rchb.f.	<i>Ephippium</i>	Qld	also called <i>B. masdevalliaceum</i> Kraenzl.
<i>Bulbophyllum minutissimum</i> F.Muell.	<i>Oncophyllum</i>	NSW Qld	
<i>Bulbophyllum nematopodium</i> F.Muell.	<i>Papulipetalum</i>	Qld	
<i>Bulbophyllum newportii</i> F.M.Bailey	<i>Adelopetalum</i>	Qld	
<i>Bulbophyllum radicans</i> F.M.Bailey	<i>Fruticicola</i>	Qld	
<i>Bulbophyllum schillerianum</i> Rchb.f. subspecies <i>maritimum</i> F.Muell.	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum schillerianum</i> Rchb.f. subspecies <i>schillerianum</i>	<i>Oxysepala</i>	NSW Qld	
<i>Bulbophyllum shepherdii</i> (F.Muell.) Rchb.f.	<i>Oxysepala</i>	NSW Qld	
<i>Bulbophyllum sladeanum</i> A.D.Hawkes	<i>Blepharochilum</i>	Qld	sometimes called <i>Bulbophyllum macphersonii</i> var. <i>spathulatum</i>
<i>Bulbophyllum wadsworthii</i> Dockrill	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum weinthalii</i> R.S.Rogers subspecies <i>striatum</i> D.L.Jones	<i>Spilorchis</i>	Qld	
<i>Bulbophyllum weinthalii</i> R.S.Rogers subspecies <i>weinthalii</i>	<i>Spilorchis</i>	NSW Qld	
<i>Bulbophyllum whitei</i> T.E.Hunt & Rupp	<i>Serpenticaulis</i>	Qld	sometimes included in <i>Bulbophyllum johnsonii</i>
<i>Bulbophyllum wilkianum</i> T.E.Hunt	<i>Adelopetalum</i>	Qld	
<i>Bulbophyllum windsorensense</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	Qld	
<i>Bulbophyllum wolfei</i> B.Gray & D.L.Jones	<i>Serpenticaulis</i>	Qld	

Australian Species	Alternate Genus	Distribution	Notes
<b>Burnettia</b> Lindl.			
<i>Burnettia cuneata</i> Lindl.		Vic Tas NSW	
<b>Caladenia</b> R.Br.			
<i>Caladenia abbreviata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia actensis</i> D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	ACT	
<i>Caladenia aestiva</i> D.L.Jones	<i>Arachnorchis</i>	Vic NSW ACT	
<i>Caladenia alata</i> R.Br.	<i>Petalochilus</i>	Vic Tas NSW Qld	
<i>Caladenia alpina</i> R.S.Rogers	<i>Stegostyla</i>	Vic Tas NSW ACT	
<i>Caladenia ambusta</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia amnicola</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia amoena</i> D.L.Jones	<i>Arachnorchis</i>	Vic	
<i>Caladenia ampla</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	doubtfully distinct from and probably a synonym of <i>Caladenia reticulata</i>
<i>Caladenia ancylosa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia angustata</i> Lindl.	<i>Stegostyla</i>	Tas	
<i>Caladenia anthracina</i> D.L.Jones	<i>Arachnorchis</i>	Tas	
<i>Caladenia applanata</i> Hopper & A.P.Br. subspecies <i>applanata</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia applanata</i> Hopper & A.P.Br. subspecies <i>erubescens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia arenaria</i> Fitzg.	<i>Arachnorchis</i>	NSW	
<i>Caladenia arenicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia argocalla</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia armata</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	ACT	
<i>Caladenia arrecta</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia atrata</i> D.L.Jones	<i>Stegostyla</i>	Tas	
<i>Caladenia atrochila</i> D.L.Jones	<i>Petalochilus</i>	Tas	
<i>Caladenia atroclavia</i> D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	Qld	
<i>Caladenia atrovespa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic NSW ACT Qld	
<i>Caladenia attenuata</i> (W.Brinsley) D.L.Jones	<i>Petalochilus</i>	NSW	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>attingens</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>effusa</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>gracillima</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia audasii</i> R.S.Rogers	<i>Arachnorchis</i>	Vic	
<i>Caladenia aurantiaca</i> (R.S.Rogers) Rupp	<i>Petalochilus</i>	Vic Tas	in Tasmania only on Deal Island Bass Strait
<i>Caladenia aurulenta</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	SA	
<i>Caladenia australis</i> G.W.Carr	<i>Arachnorchis</i>	Vic Tas	in Tasmania only on Flinders Island Bass Strait; no recent records
<i>Caladenia barbarella</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	WA	
<i>Caladenia barbarossa</i> Rchb.f.	<i>Drakonorchis</i>	WA	
<i>Caladenia behrii</i> Schldl.	<i>Arachnorchis</i>	SA	
<i>Caladenia bicalliata</i> R.S.Rogers subspecies <i>bicalliata</i>	<i>Jonesiopsis</i>	WA SA Vic	
<i>Caladenia bicalliata</i> R.S.Rogers subspecies <i>cleistogama</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA SA	
<i>Caladenia bigeminata</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia brachyscapa</i> G.W.Carr	<i>Arachnorchis</i>	Vic Tas	
<i>Caladenia branwhitei</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	NSW	doubtfully distinct from and probably a synonym of <i>Caladenia leptoclavia</i>
<i>Caladenia brevisura</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia brownii</i> Hopper	<i>Arachnorchis</i>	WA	
<i>Caladenia brumalis</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia bryceana</i> R.S.Rogers subspecies <i>bryceana</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia bryceana</i> R.S.Rogers subspecies <i>cracens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia busselliana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	



Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia cadyi</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	NSW	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>caesarea</i>	<i>Jonesiopsis</i>	WA	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>maritima</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>transiens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia cairnsiana</i> F.Muell.	<i>Jonesiopsis</i>	WA	
<i>Caladenia calcicola</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia callitrophila</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia campbellii</i> D.L.Jones	<i>Petalochilus</i>	Tas	
<i>Caladenia capillata</i> D.L.Jones	<i>Jonesiopsis</i>	SA Vic	
<i>Caladenia cardiochila</i> Tate	<i>Arachnorchis</i>	SA Vic Tas	in Tasmania only on Flinders Island Bass Strait; no recent records
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	SA Vic Tas NSW ACT Qld	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia carnea</i> variety <i>subulata</i> Nicholls	<i>Petalochilus</i>	SA Vic	recognised in one recent publication (Kuitert 2010) but probably only an aberrant form of <i>Caladenia carnea</i>
<i>Caladenia catenata</i> (Sm.) Druce	<i>Petalochilus</i>	Vic NSW Qld	
<i>Caladenia catenata</i> (Sm.) Druce variety <i>gigantea</i> (R.S.Rogers) W.M.Curtis	<i>Petalochilus</i>	Qld	taxon of uncertain status
<i>Caladenia caudata</i> Nicholls	<i>Arachnorchis</i>	Tas	
<i>Caladenia chamaephylla</i> D.L.Jones	<i>Petalochilus</i>	Qld	
<i>Caladenia chapmanii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia christineae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia citrina</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia clarkiae</i> D.L.Jones	<i>Stegostyla</i>	Vic NSW	
<i>Caladenia clavescens</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia clavigera</i> A.Cunn. ex Lindl.	<i>Arachnorchis</i>	SA Vic Tas NSW	
<i>Caladenia clavula</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia cleistantha</i> D.L.Jones	<i>Petalochilus</i>	Vic NSW	
<i>Caladenia coactilis</i> D.L.Jones	<i>Petalochilus</i>	SA	
<i>Caladenia colorata</i> D.L.Jones	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia concinna</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	NSW	
<i>Caladenia concolor</i> Fitzg.	<i>Arachnorchis</i>	Vic NSW	
<i>Caladenia conferta</i> D.L.Jones subspecies <i>conferta</i>	<i>Arachnorchis</i>	SA	
( <i>Caladenia</i> ) <i>conferta</i> D.L.Jones subspecies <i>occidentalis</i> R.J.Bates	<i>Arachnorchis</i>	SA	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	SA Vic Tas NSW ACT	
<i>Caladenia corynephora</i> A.S.George	<i>Arachnorchis</i>	WA	
<i>Caladenia cracens</i> D.L.Jones	<i>Stegostyla</i>	Tas	
<i>Caladenia crebra</i> A.S.George	<i>Arachnorchis</i>	WA	
<i>Caladenia cremna</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia cretacea</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia cristata</i> R.S.Rogers	<i>Arachnorchis</i>	WA	
<i>Caladenia cruciformis</i> D.L.Jones	<i>Arachnorchis</i>	Vic	said to have once occurred in South Australia but now extinct there (Bates 2011)
<i>Caladenia cruscula</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia cucullata</i> Fitzg.	<i>Stegostyla</i>	SA Vic NSW ACT	
<i>Caladenia curtisepala</i> D.L.Jones	<i>Petalochilus</i>	NSW	
<i>Caladenia decora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia denticulata</i> Lindl. subspecies <i>albicans</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia denticulata</i> Lindl. subspecies <i>denticulata</i>	<i>Jonesiopsis</i>	WA	
<i>Caladenia denticulata</i> Lindl. subspecies <i>rubella</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia dienema</i> D.L.Jones	<i>Arachnorchis</i>	Tas	
<i>Caladenia dilatata</i> R.Br.	<i>Arachnorchis</i>	SA Vic Tas	
<i>Caladenia dimidia</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia dimorpha</i> Fitzg.	<i>Stegostyla</i>	NSW	
<i>Caladenia discoidea</i> Lindl.	<i>Arachnorchis</i>	WA	
( <i>Caladenia</i> ) <i>diversiflora</i> R.J.Bates	<i>Arachnorchis</i>	SA	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia dorrienii</i> Domin	<i>Jonesiopsis</i>	WA	
<i>Caladenia douglasiorum</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia doutchiaie</i> O.Sarg.	<i>Jonesiopsis</i>	WA	
<i>Caladenia drakeoides</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	WA	
<i>Caladenia drummondii</i> Benth.	<i>Arachnorchis</i>	WA	
<i>Caladenia dundasiae</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia echidnachila</i> Nicholls	<i>Arachnorchis</i>	Tas	
<i>Caladenia elegans</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia ensata</i> Nicholls	<i>Arachnorchis</i>	WA	
<i>Caladenia erythrochila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia erythronema</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia evanescens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia excelsa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia exilis</i> Hopper & A.P.Br. subspecies <i>exilis</i>	<i>Jonesiopsis</i>	WA	
<i>Caladenia exilis</i> Hopper & A.P.Br. subspecies <i>vanleeuwenii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia exstans</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia falcata</i> (Nicholls) M.A.Clem. & Hopper	<i>Arachnorchis</i>	WA	
<i>Caladenia ferruginea</i> Nicholls	<i>Arachnorchis</i>	WA	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	SA Vic Tas NSW ACT Qld	
<i>Caladenia filifera</i> Lindl.	<i>Jonesiopsis</i>	WA	
<i>Caladenia fitzgeraldii</i> Rupp	<i>Arachnorchis</i>	NSW ACT	
<i>Caladenia flaccida</i> D.L.Jones	<i>Jonesiopsis</i>	SA NSW Qld	South Australian records unlikely to be true <i>Caladenia flaccida</i> ; here listed as an undescribed taxon <i>Caladenia</i> species 'Flinders Ranges red'
<i>Caladenia flava</i> R.Br. subspecies <i>flava</i>		WA	
<i>Caladenia flava</i> R.Br. subspecies <i>maculata</i> Hopper & A.P.Br.		WA	
<i>Caladenia flava</i> R.Br. subspecies <i>sylvestris</i> Hopper & A.P.Br.		WA	
<i>Caladenia flava</i> R.Br. subspecies 'late red'		WA	short, late-flowering <i>C. flava</i> with prominent red markings; wandoo woodland on granite outcrops between Beverley & Williams (Brown <i>et al.</i> 2013); probably only a colourful form of <i>C. flava</i> subsp. <i>flava</i>
<i>Caladenia flavovirens</i> G.W.Carr	<i>Arachnorchis</i>	Vic	
<i>Caladenia flindersica</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	SA	
<i>Caladenia fluvialis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia footeana</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia formosa</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia fragrantissima</i> D.L.Jones & G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia fuliginosa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	SA	
<i>Caladenia fulva</i> G.W.Carr	<i>Arachnorchis</i>	Vic	said to occur in South Australia (Bates 2011), but plants there probably fall within <i>Caladenia colorata</i>
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	SA Vic Tas NSW ACT Qld	
<i>Caladenia fuscolutescens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia gardneri</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia georgei</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia gladiolata</i> R.S.Rogers	<i>Arachnorchis</i>	SA	
<i>Caladenia gracilis</i> R.Br.	<i>Stegostyla</i>	Tas	
<i>Caladenia gracillima</i> (Rupp) D.L.Jones	<i>Petalochilus</i>	Qld	
<i>Caladenia graminifolia</i> A.S.George	<i>Arachnorchis</i>	WA	
<i>Caladenia grampiana</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic	
<i>Caladenia graniticola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia granitora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia harringtoniae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia hastata</i> (Nicholls) Rupp	<i>Arachnorchis</i>	Vic	said to occur in South Australia (Bates 2011), but records there probably based on <i>Caladenia richardsiorum</i> with clubbed petals
<i>Caladenia heberleana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia helvina</i> D.L.Jones	<i>Arachnorchis</i>	Tas	
<i>Caladenia hiemalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia hildae</i> E.E.Pescott & Nicholls	<i>Stegostyla</i>	Vic NSW ACT	
<i>Caladenia hillmanii</i> D.L.Jones	<i>Petalochilus</i>	NSW	
<i>Caladenia hirta</i> Lindl. subspecies <i>hirta</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia hirta</i> Lindl. subspecies <i>rosea</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia hoffmanii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia hopperiana</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia horistes</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia huegelii</i> Rchb.f.	<i>Arachnorchis</i>	WA	
<i>Caladenia incensa</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia incrassata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia infundibularis</i> A.S.George	<i>Arachnorchis</i>	WA	
<i>Caladenia insularis</i> G.W.Carr	<i>Arachnorchis</i>	Vic	
<i>Caladenia integra</i> E.Coleman	<i>Arachnorchis</i>	WA	
<i>Caladenia interanea</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	SA	
<i>Caladenia interjacens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia intuta</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia iridescens</i> R.S.Rogers	<i>Stegostyla</i>	SA Vic	
<i>Caladenia latifolia</i> R.Br.		WA SA Vic Tas	
<i>Caladenia leptochila</i> Fitzg. subspecies <i>dentata</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	SA	
<i>Caladenia leptochila</i> Fitzg. subspecies <i>leptochila</i>	<i>Arachnorchis</i>	SA	records from Victoria and New South Wales are erroneous, probably based on collapsing flowers of another clubbed spider-orchid species
<i>Caladenia leptoclavia</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia leucochila</i> A.P.Br., R.D.Phillips & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia lindleyana</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Arachnorchis</i>	Tas	
<i>Caladenia littoricola</i> R.J.Bates	<i>Petalochilus</i>	SA	
<i>Caladenia lobata</i> Fitzg.	<i>Arachnorchis</i>	WA	
<i>Caladenia lodgeana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>albella</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>australora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>borealis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>calcigena</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>clivicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>crassa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>eminens</i> (Domin) Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>extrema</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>insularis</i> Hopper & A.P.Br. ex A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>longicauda</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>merrittii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>minima</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>redacta</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies <i>rigidula</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia longicauda</i> Lindl. subspecies 'Bulbarnet'	<i>Arachnorchis</i>	WA	<i>longicauda</i> subspecies with robust flowers with relatively short stiff petals and sepals with broad bases, sepals often clubbed, labellum lamina calli in numerous rows; woodland near Moora



Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia longiclavata</i> E.Coleman	<i>Arachnorchis</i>	WA	
<i>Caladenia longifimbriata</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia lorea</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia lowanensis</i> G.W.Carr	<i>Arachnorchis</i>	Vic	
<i>Caladenia luteola</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia macroclavia</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia macrostylis</i> Fitzg.	<i>Arachnorchis</i>	WA	
<i>Caladenia magniclavata</i> Nicholls	<i>Arachnorchis</i>	WA	
<i>Caladenia magnifica</i> (Nicholls) D.L.Jones & G.W.Carr	<i>Arachnorchis</i>	Vic	probably extinct
<i>Caladenia marginata</i> Lindl.		WA	
<i>Caladenia maritima</i> D.L.Jones	<i>Petalochilus</i>	Vic	
<i>Caladenia melanema</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia mentiens</i> D.L.Jones	<i>Petalochilus</i>	SA Vic Tas NSW ACT	
<i>Caladenia meridionalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia mesocera</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	WA	
<i>Caladenia microchila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA SA	
<i>Caladenia minor</i> Hook.f.	<i>Petalochilus</i>	SA NSW	a taxon of highly uncertain identity; Australian records from SA and NSW but generally considered to be endemic to New Zealand
<i>Caladenia montana</i> G.W.Carr	<i>Arachnorchis</i>	Vic NSW ACT	
<i>Caladenia moschata</i> D.L.Jones	<i>Stegostyla</i>	SA Vic NSW ACT	
<i>Caladenia multiclavia</i> Rchb.f.	<i>Jonesiopsis</i>	WA	
<i>Caladenia nana</i> Endl. subspecies <i>nana</i>		WA	
<i>Caladenia nana</i> Endl. subspecies <i>unita</i> (Fitzg.) Hopper & A.P.Br.		WA	sometimes considered a full species <i>Caladenia unita</i>
<i>Caladenia necrophylla</i> D.L.Jones	<i>Arachnorchis</i>	SA	doubtfully distinct from and probably a synonym of <i>Caladenia dilatata</i>
<i>Caladenia nivalis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia nobilis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia occidentalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia oenochila</i> G.W.Carr	<i>Arachnorchis</i>	Vic	
<i>Caladenia oreophila</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic NSW	
<i>Caladenia orestes</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	NSW	
<i>Caladenia orientalis</i> (G.W.Carr) Hopper & A.P.Br.	<i>Arachnorchis</i>	Vic	
<i>Caladenia ornata</i> (Nicholls) D.L.Jones	<i>Petalochilus</i>	SA Vic	
<i>Caladenia osmera</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic NSW	distinguishable from <i>Caladenia montana</i> only by altitude
<i>Caladenia ovata</i> R.S.Rogers	<i>Arachnorchis</i>	SA	
<i>Caladenia pachychila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia pallida</i> Lindl.	<i>Arachnorchis</i>	Tas	possibly extinct
<i>Caladenia paludosa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia paradoxa</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia parva</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia patersonii</i> R.Br.	<i>Arachnorchis</i>	Tas	
<i>Caladenia pectinata</i> R.S.Rogers	<i>Arachnorchis</i>	WA	
<i>Caladenia peisleyi</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	Vic NSW	
<i>Caladenia pendens</i> Hopper & A.P.Br. subspecies <i>pendens</i>	<i>Jonesiopsis</i>	WA	
<i>Caladenia pendens</i> Hopper & A.P.Br. subspecies <i>talbotii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia perangusta</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia petrensis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia phaeoclavia</i> D.L.Jones	<i>Arachnorchis</i>	SA Vic NSW ACT	doubtfully distinct from and probably a synonym of <i>Caladenia parva</i>
<i>Caladenia pholcoidea</i> Hopper & A.P.Br. subspecies <i>augustensis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia pholcoidea</i> Hopper & A.P.Br. subspecies <i>pholcoidea</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia picta</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	NSW Qld	
<i>Caladenia pilotensis</i> D.L.Jones	<i>Arachnorchis</i>	Vic	

Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia plicata</i> Fitzg.	<i>Arachnorchis</i>	WA	
<i>Caladenia pluvialis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia polychroma</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia porphyrea</i> D.L.Jones	<i>Petalochilus</i>	NSW	doubtfully distinguishable from <i>Caladenia quadrifaria</i>
<i>Caladenia postea</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia praecox</i> Nicholls	<i>Stegostyla</i>	Vic NSW ACT	
<i>Caladenia procera</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia prolata</i> D.L.Jones	<i>Petalochilus</i>	SA Vic Tas	
<i>Caladenia pulchra</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia pumila</i> R.S.Rogers	<i>Arachnorchis</i>	Vic	
<i>Caladenia pusilla</i> W.M.Curtis	<i>Petalochilus</i>	SA Vic Tas NSW	
<i>Caladenia pygmaea</i> (R.S.Rogers) R.J.Bates	<i>Petalochilus</i>	SA	
<i>Caladenia quadrifaria</i> D.L.Jones	<i>Petalochilus</i>	NSW Qld	
<i>Caladenia radialis</i> R.S.Rogers	<i>Jonesiopsis</i>	WA	
<i>Caladenia radiata</i> Nicholls	<i>Arachnorchis</i>	WA	
<i>Caladenia remota</i> Hopper & A.P.Br. subspecies <i>parva</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia remota</i> Hopper & A.P.Br. subspecies <i>remota</i>	<i>Jonesiopsis</i>	WA	
<i>Caladenia reptans</i> Lindl. subspecies <i>impensa</i> Hopper & A.P.Br.		WA	
<i>Caladenia reptans</i> Lindl. subspecies <i>reptans</i>		WA	
<i>Caladenia reticulata</i> Fitzg.	<i>Arachnorchis</i>	SA	almost certainly occurs elsewhere in south-eastern Australia and more widely distributed than <i>reticulata</i> as narrowly defined by Jones (2006)
<i>Caladenia rhomboidiformis</i> (E.Coleman) M.A.Clem. & Hopper	<i>Arachnorchis</i>	WA	
<i>Caladenia richardsiorum</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia rigens</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia rigida</i> R.S.Rogers	<i>Arachnorchis</i>	SA	
<i>Caladenia rileyi</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia robinsonii</i> G.W.Carr	<i>Arachnorchis</i>	Vic	
<i>Caladenia roei</i> Benth.	<i>Jonesiopsis</i>	WA	
<i>Caladenia rosella</i> G.W.Carr	<i>Arachnorchis</i>	Vic NSW	almost certainly extinct in NSW, if it ever occurred there
<i>Caladenia saggicola</i> D.L.Jones	<i>Arachnorchis</i>	Tas	
<i>Caladenia sanguinea</i> D.L.Jones	<i>Jonesiopsis</i>	SA	
<i>Caladenia saxatilis</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	SA	
<i>Caladenia saxicola</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia septuosa</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia serotina</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia sigmoidea</i> R.S.Rogers	<i>Arachnorchis</i>	WA	
<i>Caladenia speciosa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia splendens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia starteorum</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia stellata</i> D.L.Jones	<i>Arachnorchis</i>	NSW	erroneously recorded for South Australia, the records there referable to the undescribed <i>Caladenia</i> species 'Murray mallee'
<i>Caladenia straminichila</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia stricta</i> (R.Bates) R.Bates	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia strigosa</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	SA	
<i>Caladenia subglabriphylla</i> (R.J.Bates) M.A.Clem.	<i>Arachnorchis</i>	SA	
<i>Caladenia subtilis</i> D.L.Jones	<i>Arachnorchis</i>	NSW	
<i>Caladenia swartsiorum</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	WA	
<i>Caladenia sylvicola</i> D.L.Jones	<i>Petalochilus</i>	Tas	
<i>Caladenia tensa</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia tentaculata</i> Schldl.	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia tessellata</i> Fitzg.	<i>Arachnorchis</i>	Vic NSW	
<i>Caladenia testacea</i> R.Br.	<i>Stegostyla</i>	NSW	
<i>Caladenia thinicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia thysanochila</i> G.W.Carr	<i>Arachnorchis</i>	Vic	probably extinct
<i>Caladenia tonellii</i> D.L.Jones	<i>Petalochilus</i>	Tas	
<i>Caladenia toxochila</i> Tate	<i>Arachnorchis</i>	SA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia transitoria</i> D.L.Jones	<i>Stegostyla</i>	Vic Tas NSW	
<i>Caladenia transitoria</i> D.L.Jones subspecies 'Kangaroo Island'	<i>Stegostyla</i>	SA	small aff. <i>transitoria</i> ; heathy woodland on limestone on Kangaroo Island; (R. Bates unpublished) subsp. <i>isolata</i> ms
<i>Caladenia uliginosa</i> A.S.George subspecies <i>candicans</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia uliginosa</i> A.S.George subspecies <i>patulens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia uliginosa</i> A.S.George subspecies <i>uliginosa</i>	<i>Arachnorchis</i>	WA	
<i>Caladenia ultima</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia ustulata</i> D.L.Jones	<i>Stegostyla</i>	NSW ACT	
<i>Caladenia valida</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia validinerva</i> Hopper & A.P.Br. ex A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	WA	
<i>Caladenia venusta</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia verrucosa</i> G.W.Carr	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia versicolor</i> G.W.Carr	<i>Arachnorchis</i>	Vic	unconfirmed records from South Australia; if it ever occurred there, it is now extinct in that State
<i>Caladenia villosissima</i> (G.W.Carr) Hopper & A.P.Br.	<i>Arachnorchis</i>	SA Vic	
<i>Caladenia viridescens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
( <i>Caladenia</i> ) <i>viriosa</i> R.J.Bates	<i>Arachnorchis</i>	SA	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia voigtii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia vulgaris</i> D.L.Jones	<i>Petalochilus</i>	SA Vic Tas	
<i>Caladenia vulgata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia wanosa</i> A.S.George	<i>Jonesiopsis</i>	WA	
<i>Caladenia whiteheadii</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	NSW	
<i>Caladenia williamsiae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia winfieldii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	WA	
<i>Caladenia woolcockiorum</i> D.L.Jones	<i>Arachnorchis</i>	SA	
<i>Caladenia xantha</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	WA	
<i>Caladenia xanthochila</i> D.Beardsell & C.Beardsell	<i>Arachnorchis</i>	Vic	unconfirmed records from South Australia; if it ever occurred there, it is now extinct in that State
<i>Caladenia xantholeuca</i> D.L.Jones	<i>Petalochilus</i>	SA	
<i>Caladenia zephyra</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	SA	
<i>Caladenia</i> species 'Ararat hills'	<i>Arachnorchis</i>	Vic	late winter flowering aff. <i>venusta</i> with smaller flowers, short marginal teeth and sparser calli; Ararat Hills (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'Bindoon'	<i>Jonesiopsis</i>	WA	aff. <i>hiemalis</i> with larger often yellowish-white flowers; granite hills between Bindoon and Gillingarra
<i>Caladenia</i> species 'Buchan'	<i>Arachnorchis</i>	Vic	large aff. <i>montana</i> with longer, more slender sepals and petals with long indistinct clubs and more numerous marginal teeth; open forest near Buchan (Turner <i>et al.</i> 2014)
<i>Caladenia</i> species 'Carboor'	<i>Arachnorchis</i>	Vic	aff. <i>concolor</i> with dull red flowers; grassy open forest in nth-eastern Victoria (Backhouse 2011); possibly the same as <i>Caladenia clavescens</i>
<i>Caladenia</i> species 'central NSW'	<i>Petalochilus</i>	NSW	small, self-pollinating aff. <i>vulgaris</i> with colourful flowers with narrow petals and sepals with rounded tips; woodland from Cowra to Armidale (Bishop 2000)
<i>Caladenia</i> species 'central Victoria'	<i>Arachnorchis</i>	Vic	aff. <i>formosa</i> with smaller, dull maroon flowers with long filamentous tails; dry woodland in central Vic (Jeanes & Backhouse 2006; Backhouse 2011)
<i>Caladenia</i> species 'Chapple Vale'	<i>Stegostyla</i>	Vic	summer-flowering aff. <i>iridescens</i> with golden yellow flowers; Otway Ranges (Jeanes & Backhouse 2006); probably only a colour form of <i>Caladenia iridescens</i>
<i>Caladenia</i> species 'coastal dunes'	<i>Arachnorchis</i>	SA Vic	aff. <i>orientalis</i> with large pale yellow flowers; coastal heath and heathy woodland, often on coastal dunes; sporadic between Portland (Vic) and Millicent (SA) (Backhouse 2011; Bates 2011)
<i>Caladenia</i> species 'coastal Eyre Peninsula'	<i>Arachnorchis</i>	SA	aff. <i>macroclavia</i> with smaller flowers with more distinctly clubbed petals; heath and heathy woodland on limestone, coastal and hinterland Eyre Peninsula (Bates 2011)
<i>Caladenia</i> species 'Coonawarra'	<i>Arachnorchis</i>	SA Vic	aff. <i>colorata</i> with larger, variably coloured flowers with paler brown tails; heathy woodland between Tallageira (Vic), Lucindale and Penola (SA) (Jeanes & Backhouse 2006; Backhouse 2011); usually included in <i>Caladenia colorata</i>
<i>Caladenia</i> species 'Coorong'	<i>Petalochilus</i>	SA	aff. <i>carnea</i> with brighter pink flowers with broader sepals and petals; coastal scrubs on alkaline sands in the Coorong region (Bates 2011)



Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia</i> species 'crowded calli'	<i>Arachnorchis</i>	NSW	aff. <i>concinna</i> with larger flowers with larger, more numerous crowded labellum calli extending almost to the apex, and short sepal clubs; shrubby woodland between Temora and Rankin Springs (Bower 2008; Backhouse 2011)
<i>Caladenia</i> species 'Dorrigo'	<i>Stegostyla</i>	NSW	aff. <i>transitoria</i> with internally white sepals and petals; montane scrub on rainforest margin near Dorrigo (Copeland unpubl.; Backhouse 2011) <i>dorrigoensis</i> ms
<i>Caladenia</i> species 'Earlston'	<i>Arachnorchis</i>	Vic	aff. <i>concolor</i> with glossy red flowers with dark red labellum & thickened tepal tails; dry woodland on low hills at Earlston (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'east of Esperance'	<i>Arachnorchis</i>	WA	late flowering aff. <i>hiemalis</i> with larger flowers from east of Esperance; brief mention in Brown <i>et al.</i> 2013
<i>Caladenia</i> species 'Finnis'	<i>Arachnorchis</i>	SA	extinct: the photo of this taxon in Bates (2011) suggests it was probably a hybrid, perhaps involving <i>Caladenia cardiochila</i> and <i>Caladenia parva</i>
<i>Caladenia</i> species 'Fleurieu Peninsula'	<i>Arachnorchis</i>	SA	aff. <i>patersonii</i> with large white to pink to red flowers; heathy woodland on Fleurieu Peninsula (Bates 2011; Backhouse 2011)
<i>Caladenia</i> species 'Flinders Ranges red'	<i>Jonesiopsis</i>	SA	aff. <i>filamentosa</i> with larger flowers with white-margined lateral sepals; dry woodland on rocky hills in the central Flinders Ranges (Bates 2011; Backhouse 2011); usually included in <i>Caladenia flaccida</i>
<i>Caladenia</i> species 'glowing green'	<i>Petalochilus</i>	SA Vic	aff. <i>prolata</i> with larger, more translucent flowers; seasonally damp River Red Gum flats in lower south-east SA & western Vic (Bates 2011)
<i>Caladenia</i> species 'inland NSW'	<i>Arachnorchis</i>	NSW	aff. <i>atrovespa</i> with smaller, stiffer flowers; has a different wasp pollinator to <i>atrovespa</i> ; dry woodland on inland plains between Rankin Springs, West Wyalong and Lockhart (Colin Bower personal communication; Backhouse 2011)
<i>Caladenia</i> species 'Inverleigh'	<i>Arachnorchis</i>	Vic	aff. <i>patersonii</i> with large cream to pinkish flowers; open heathy woodland on sandy loam at Inverleigh (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'Kangaroo Island'	<i>Stegostyla</i>	SA	aff. <i>transitoria</i> with self-pollinating flowers that do not open and have very short barely distinguishable teeth on labellum mid-lobe; heathy woodland on Kangaroo Island (Bates 2011)
<i>Caladenia</i> species 'Kaniva'	<i>Petalochilus</i>	Vic	aff. <i>pusilla</i> with 1-5 pink flowers with cupped tepals; dry heathy woodland near Kaniva (Backhouse 2011)
<i>Caladenia</i> species 'Kaniva Giant'	<i>Jonesiopsis</i>	Vic	aff. <i>filamentosa</i> with large dark red flowers with long drooping segments; mallee heath from Kiata to Kaniva (Backhouse 2011)
<i>Caladenia</i> species 'Kilkoy Creek'	<i>Petalochilus</i>	Qld	aff. <i>carnea</i> of uncertain identity; Kilkoy Creek (Qld Herbarium record)
<i>Caladenia</i> species 'Kilsyth South'	<i>Arachnorchis</i>	Vic	aff. <i>venusta</i> with smaller dull white flowers with narrower labellum; heathy open forest at Kilsyth South (Backhouse & Jeanes 1995) <i>consternans</i> ms
<i>Caladenia</i> species 'Kooyera'	<i>Arachnorchis</i>	Vic	aff. <i>venusta</i> with smaller flower with dark reddish black glandular tails; grassy woodland on sandy loam near Rheola (Backhouse 2011)
<i>Caladenia</i> species 'large scented'	<i>Petalochilus</i>	Vic	aff. <i>carnea</i> with large, pale, strongly scented flowers; hilly dry woodland between Ararat and Stuart Mill (Backhouse 2011)
<i>Caladenia</i> species 'limestone mallee'	<i>Jonesiopsis</i>	SA	aff. <i>capillata</i> with smaller flowers with very small labellum and relatively stiff petals and sepals; limestone mallee woodland and broombush on Eyre Peninsula; mentioned in NOSSA Feb 2015 (Bates unpublished)
<i>Caladenia</i> species 'little clowns'	<i>Arachnorchis</i>	SA Vic	aff. <i>lowanensis</i> with slightly smaller flowers; heathy woodland near Bordertown SA & western Vic (Bates 2011); probably falls within variation of <i>Caladenia lowanensis</i> (Backhouse 2011)
<i>Caladenia</i> species 'lowan mallee'	<i>Arachnorchis</i>	SA Vic	aff. <i>toxochila</i> with smaller flowers with shorter, thinner, more spaced calli and a more sharply curved column; mallee heath in Little Desert, also dry woodland near Rostron (Backhouse 2011); usually included in <i>Caladenia toxochila</i>
<i>Caladenia</i> species 'lower Glenelg River'	<i>Arachnorchis</i>	Vic	aff. <i>fragrantissima</i> with variable cream to red flowers; heathy woodland on red clays over limestone on Glenelg River near Nelson (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'mallee'	<i>Arachnorchis</i>	SA Vic	small aff. <i>tensa</i> with more sparsely hairy stem and leaf and colourful flowers with almost no white on the labellum; widespread in mallee on sandy soils in SA, extending to west Vic (Bates 2011)
<i>Caladenia</i> species 'mallee heath'	<i>Arachnorchis</i>	SA Vic	aff. <i>arenaria</i> with lge cream flower with long dark red tails; mallee heath & woodland, Little Desert & Miram Piram (Backhouse 2011)
<i>Caladenia</i> species 'Mandurang'	<i>Arachnorchis</i>	Vic	aff. <i>fragrantissima</i> with large stiff white flowers with reddish calli and marginal teeth; dry sparsely shrubby woodland near Mandurang (Backhouse & Jeanes 1995)
<i>Caladenia</i> species 'Monarto South'	<i>Jonesiopsis</i>	SA	aff. <i>capillata</i> with shorter pale yellow sepals and petals; open mallee woodland on sand and clay loams over limestone, Murray Bridge to Middleton (Bates 2011)

Australian Species	Alternate Genus	Distribution	Notes
<i>Caladenia</i> species 'Murray mallee'	<i>Arachnorchis</i>	SA	aff. <i>stellata</i> with larger flowers with the labellum apex usually rolled under; mallee and dry woodland between Mt Pleasant and Langhorne Creek (Backhouse 2011; Bates 2011); usually included in <i>C. stellata</i>
<i>Caladenia</i> species 'Orange'	<i>Arachnorchis</i>	NSW	aff. <i>patersonii</i> with large pinkish mauve flowers with darker calli and teeth and brownish purple tails; hilly open forest near Orange and the Blue Mountains; last seen in the late 1980s and probably extinct (Bishop 2000)
<i>Caladenia</i> species 'pointed flowers'	<i>Petalochilus</i>	NSW ACT	aff. <i>mentiens</i> with sharply pointed sepals and petals and incurved dorsal sepal; hilly dry woodland near Gungahlin (J. Egan & T. Wood personal communication)
<i>Caladenia</i> species 'Raymond Island'	<i>Arachnorchis</i>	Vic	aff. <i>ancylosa</i> with relatively large clubbed greenish flower with long marginal teeth; heathy woodland on Raymond Island (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'Richmond River'	<i>Arachnorchis</i>	NSW	aff. <i>patersonii</i> with creamy yellow flowers with long dark glandular tails; heathy open forest near the Richmond River; collected in the late 1800s and not seen since, probably extinct (Bishop 2000)
<i>Caladenia</i> species 'sand dunes'	<i>Petalochilus</i>	SA	tall aff. <i>carnea</i> with multiple self-pollinating flowers, coastal scrubs on calcareous sand dunes (Bates 2011); <i>brevicaulis</i> ms
<i>Caladenia</i> species 'short yellow clubs'	<i>Arachnorchis</i>	SA	aff. <i>aurulenta</i> with shorter, thicker sepaline clubs; mallee and dry woodland on the Eyre Peninsula (Bates 2011)
<i>Caladenia</i> species 'small tensa'	<i>Arachnorchis</i>	SA	small aff. <i>tensa</i> ; no further information (R. Bates unpublished)
<i>Caladenia</i> species 'south east'	<i>Arachnorchis</i>	SA	large aff. <i>reticulata</i> with yellow flowers; sandplains, flowering after fire (R. Bates unpublished)
<i>Caladenia</i> species 'south Gippsland'	<i>Arachnorchis</i>	Vic	early flowering aff. <i>australis</i> with large flowers with long indistinct clubs; moist forest in south Gippsland (Backhouse 2011); usually included in <i>Caladenia australis</i>
<i>Caladenia</i> species 'southern slopes'	<i>Arachnorchis</i>	Vic	aff. <i>australis</i> with large flowers with long indistinct sepal clubs and shorter petal clubs and has a different wasp pollinator; moist foothill forest in Gippsland from Tonimbuk to Erica (Backhouse 2011)
<i>Caladenia</i> species 'St Andrews'	<i>Arachnorchis</i>	Vic	aff. <i>oenochila</i> with large pale flowers; shrubby woodland on gravelly clay soil at St Andrews Vic (Jeanes & Backhouse 2006); probably only a form of <i>Caladenia oenochila</i>
<i>Caladenia</i> species 'swamp heath'	<i>Arachnorchis</i>	Vic	aff. <i>venusta</i> with small creamy white flowers with dark tails; swampy heath west of Casterton, flowering mostly after fires (Backhouse 2011)
<i>Caladenia</i> species 'tiny white'	<i>Petalochilus</i>	SA	aff. <i>pusilla</i> with tiny (segments 5 mm long), short-lived (1 day), self-pollinating white flowers; damp shaded sites in woodland in the southern Mt Lofty Ranges and lower south-east (Bates 2011)
<i>Caladenia</i> species 'Wandoo'	<i>Jonesiopsis</i>	WA	aff. <i>fluvialis</i> with slightly smaller flowers and upland habitat; woodland west of York and Brookton (A. Brown unpublished)
<i>Caladenia</i> species 'Watheroo'	<i>Jonesiopsis</i>	WA	aff. <i>lorea</i> with creamy white flowers with red apex to labellum; heathy woodland near Watheroo (Backhouse 2011)
<i>Caladenia</i> species 'western Murray Flats'	<i>Arachnorchis</i>	SA	unknown affinity; mentioned in NOSSA Feb 2015; Hartley to Sedan (R. Bates unpublished)
<i>Caladenia</i> species 'western wheatbelt'	<i>Jonesiopsis</i>	WA	aff. <i>dimidia</i> with larger creamy white to yellow flowers with longer drooping segments with brighter red tails; dry woodland from York to Quairadding (A. Brown unpublished)
<i>Caladenia</i> species 'Woronora Plateau'	<i>Petalochilus</i>	NSW	aff. <i>alata</i> with calli coalescing into wrinkled ridges and backward pointing large teeth at mid-lob base; heathy woodland on Woronora Plateau (Bishop 2000)
<i>Caladenia</i> species 'Yarram'	<i>Petalochilus</i>	Vic	aff. <i>catenata</i> with variable white to crimson flowers; heathy open forest on sandy loam near Yarram (Jeanes & Backhouse 2006); most likely a hybrid swarm involving <i>Caladenia carnea</i> and <i>Caladenia catenata</i>
<b><i>Calanthe</i> R.Br.</b>			
<i>Calanthe australasica</i> D.L.Jones & M.A.Clem.		NSW Qld	
<b><i>Caleana</i> R.Br.</b>			
<i>Caleana major</i> R.Br.		SA Vic Tas NSW Qld	
<b><i>Calochilus</i> R.Br.</b>			
<i>Calochilus ammobius</i> D.L.Jones & B.Gray		Qld	
<i>Calochilus caeruleus</i> L.O. Williams		Qld NT	
<i>Calochilus caesius</i> D.L.Jones		WA NT	
<i>Calochilus campestris</i> R.Br.		Vic Tas NSW Qld	
<i>Calochilus cleistanthus</i> D.L.Jones		Qld	
<i>Calochilus cupreus</i> R.S.Rogers		SA	
<i>Calochilus gracillimus</i> Rupp		NSW	

Australian Species	Alternate Genus	Distribution	Notes
<i>Calochilus grandiflorus</i> (Benth.) Domin		NSW Qld	
<i>Calochilus herbaceus</i> Lindl.		SA Vic Tas	
<i>Calochilus holtzei</i> F.Muell.		WA NT	also recorded for Qld but these records are in doubt, probably based on misidentifications of <i>Calochilus imperiosus</i>
<i>Calochilus imberbis</i> R.S.Rogers		Vic Tas	
<i>Calochilus imperiosus</i> D.L.Jones		Qld NT	
<i>Calochilus kalaru</i> D.L.Jones		NSW	
<i>Calochilus metallicus</i> D.L.Jones		Qld	
<i>Calochilus montanus</i> D.L.Jones		NSW ACT	
<i>Calochilus paludosus</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Calochilus platyichilus</i> D.L.Jones		SA Vic Tas NSW ACT Qld	virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus praeltus</i> D.L.Jones		NSW	
<i>Calochilus pruinus</i> D.L.Jones		WA SA Vic	
<i>Calochilus psednus</i> D.L.Jones & Lavarack		Qld	
<i>Calochilus pulchellus</i> D.L.Jones		NSW	
<i>Calochilus richiae</i> Nicholls		Vic	
<i>Calochilus robertsonii</i> Benth.		SA Vic NSW	
<i>Calochilus russeus</i> D.L.Jones		NSW	very similar to, and possibly just a montane form of, <i>Calochilus therophilus</i>
<i>Calochilus sandrae</i> D.L.Jones		NSW ACT	
<i>Calochilus saprophyticus</i> R.S.Rogers		Vic NSW ACT	
<i>Calochilus stramenicola</i> D.L.Jones		WA	
<i>Calochilus therophilus</i> D.L.Jones		Vic NSW ACT Qld	in Qld recorded as <i>Calochilus gracillimus</i>
<i>Calochilus uliginosus</i> D.L.Jones		WA	
<i>Calochilus</i> species 'Booyup Brook'		WA	aff. <i>stramenicola</i> with different habitat and well-spaced flowers; heathy woodland near Booyup Brook (Brown <i>et al.</i> 2013)
<i>Calochilus</i> species 'Coaldale'		NSW	aff. <i>campestris</i> with short flowers with shorter basal plates and broader column wings; heathy woodland on rocky hills near Coaldale (L. Copeland unpublished)
<i>Calochilus</i> species 'Gibraltar Range'		NSW	aff. <i>campestris</i> with slim reddish stem with several longish slender bracts held well away from stem and small, narrow flowers with long narrow rostrum; heathy woodland in Gibraltar Range
<i>Calochilus</i> species 'Kangaroo Island'		SA	small. aff. <i>campestris</i> ; heathy woodland on limestone on Kangaroo Island (R. Bates unpublished)
<i>Calochilus</i> species 'Moondarra'		Vic	small, colourful aff. <i>campestris</i> with short irregular basal plates and short very sparse bristles on labellum; wet heathy woodland near Erica (Backhouse <i>et al.</i> 2016)
<i>Calochilus</i> species 'New England'		NSW	aff. <i>robertsonii</i> with small flowers with reddish eye-like column glands and a sparsely bristled labellum with short apical bristles and a visible naked tail; basalt grasslands and grassy woodlands between Ebor and Ben Lomond
<i>Calochilus</i> species 'Theda Station'		WA	aff. <i>caesius</i> with larger flowers with longer basal hairs and a naked, cupped labellum apex; on sandy soils amongst sedges on seasonally wet sites in the central Kimberley region (Brown <i>et al.</i> 2013); <i>barbarossa</i> ms
<b><i>Cestichis</i> Thouars</b>			
<i>Cestichis angustilabris</i> (F.Muell.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Cestichis bracteata</i> (T.E.Hunt) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Cestichis coelogynoides</i> (F.Muell.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	NSW Qld	
<i>Cestichis condylobulbon</i> (Rchb.f.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Cestichis fleckeri</i> (Nicholls) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Cestichis nugentiae</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Cestichis reflexa</i> (R.Br.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	NSW	
<i>Cestichis swenssonii</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	NSW Qld	



Australian Species	Alternate Genus	Distribution	Notes
<b>Cheirostylis</b> Blume			
<i>Cheirostylis notialis</i> D.L.Jones		NSW Qld	
<i>Cheirostylis ovata</i> (F.M.Bailey) Schltr.		Qld	
<b>Chiloglottis</b> R.Br.			
<i>Chiloglottis anatriceps</i> D.L.Jones		NSW	
<i>Chiloglottis chlorantha</i> D.L.Jones	<i>Simpliglottis</i>	NSW	
<i>Chiloglottis cornuta</i> Hook.f.	<i>Simpliglottis</i>	SA Vic Tas NSW	
<i>Chiloglottis diphylla</i> R.Br.		NSW Qld	
<i>Chiloglottis formicifera</i> Fitzg.	<i>Myrmechila</i>	NSW Qld	
<i>Chiloglottis grammata</i> G.W.Carr	<i>Simpliglottis</i>	Tas	
<i>Chiloglottis gunnii</i> Lindl.	<i>Simpliglottis</i>	Tas	
<i>Chiloglottis jeanesii</i> D.L.Jones	<i>Simpliglottis</i>	Vic	
<i>Chiloglottis longiclavata</i> D.L.Jones		Qld	
<i>Chiloglottis palachila</i> D.L.Jones		NSW	
<i>Chiloglottis platyptera</i> D.L.Jones	<i>Myrmechila</i>	NSW	
<i>Chiloglottis pluricallata</i> D.L.Jones	<i>Simpliglottis</i>	NSW	
<i>Chiloglottis reflexa</i> (Labill.) Druce		Vic Tas NSW	
<i>Chiloglottis seminuda</i> D.L.Jones		Vic NSW ACT	
<i>Chiloglottis sphaerula</i> D.L.Jones		NSW	
<i>Chiloglottis sphyrnoides</i> D.L.Jones		NSW Qld	
<i>Chiloglottis sylvestris</i> D.L.Jones & M.A.Clem.		NSW Qld	
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmechila</i>	SA Vic Tas NSW ACT	
<i>Chiloglottis triceratops</i> D.L.Jones	<i>Simpliglottis</i>	Tas	
<i>Chiloglottis trilabra</i> Fitzg.		Vic NSW ACT	
<i>Chiloglottis trullata</i> D.L.Jones	<i>Myrmechila</i>	Qld	
<i>Chiloglottis truncata</i> D.L.Jones & M.A.Clem.	<i>Myrmechila</i>	Qld	
<i>Chiloglottis turfosa</i> D.L.Jones	<i>Simpliglottis</i>	NSW ACT	
<i>Chiloglottis valida</i> D.L.Jones	<i>Simpliglottis</i>	Vic NSW ACT	'true' <i>valida</i> occurs in the mountains above 500 m but is morphologically indistinguishable from two related cryptic species (Bower 2007; see sp. 'Falls Creek' and sp. 'lowlands'), distribution overlaps with sp. 'Falls Creek'
<i>Chiloglottis</i> species 'Bald Rock'	<i>Myrmechila</i>	NSW Qld	small, spring flowering aff. <i>formicifera</i> with recurved labellum apex, calli extending almost to apex and narrower column wings; grassy open forest on rich loams; border region in Girraween and Bald Rock National Parks (Bishop 2000)
<i>Chiloglottis</i> species 'Barrington Tops'	<i>Simpliglottis</i>	NSW	aff. <i>pluricallata</i> with more numerous calli and different wasp pollinator; heathy swamps on Barrington Tops (Bishop 2000)
<i>Chiloglottis</i> species 'Bathurst'		NSW	small aff. <i>palachila</i> with yellow clubs on lateral sepals and much smaller calli mass; also has a different wasp pollinator; shrubby open forest and woodland near Bathurst (Bishop 2000)
<i>Chiloglottis</i> species 'Ebor'	<i>Simpliglottis</i>	NSW	aff. <i>pluricallata</i> with more numerous calli and different wasp pollinator; wet forest and sphagnum swamps in New England National Park around Point Lookout (Bishop 2000)
<i>Chiloglottis</i> species 'highlands'	<i>Simpliglottis</i>	Vic NSW ACT	aff. <i>valida</i> with slightly smaller labellum with fewer calli and different wasp pollinator; montane damp forest and woodland above 800 m (Bower 2007; Jones <i>et al.</i> 2008); indistinguishable from <i>C. valida</i>
<i>Chiloglottis</i> species 'lowlands'	<i>Simpliglottis</i>	SA Vic Tas NSW	aff. <i>valida</i> indistinguishable from <i>valida</i> but with a different wasp pollinator (Bower 2007); widespread in damp shady sites in south-eastern Aust. below 1,000 m, mostly at low altitudes (Jeanes & Backhouse 2006)
<i>Chiloglottis</i> species 'Maryborough'		Qld	no information available (Qld. Herbarium record)
<i>Chiloglottis</i> species 'Weilangta'	<i>Simpliglottis</i>	Tas	no information available (Tas. Herbarium record)
<b>Chiloschista</b> Lindl.			
<i>Chiloschista phyllorhiza</i> (F.Muell.) Schltr.		Qld NT	
<b>Cooktownia</b> D.L.Jones			
<i>Cooktownia robertsii</i> D.L.Jones		Qld	
<b>Corunastylis</b> Fitzg.			
<i>Corunastylis acuminata</i> (R.S.Rogers) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW Qld	
<i>Corunastylis alticola</i> (D.L.Jones & B. Gray) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis anomala</i> D.L.Jones	<i>Genoplesium</i>	NSW	

Australian Species	Alternate Genus	Distribution	Notes
<i>Corunastylis anthracina</i> D.L.Jones	<i>Genoplesium</i>	NSW	
<i>Corunastylis apostasioides</i> Fitzg.	<i>Genoplesium</i>	NSW	
<i>Corunastylis archeri</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic Tas NSW Qld	'true' <i>archeri</i> almost certainly does not occur in NSW & Qld, with these plants probably comprising a complex of several different, unnamed species
<i>Corunastylis arrecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic NSW ACT	
<i>Corunastylis bishopii</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis brachystachya</i> (Lindl.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Tas	
<i>Corunastylis carecta</i> D.L.Jones & L.M.Copel.	<i>Genoplesium</i>	NSW	
<i>Corunastylis ciliata</i> (Ewart & B.Rees) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	SA Vic	
<i>Corunastylis citriodora</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis clivicola</i> D.L.Jones	<i>Genoplesium</i>	NSW ACT	
<i>Corunastylis conferta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis cornuta</i> D.L.Jones	<i>Genoplesium</i>	NSW ACT	
<i>Corunastylis cranei</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis densa</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic NSW	erroneously recorded for Qld
<i>Corunastylis despectans</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	SA Vic Tas NSW	
<i>Corunastylis ectopa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	ACT	
<i>Corunastylis eriochila</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis filiformis</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW Qld	
<i>Corunastylis fimbriata</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW Qld	
<i>Corunastylis firthii</i> (L.Cady) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Tas	
<i>Corunastylis formosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis insignis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis laminata</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis littoralis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis morina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic NSW	
<i>Corunastylis morrisii</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	SA Vic Tas NSW	
<i>Corunastylis nigricans</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	SA	
<i>Corunastylis nuda</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic Tas NSW ACT	
<i>Corunastylis nudiscapa</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic Tas	
<i>Corunastylis oligantha</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis ostrina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic NSW	
<i>Corunastylis parvicalla</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis pedersonii</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis plumosa</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis psammophila</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis pumila</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic Tas NSW Qld	

Australian Species	Alternate Genus	Distribution	Notes
<i>Corunastylis rhyolitica</i> (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis rufa</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW Qld	
<i>Corunastylis ruppii</i> (R.S.Rogers) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis sagittifera</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW Qld	
<i>Corunastylis sigmoidea</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis simulans</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis stephensonii</i> D.L.Jones	<i>Genoplesium</i>	NSW	
<i>Corunastylis superba</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis systema</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis tasmanica</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Tas	
<i>Corunastylis tecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis tenella</i> D.L.Jones & L.M.Copel.	<i>Genoplesium</i>	NSW	
<i>Corunastylis tepperi</i> (F.Muell. ex. Tepper) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	WA SA Vic	
<i>Corunastylis trifida</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis turfosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Vic NSW ACT	
<i>Corunastylis valida</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	Qld	
<i>Corunastylis vernalis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis woollsii</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	NSW	
<i>Corunastylis</i> species 'Adelaide Hills'	<i>Genoplesium</i>	SA	aff. <i>tasmanica</i> with dark purplish brown and green flowers with a broad labellum with a shallowly toothed, upturned margin; widespread in woodland & heath, often on stony sites (Bates 2011)
<i>Corunastylis</i> species 'Beechworth'	<i>Genoplesium</i>	Vic	aff. <i>archeri</i> with narrower, bluntly pointed labellum with curved but not upturned apex; sparsely shrubby woodland on granite hills at Beechworth (Backhouse <i>et al.</i> 2016)
<i>Corunastylis</i> species 'Charmhaven' <i>Genoplesium</i>		NSW	aff. <i>archeri</i> with fewer larger flowers in a shorter inflorescence; heathy open forest around Charmhaven
<i>Corunastylis</i> species 'dark flowers'	<i>Genoplesium</i>	SA Vic NSW	aff. <i>rufa</i> with pointed dark reddish black labellum; widespread in dry woodland across inland south-eastern Australia (=C. sp. 'inland' in Backhouse <i>et al.</i> 2016; C. sp. 'dark midge' in Bates 2011)
<i>Corunastylis</i> species 'Fraser Island'	<i>Genoplesium</i>	Qld	no further information (Qld Herbarium record)
<i>Corunastylis</i> species 'Gippsland Plains'	<i>Genoplesium</i>	Vic	aff. <i>rufa</i> with fewer, slightly smaller paler flowers with a broader labellum and more broadly triangular petals; heathy open forest Bairnsdale to Stratford (Jeanes & Backhouse 2006)
<i>Corunastylis</i> species 'intermediate'	<i>Genoplesium</i>	SA	aff. <i>rufa</i> with numerous crowded dark purplish brown flowers with contrasting pale (sometimes white-edged) petals; woodland and heath on calcareous soils (Bates 2011)
<i>Corunastylis</i> species 'large bracts'	<i>Genoplesium</i>	SA	robust aff. <i>rufa</i> with numerous crowded maroon and green flowers with long sepals and long pointed floral bracts (most obvious in bud); <i>Triodia</i> hummocks on rocky ridges in dry woodland (Bates 2011)
<i>Corunastylis</i> species 'Mount Billy'	<i>Genoplesium</i>	SA	aff. <i>rufa</i> with uniformly maroon flowers with a longish pointed labellum lacking any raised plates; seasonally wet heaths on clay soils on the Fleurieu Peninsula (Bates 2011)
<i>Corunastylis</i> species 'Mount Kaputar'	<i>Genoplesium</i>	NSW	summer-flowering aff. <i>arrecta</i> with numerous crowded flowers with upswept lateral sepals and hairless petals; heathy woodland and rocky outcrops in the Nandewar Range (Bishop 2000)
<i>Corunastylis</i> species 'Mount Mee'	<i>Genoplesium</i>	Qld	late summer-flowering aff. <i>densa</i> with numerous somewhat spaced flowers with broad lateral sepals and a relatively short leaf ending below the lowest flower; 'clay pan' heath on Mt Mee (David James personal communication)
<i>Corunastylis</i> species 'granite ranges'	<i>Genoplesium</i>	NSW Qld	spring-summer flowering, robust aff. <i>archeri</i> with more crowded flowers with strongly upcurved lateral sepals; open forest on granite ranges of north-eastern NSW (L. Copeland unpublished) and south-eastern Qld



Australian Species	Alternate Genus	Distribution	Notes
<i>Corunastylis</i> species 'Pyrete Range'	<i>Genoplesium</i>	Vic	tall, robust aff. <i>morrisii</i> with more flowers in a denser spike, a broader labellum and lateral sepals pouched at the base; heathy woodland in Otway & Pyrete Ranges (Jeanes & Backhouse 2006); probably a form of <i>C. morrisii</i>
<i>Corunastylis</i> species 'Raby Bay'	<i>Genoplesium</i>	Qld	no further information (Qld Herbarium record)
<i>Corunastylis</i> species 'Wyberba'	<i>Genoplesium</i>	Qld	no further information (Qld Herbarium record)
<b><i>Corybas</i> Salisb.</b>			
<i>Corybas abditus</i> D.L.Jones	<i>Anzybas</i>	WA	
<i>Corybas abellianus</i> Dockrill		Qld	
<i>Corybas aconitiflorus</i> Salisb.		Vic Tas NSW Qld	
<i>Corybas acuminatus</i> M.A.Clem. & Hatch		AIT	Norfolk Island
<i>Corybas barbarae</i> D.L.Jones		NSW Qld	also on Lord Howe Island
<i>Corybas cerasinus</i> D.L.Jones & B.Gray		Qld	
<i>Corybas dentatus</i> D.L.Jones	<i>Corysanthes</i>	SA	possibly of hybrid origin: <i>C. despectans</i> X <i>C. diemenicus</i>
<i>Corybas despectans</i> D.L.Jones & R.C.Nash	<i>Corysanthes</i>	WA SA Vic	
<i>Corybas diemenicus</i> (Lindl.) Rchb.f.	<i>Corysanthes</i>	SA Vic Tas NSW	
<i>Corybas dienemus</i> D.L.Jones	<i>Nematoceras</i>	Tas	Macquarie Island only
<i>Corybas dilatatus</i> (Rupp & Nicholls) Rupp	<i>Corysanthes</i>	SA	
<i>Corybas downlingii</i> D.L.Jones		NSW	doubtfully distinct from, and probably just a form of, <i>Corybas aconitiflorus</i> (Mark Clements CSIRO personal communication)
<i>Corybas expansus</i> D.L.Jones	<i>Corysanthes</i>	SA	
<i>Corybas fimbriatus</i> (R.Br.) Rchb.f.	<i>Corysanthes</i>	Vic Tas NSW Qld	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	SA Vic Tas NSW Qld	
<i>Corybas grumulus</i> D.L.Jones	<i>Corysanthes</i>	Vic NSW ACT	
<i>Corybas hispidus</i> D.L.Jones	<i>Corysanthes</i>	Vic NSW ACT Qld	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	SA Vic Tas NSW ACT	
<i>Corybas limpidus</i> D.L.Jones	<i>Corysanthes</i>	WA	
( <i>Corybas</i> ) <i>minimus</i> (R.J.Bates)	<i>Corysanthes</i>	SA	described in <i>Corysanthes</i> , new combination in <i>Corybas</i> not yet made
<i>Corybas montanus</i> D.L.Jones	<i>Anzybas</i>	Qld	
<i>Corybas pruinosus</i> (A.Cunn.) Rchb.f.	<i>Corysanthes</i>	NSW	
<i>Corybas recurvus</i> D.L.Jones	<i>Corysanthes</i>	WA	
<i>Corybas sulcatus</i> (M.A.Clem. & D.L.Jones) G.N.Backh.	<i>Nematoceras</i>	Tas	Macquarie Island only
<i>Corybas undulatus</i> (A.Cunn.) Rupp	<i>Corysanthes</i>	NSW Qld	
<i>Corybas unguiculatus</i> (R.Br.) Rchb.f.	<i>Anzybas</i>	SA Vic Tas NSW	
<i>Corybas</i> species 'Busselton'	<i>Corysanthes</i>	WA	aff. <i>despectans</i> with tiny dorsal sepal; swampy habitats near Busselton (Brown <i>et al.</i> 2013)
<i>Corybas</i> species 'dark labellum'	<i>Corysanthes</i>	Vic	small aff. <i>diemenicus</i> with dark labellum; heathy woodland on Wilsons Promontory (Jeanes & Backhouse 2006); possibly <i>C. grumulus</i> but not seen for over a decade now
<i>Corybas</i> species 'Gibraltar Range'	<i>Anzybas</i>	NSW	aff. <i>unguiculatus</i> with shorter beak on labellum orifice and with calli on the main and lateral veins inside the orifice (D.L. Jones personal communication comm.); damp montane forest in the Gibraltar Range
<i>Corybas</i> species 'Hammersley Inlet'	<i>Corysanthes</i>	WA	aff. <i>limpidus</i> with smaller flowers and a proportionately smaller dorsal sepal; under tea-tree on sandy soils around Hammersley Inlet (Fitzgerald River National Park) (Brown <i>et al.</i> 2013)
<i>Corybas</i> species 'Naturaliste'	<i>Corysanthes</i>	WA	aff. <i>despectans</i> with very small self-pollinating flowers with incurved labellum margins and peaked dorsal sepal; low coastal heath south between Busselton and Cowaramup (Brown <i>et al.</i> 2013)
<i>Corybas</i> species 'Portland'	<i>Corysanthes</i>	Vic	aff. <i>diemenicus</i> with smaller, pale flower with fewer marginal teeth; coastal heath near Portland (= sp. aff. <i>diemenicus</i> 3 Jeanes & Backhouse 2006); possibly just a mutant form of <i>C. diemenicus</i>
<i>Corybas</i> species 'red-eyed ghost'	<i>Corysanthes</i>	SA	aff. <i>despectans</i> with greenish white, more upright flowers with a narrower dorsal sepal and a red column; calcareous sands under mallee and cypress-pines <i>Callitris</i> spp., Marion Bay, Fleurieu Peninsula and Kangaroo Island (Bates 2011)
<i>Corybas</i> species 'Wilson's Promontory'	<i>Corysanthes</i>	Vic	aff. <i>diemenicus</i> with smaller, darker flower with fewer marginal teeth; damp heathy woodland at Wilsons Promontory (= sp. aff. <i>diemenicus</i> 2 Jeanes & Backhouse 2006); possibly a misidentification of the named hybrid <i>C. xmiscellus</i>

Australian Species	Alternate Genus	Distribution	Notes
<i>Corybas</i> species 'Woolly Tea-tree'	<i>Corysanthes</i>	SA Vic	late spring flowering aff. <i>diemenicus</i> with dark labellum; Woolly Tea-Tree swamps (Jeanes & Backhouse 2006 as sp. aff. <i>diemenicus</i> 1; Bates 2011 as sp. Silky Tea-tree swamps); <i>arcanus</i> ms
<b><i>Corymborkis</i> Thouars</b>			
<i>Corymborkis veratrifolia</i> (Reinw.) Blume		Qld AIT	also on Christmas Island
<b><i>Crepidium</i> Blume</b>			
<i>Crepidium acuminatum</i> (D.Don) Szlach.	<i>Malaxis</i>	NT	
<i>Crepidium fimbriatum</i> (Lavarack) Szlach.	<i>Malaxis</i>	Qld	
<i>Crepidium flavovirens</i> D.L.Jones & M.A.Clem.	<i>Malaxis</i>	Qld	
<i>Crepidium fontinale</i> D.L.Jones & M.A.Clem.	<i>Malaxis</i>	NT	
<i>Crepidium lawleri</i> (Lavarack & B.Gray) Szlach.	<i>Malaxis</i>	Qld	
<i>Crepidium marsupichilum</i> (Upton) Szlach.	<i>Malaxis</i>	Qld NT	
<b><i>Cryptostylis</i> R.Br.</b>			
<i>Cryptostylis erecta</i> R.Br.		Vic NSW Qld	
<i>Cryptostylis hunteriana</i> Nicholls		Vic NSW Qld	
<i>Cryptostylis leptochila</i> Benth.		Vic Tas NSW Qld	
<i>Cryptostylis ovata</i> R.Br.		WA	
<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		SA Vic Tas NSW Qld	
<b><i>Cyanicula</i> Hopper &amp; A.P.Br.</b>			
<i>Cyanicula amplexans</i> (A.S.George) Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula aperta</i> Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula ashbyae</i> Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula caerulea</i> (R.Br.) Hopper & A.P.Br.	<i>Caladenia</i>	Vic NSW ACT Qld	
<i>Cyanicula fragrans</i> Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula gemmata</i> (Lindl.) Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula gertrudeae</i> (Ostenf.) Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula ixioides</i> (Lindl.) Hopper & A.P.Br. subspecies <i>candida</i> Lindl. Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula ixioides</i> (Lindl.) Hopper & A.P.Br. subspecies <i>ixioides</i>	<i>Caladenia</i>	WA	
<i>Cyanicula nikulinskyae</i> Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula sericea</i> (Lindl.) Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<i>Cyanicula</i> species 'Dale'	<i>Caladenia</i>	WA	aff. <i>gemma</i> with an all-green leaf and pale blue flowers with a prominently striped labellum; shrubby woodland adjacent to seasonally wet drainage lines, near Dale, flowering mostly after fire (Brown <i>et al.</i> 2013)
<i>Cyanicula</i> species 'Esperance'	<i>Caladenia</i>	WA	aff. <i>gemma</i> with purplish underside to leaf and purple and white labellum with tall calli, scrubs and woodland, also on granite outcrops, east from Esperance (Brown <i>et al.</i> 2013)
<b><i>Cymbidium</i> Sw.</b>			
<i>Cymbidium canaliculatum</i> R.Br.		WA NSW Qld NT	many forms and varieties have been named (eg. var. <i>sparksii</i> ) although none have any formal taxonomic recognition
<i>Cymbidium madidum</i> Lindl.		NSW Qld	
<i>Cymbidium suave</i> R.Br.		NSW Qld	
<b><i>Cyrtostylis</i> R.Br.</b>			
<i>Cyrtostylis huegelii</i> Endl.		WA	
<i>Cyrtostylis reniformis</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Cyrtostylis robusta</i> D.L.Jones & M.A.Clem.		WA SA Vic Tas	
<i>Cyrtostylis tenuissima</i> (Nicholls & Goadby) D.L.Jones & M.A.Clem.		WA	
<i>Cyrtostylis</i> species 'Cape Naturaliste'		WA	short aff. <i>robusta</i> with smaller flowers and narrower labellum; coastal scrubs and heathy woodland on the Leeuwin-Naturaliste Ridge near Dunsborough (Brown <i>et al.</i> 2013)
<b><i>Danhatchia</i> Garay &amp; Christenson</b>			
<i>Danhatchia australis</i> (Hatch) Garay & Christenson	<i>Yoania</i>	NSW	
<b><i>Demorchis</i> D.L.Jones &amp; M.A.Clem.</b>			
<i>Demorchis queenslandica</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Gastrodia</i>	Qld	

Australian Species	Alternate Genus	Distribution	Notes
<b>Dendrobium Sw.</b>			
<i>Dendrobium adae</i> F.M.Bailey	<i>Thelychiton</i>	Qld	
<i>Dendrobium aemulum</i> R.Br.	<i>Tropilis</i>	NSW Qld	
<i>Dendrobium agrostophyllum</i> F.Muell.	<i>Trachyrhizum</i>	Qld	
( <i>Dendrobium</i> ) <i>angustum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	NSW Qld	described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium antennatum</i> Lindl.	<i>Ceratobium</i>	Qld	also called <i>Dendrobium dalbertisii</i> Rchb.f.
<i>Dendrobium aphyllum</i> (Roxb.) C.E.C.Fisch.		Qld	occurrence in Australia doubtful
<i>Dendrobium aurantiacopurpureum</i> Nicholls	<i>Davejonesia</i>	Qld	doubtfully distinct from <i>Dendrobium lichenastrum</i>
<i>Dendrobium baileyi</i> F.Muell.	<i>Grastidium</i>	Qld	
( <i>Dendrobium</i> ) <i>banksii</i> (D.L.Jones & M.A.Clem.)	<i>Dockrillia</i>	NSW	described in <i>Dockrillia</i> , new combination in <i>Dendrobium</i> not yet made; sometimes included in <i>Dendrobium striolatum</i>
<i>Dendrobium bifalce</i> Lindl.	<i>Leioanthum</i>	Qld	
<i>Dendrobium bigibbum</i> Lindl.	<i>Vappodes</i>	Qld	
<i>Dendrobium bowmanii</i> Benth.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium brachypus</i> (Endl.) Rchb.f.	<i>Thelychiton</i>	AIT	Norfolk Island
<i>Dendrobium brevicaudum</i> D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	Qld	
<i>Dendrobium calamiformis</i> Lodd. ex. Lindl.	<i>Dockrillia</i>	Qld	
<i>Dendrobium callitrophilum</i> B.Gray & D.L.Jones	<i>Tropilis</i>	Qld	usually treated as a variety of <i>Dendrobium aemulum</i>
<i>Dendrobium canaliculatum</i> R.Br.	<i>Cepobaculum</i>	Qld	
<i>Dendrobium cancroides</i> T.E.Hunt	<i>Grastidium</i>	Qld	
<i>Dendrobium carrii</i> Rupp & C.T.White	<i>Australorchis</i>	Qld	
<i>Dendrobium carronii</i> Lavarack & P.J.Cribb	<i>Cepobaculum</i>	Qld	
<i>Dendrobium collinsii</i> (Lavarack) Schuit. & P.B.Adams	<i>Cadetia</i>	Qld	
<i>Dendrobium comatum</i> (Blume) Lindl.	<i>Flickingeria</i>	Qld	described as <i>Flickingeria clemensii</i>
<i>Dendrobium convexum</i> (Blume) Lindl.	<i>Abaxianthus</i>	Qld	
( <i>Dendrobium</i> ) <i>crassum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	Qld	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium crumenatum</i> Sw.	<i>Ceraia</i>	AIT	Christmas Island; species name <i>saaronica</i> sometimes used
<i>Dendrobium cucumerinum</i> MacLeay ex Lindl.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium dicuphum</i> F.Muell.	<i>Vappodes</i>	WA NT	
<i>Dendrobium discolor</i> R.Br. subspecies <i>discolor</i>	<i>Durabaculum</i>	Qld	sometimes called <i>Dendrobium undulatum</i>
<i>Dendrobium discolor</i> R.Br. subspecies <i>incurvata</i> Liddle & P.I. Forst.	<i>Durabaculum</i>	Qld	
<i>Dendrobium discolor</i> R.Br. variety <i>broomfieldii</i> Fitzg.	<i>Durabaculum</i>	Qld	
<i>Dendrobium discolor</i> R.Br. variety <i>fimbrilabium</i> (Rchb.f.) Dockrill	<i>Durabaculum</i>	Qld	
<i>Dendrobium discolor</i> R.Br. variety <i>fuscum</i> (Fitzg.) Dockrill	<i>Durabaculum</i>	Qld	
<i>Dendrobium dolichophyllum</i> D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	NSW Qld	
( <i>Dendrobium</i> ) <i>eburneum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	NSW Qld	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
( <i>Dendrobium</i> ) <i>eungellensis</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	Qld	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium fairfaxii</i> F.Muell. ex Fitzg.	<i>Dockrillia</i>	NSW	
<i>Dendrobium falcorostrum</i> Fitzg.	<i>Thelychiton</i>	NSW Qld	
<i>Dendrobium fellowsii</i> F.Muell.	<i>Eleutheroglossum</i>	Qld	
<i>Dendrobium finniganense</i> D.L.Jones	<i>Thelychiton</i>	Qld	
<i>Dendrobium fleckeri</i> Rupp & C.T.White	<i>Thelychiton</i>	Qld	
<i>Dendrobium foelschei</i> F.Muell.	<i>Cepobaculum</i>	Qld NT	usually treated as a variety of <i>Dendrobium canaliculatum</i>
<i>Dendrobium funiforme</i> Blume	<i>Cadetia</i> , <i>Sarcocadetia</i>	Qld	formerly called <i>Cadetia wariana</i> & <i>Sarcocadetia wariana</i>
<i>Dendrobium glabrum</i> J.J.Sm.	<i>Diplocaulobium</i>	Qld	
<i>Dendrobium gracilicaule</i> F.Muell.	<i>Thelychiton</i>	NSW Qld	
<i>Dendrobium howeanum</i> (Maiden) D.L.Jones	<i>Thelychiton</i>	NSW	Lord Howe Island only; variously considered a variety of <i>Dendrobium gracilicaule</i> and <i>Dendrobium macropus</i>
<i>Dendrobium insigne</i> Blume	<i>Dichopus</i> , <i>Grastidium</i>	AIT	Dauan Island, Torres Strait
<i>Dendrobium johannis</i> Rchb.f.	<i>Cepobaculum</i>	Qld NT	

Australian Species	Alternate Genus	Distribution	Notes
<i>Dendrobium jonesii</i> Rendle subspecies <i>bancroftianum</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Thelychiton</i>	Qld	doubtfully distinct from subsp. <i>jonesii</i>
<i>Dendrobium jonesii</i> Rendle subspecies <i>blackburnii</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Thelychiton</i>	Qld	doubtfully distinct from subsp. <i>jonesii</i>
<i>Dendrobium jonesii</i> Rendle subspecies <i>jonesii</i>	<i>Thelychiton</i>	Qld	
<i>Dendrobium jonesii</i> Rendle subspecies <i>magnificum</i> (Dockrill) D.P.Banks & Clemesha	<i>Thelychiton</i>	Qld	usually called var. <i>magnificum</i>
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>kingianum</i>	<i>Thelychiton</i>	NSW Qld	several varieties have been named although these are recognised mainly in cultivation and not taxonomically; one variety is recognised here
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>kingianum</i> variety <i>pulcherrimum</i> Rupp	<i>Thelychiton</i>	NSW	considered a full species <i>D. pulcherrimum</i> by Jones (2006)
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>carnarvonense</i> P.B.Adams	<i>Thelychiton</i>	Qld	considered a full species <i>D. canarvonense</i> by Jones (2006)
<i>Dendrobium lichenastrum</i> (F.Muell.) Rolfe	<i>Davejonesia</i>	Qld	
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>huntianum</i> Rupp	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>linguiforme</i>	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium lithocola</i> D.L.Jones & M.A.Clem.	<i>Vappodes</i>	Qld	usually treated as a variety of <i>Dendrobium bigibbum</i>
<i>Dendrobium litorale</i> Schltr.	<i>Aporopsis</i>	AIT	Dauan Island, Torres Strait; also called <i>Dendrobium macfarlanei</i>
<i>Dendrobium lobbii</i> Teijsm. & Binn.	<i>Conostalix</i>	Qld NT	also called <i>Dendrobium paludica</i> Schltr.
<i>Dendrobium luteocilium</i> Rupp	<i>Grastidium</i>	Qld	sometimes included in <i>Dendrobium pruinosum</i> Teijsm. & Binn
<i>Dendrobium macropus</i> (Endl.) Rchb.f. ex Lindl.	<i>Thelychiton</i>	AIT	Norfolk Island
<i>Dendrobium maidenianum</i> Schltr.	<i>Cadetia</i>	Qld	
<i>Dendrobium malbrowii</i> Dockrill	<i>Monanthos</i>	Qld	
<i>Dendrobium melaleucaphilum</i> M.A.Clem. & D.L.Jones	<i>Tetrabaculum</i>	NSW	alternative name <i>Dendrobium tetragonum</i> subspecies <i>tetragonum</i> variety <i>melaleucaphilum</i>
<i>Dendrobium mirbelianum</i> Gaudich.	<i>Durabaculum</i>	Qld	
<i>Dendrobium monophyllum</i> F.Muell.	<i>Australorchis</i>	NSW Qld	
<i>Dendrobium moorei</i> F.Muell.	<i>Thelychiton</i>	NSW	Lord Howe Island only
<i>Dendrobium mortii</i> F.Muell.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium nativitatis</i> Ridl.	<i>Flickingeria</i>	AIT	Christmas Island
<i>Dendrobium nindii</i> W.Hill	<i>Durabaculum</i>	Qld	
<i>Dendrobium nitidum</i> F.M.Bailey	<i>Thelychiton</i>	Qld	a taxon of uncertain status; variously included in <i>Dendrobium speciosum</i> subs. <i>pedunculatum</i> var. <i>boreale</i> ; also considered a hybrid, a form of <i>Dendrobium</i> × <i>gracillimum</i>
<i>Dendrobium nugentii</i> (F.M.Bailey) D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	Qld	
<i>Dendrobium phalaenopsis</i> Fitzg.	<i>Vappodes</i>	Qld	usually treated as a variety of <i>Dendrobium bigibbum</i>
<i>Dendrobium prenticei</i> (F.Muell.) Nicholls	<i>Davejonesia</i>	Qld	sometimes included in <i>Dendrobium lichenastrum</i>
<i>Dendrobium pugioniforme</i> A.Cunn.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium racemosum</i> (Nicholls) Clemesha & Dockrill	<i>Dockrillia</i>	Qld	
( <i>Dendrobium</i> ) <i>radiata</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	NSW Qld	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium reconditum</i> Schuit. & P.B.Adams	<i>Cadetia</i>	AIT	Moa Island, Torres Strait; also called <i>Cadetia clausa</i>
<i>Dendrobium rigidum</i> R.Br.	<i>Dockrillia</i>	Qld	
<i>Dendrobium schneiderae</i> F.M.Bailey variety <i>majus</i> Rupp	<i>Australorchis</i>	Qld	renamed as <i>Australorchis eungellensis</i> D.L. Jones & M.A. Clem.; that epithet combination in <i>Dendrobium</i> not yet made
<i>Dendrobium schneiderae</i> F.M.Bailey variety <i>schneiderae</i>	<i>Australorchis</i>	NSW Qld	
<i>Dendrobium schoeninum</i> Lindl.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium semifuscum</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Cepobaculum</i>	Qld	sometimes included in <i>Dendrobium trilamellatum</i>
<i>Dendrobium smillieae</i> (F.Muell.) Fitzg.	<i>Coelandria</i>	Qld	
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>blackdownense</i> P.B.Adams	<i>Thelychiton</i>	Qld	alternative name (in part) <i>Dendrobium (Thelychiton) coriaceum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>capricornicum</i> Clemesha	<i>Thelychiton</i>	Qld	alternative name <i>Dendrobium (Thelychiton) capricornicum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>carnarvonense</i> P.B.Adams	<i>Thelychiton</i>	Qld	alternative name (in part) <i>Dendrobium (Thelychiton) coriaceum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>curvicaule</i> F.M.Bailey	<i>Thelychiton</i>	Qld	alternative name <i>Dendrobium (Thelychiton) spectabilis</i>



Australian Species	Alternate Genus	Distribution	Notes
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>grandiflorum</i> F.M.Bailey	<i>Thelychiton</i>	Qld	alternative name <i>Dendrobium (Thelychiton) rex</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>hillii</i> Masters	<i>Thelychiton</i>	NSW Qld	alternative name <i>Dendrobium (Thelychiton) tarberi</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>speciosum</i>	<i>Thelychiton</i>	Vic NSW	alternative name (in part) <i>Dendrobium (Thelychiton) epiphyticus</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>pedunculatum</i> variety <i>pedunculatum</i> (Clemesha) D.P.Banks & Clemesha	<i>Thelychiton</i>	Qld	alternative name <i>Dendrobium (Thelychiton) pedunculatum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>pedunculatum</i> variety <i>boreale</i> P.B.Adams, J.M.Burke and S.D.Lawson	<i>Thelychiton</i>	Qld	alternative names <i>Dendrobium (Thelychiton) biconvexum</i> , <i>D. (T.) curvicaule</i> , <i>D. (T.) rupicolum</i>
<i>Dendrobium stelliferum</i> J.J.Sm.	<i>Diplocaulobium</i>	Qld	Australian record doubtful
<i>Dendrobium striolatum</i> Rchb.f. subspecies <i>chrysanthum</i> D.L.Jones	<i>Dockrillia</i>	Tas	
<i>Dendrobium striolatum</i> Rchb.f. Rauschert subspecies <i>striolatum</i>	<i>Dockrillia</i>	Vic Tas NSW	
<i>Dendrobium stuartii</i> F.M.Bailey		Qld	sometimes included in <i>Dendrobium macrostachyum</i> Lindl.
<i>Dendrobium sulphureum</i> Schltr.	<i>Dockrillia</i>	Qld	
<i>Dendrobium tattonianum</i> Bateman ex Rchb.f.		Qld	sometimes included in <i>Dendrobium canaliculatum</i>
<i>Dendrobium taylori</i> (F.Muell.) F.M.Bailey	<i>Cadetia</i>	Qld	
<i>Dendrobium teretifolium</i> R.Br.	<i>Dockrillia</i>	NSW Qld	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>cataractum</i> P.B.Adams, S.D.Lawson and G.A.Paterson	<i>Tetrabaculum</i>	Qld	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>giganteum</i> variety <i>giganteum</i> (Leaney) P.B.Adams	<i>Tetrabaculum</i>	Qld	alternative name <i>Dendrobium (Tetrabaculum) capitisyork</i>
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>giganteum</i> variety <i>cacatua</i> M.A.Clem. & D.L.Jones	<i>Tetrabaculum</i>	Qld	alternative name <i>Dendrobium (Tetrabaculum) cacatua</i>
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>tetragonum</i> variety <i>serpentis</i> P.B.Adams	<i>Tetrabaculum</i>	Qld	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>tetragonum</i> variety <i>tetragonum</i>	<i>Tetrabaculum</i>	NSW Qld	
<i>Dendrobium toressae</i> (F.M.Bailey) Dockrill	<i>Stilbophyllum</i>	Qld	
<i>Dendrobium tozerense</i> Lavarack	<i>Grastidium</i>	Qld	
<i>Dendrobium trilamellatum</i> J.J.Sm.	<i>Cepobaculum</i>	Qld NT	
<i>Dendrobium uniflos</i> F.M.Bailey	<i>Cadetia</i>	Qld	possibly a synonym of <i>Dendrobium taylori</i>
<i>Dendrobium wassellii</i> S.T.Blake	<i>Dockrillia</i>	Qld	
<b><i>Didymoplexis</i> Griff.</b>			
<i>Didymoplexis pallens</i> Griffith		WA Qld NT AIT	also on Christmas Island
<b><i>Dienia</i> Lindl.</b>			
<i>Dienia montana</i> (Blume) M.A.Clem. & D.L.Jones	<i>Malaxis</i>	Qld NT	
<b><i>Dipodium</i> R.Br.</b>			
<i>Dipodium atropurpureum</i> D.L.Jones		NSW Qld	
<i>Dipodium campanulatum</i> D.L.Jones		SA Vic	
<i>Dipodium elegantulum</i> D.L.Jones		Qld	
<i>Dipodium ensifolium</i> F.Muell.		Qld	
<i>Dipodium hamiltonianum</i> F.M.Bailey		Vic NSW ACT Qld	
<i>Dipodium pandanum</i> F.M.Bailey		Qld	
<i>Dipodium pardalinum</i> D.L.Jones		SA Vic	
<i>Dipodium pictum</i> (Lindl.) Rchb.f.		Qld	taxon of uncertain status
<i>Dipodium pulchellum</i> D.L.Jones & M.A.Clem		NSW Qld	
<i>Dipodium punctatum</i> (Sm.) R.Br.		SA Vic NSW ACT Qld	
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		SA Vic Tas NSW ACT	
<i>Dipodium stenochilum</i> O.Schwarz		WA Qld NT	it is likely that plants identified as <i>D. stenochilum</i> in WA will be redetermined as <i>Dipodium</i> species 'basalt woodland' and <i>Dipodium</i> species 'sandstone'
<i>Dipodium variegatum</i> M.A.Clem. & D.L.Jones		Vic NSW Qld	
<i>Dipodium</i> species 'basalt woodland'		WA	aff. <i>roseum</i> with spotted deep pink flowers with a rose pink labellum; woodland in the Kimberley region; <i>basalticum</i> ms (Brown <i>et al.</i> 2013)

Australian Species	Alternate Genus	Distribution	Notes
<i>Dipodium</i> species 'sandstone'		WA	aff. <i>roseum</i> with unspotted pale pink flowers with a rose pink labellum; woodland over sandstone in the Kimberly region; <i>ammolithum</i> ms (Brown <i>et al.</i> 2013)
<b><i>Diteilis</i> Raf.</b>			
<i>Diteilis collinsii</i> (B.Gray) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<i>Diteilis petricola</i> D.L.Jones & B.Gray	<i>Liparis</i>	Qld	
<i>Diteilis simmondsii</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	Qld	
<b><i>Diuris</i> Sm</b>			
<i>Diuris abbreviata</i> Benth.		NSW Qld	
<i>Diuris aequalis</i> F.Muell. ex Fitzg.		NSW	
<i>Diuris alba</i> R.Br.		NSW Qld	
<i>Diuris amplissima</i> D.L.Jones		WA	
<i>Diuris arenaria</i> D.L.Jones		NSW	
<i>Diuris aurea</i> Sm.		NSW Qld	Qld records possibly erroneous, perhaps misidentifications of <i>Diuris chrysantha</i>
<i>Diuris basaltica</i> D.L.Jones		Vic	
<i>Diuris behrii</i> Schldl.		SA Vic NSW ACT	
<i>Diuris bracteata</i> Fitzg.		NSW	
<i>Diuris brevifolia</i> R.S.Rogers		SA	
<i>Diuris brevissima</i> Fitzg. ex Nicholls		NSW	included on PlantNET (NSW Flora Online) but status doubtful, usually included in <i>Diuris maculata</i>
<i>Diuris brumalis</i> D.L.Jones		WA	
<i>Diuris byronensis</i> D.L.Jones		NSW	
<i>Diuris callitrophila</i> D.L.Jones		NSW	
<i>Diuris calcicola</i> R.J.Bates		SA	
<i>Diuris carinata</i> Lindl.		WA	
<i>Diuris chrysantha</i> D.L.Jones & M.A.Clem.		NSW Qld	
<i>Diuris chryseopsis</i> D.L.Jones		SA Vic Tas NSW ACT	
<i>Diuris concinna</i> D.L.Jones		WA	
<i>Diuris conspicillata</i> D.L.Jones		WA	
<i>Diuris corymbosa</i> Lindl.		WA	
<i>Diuris cuneilabris</i> Rupp		NSW	included on Atlas of Living Australia but status doubtful, usually included in <i>Diuris platichila</i>
<i>Diuris curta</i> D.L.Jones		NSW Qld	
<i>Diuris daltonii</i> (C.Walter) D.L.Jones & M.A.Clem.		SA Vic	probably the same as <i>Diuris punctata</i> , from which it is poorly differentiated
<i>Diuris decrementa</i> D.L.Jones & C.J.French		WA	
<i>Diuris dendrobioides</i> Fitzg.		Vic NSW ACT	
<i>Diuris disposita</i> D.L.Jones		NSW	
<i>Diuris drummondii</i> Lindl.		WA	
<i>Diuris eborensis</i> D.L.Jones		NSW	
<i>Diuris eburnea</i> D.L.Jones		WA	
<i>Diuris emarginata</i> R.Br.		WA	
<i>Diuris exitela</i> D.L.Jones		Qld	
<i>Diuris filifolia</i> Lindl.		WA	
<i>Diuris flavescens</i> D.L.Jones		NSW	
<i>Diuris fragrantissima</i> D.L.Jones & M.A.Clem.		Vic	
<i>Diuris fucosa</i> D.L.Jones		Vic NSW	
<i>Diuris goonooensis</i> Rupp		NSW Qld	called <i>Diuris maculata</i> in Qld
<i>Diuris gregaria</i> D.L.Jones		Vic	
<i>Diuris hazeliae</i> D.L.Jones & C.J.French		WA	
<i>Diuris heberlei</i> D.L.Jones		WA	
<i>Diuris immaculata</i> D.L.Jones		WA	
<i>Diuris insignis</i> D.L.Jones & C.J.French		WA	
<i>Diuris jonesii</i> C.J.French & G.Brockman		WA	
<i>Diuris laevis</i> Fitzg.		WA	
<i>Diuris lanceolata</i> Lindl.		Tas	
<i>Diuris laxiflora</i> Lindl.		WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Diuris longifolia</i> R.Br.		WA	
<i>Diuris luteola</i> D.L.Jones & B.Gray		Qld	
<i>Diuris maculata</i> Sm.		NSW	
<i>Diuris magnifica</i> D.L.Jones		WA	
<i>Diuris micrantha</i> D.L.Jones		WA	
<i>Diuris minor</i> (Benth.) D.L.Jones & M.A.Clem.		NSW	
<i>Diuris monticola</i> D.L.Jones		Vic Tas NSW ACT	
<i>Diuris nigromontana</i> D.L.Jones		NSW ACT	
<i>Diuris ochroma</i> D.L.Jones		Vic NSW	
<i>Diuris oporina</i> D.L.Jones		Qld	
<i>Diuris orientis</i> D.L.Jones		SA Vic Tas NSW	
<i>Diuris palustris</i> Lindl.		SA Vic Tas	
<i>Diuris pardina</i> Lindl.		SA Vic Tas NSW ACT	
<i>Diuris parvipetala</i> (Dockrill) D.L.Jones & M.A.Clem.		NSW Qld	
<i>Diuris pauciflora</i> R.Br.		WA	
<i>Diuris pedunculata</i> R.Br.		NSW Qld	
<i>Diuris perialla</i> D.L.Jones & C.J.French		WA	
<i>Diuris picta</i> J. Drummond		WA	
<i>Diuris platichila</i> Fitzg.		NSW	
<i>Diuris porrifolia</i> Lindl.		WA	
<i>Diuris praecox</i> D.L.Jones		NSW	
<i>Diuris protena</i> D.L.Jones		Vic	
<i>Diuris pulchella</i> D.L.Jones		WA	
<i>Diuris punctata</i> Sm. variety <i>punctata</i>		Vic NSW ACT Qld	Qld records are most likely attributable to several undescribed species including 'granite ranges' and 'NSW North Coast'; possibly also misidentifications of <i>D. alba</i> , <i>D. oporina</i> and <i>D. parvipetala</i>
<i>Diuris punctata</i> Sm. variety <i>sulfurea</i> Rupp		NSW	most likely a hybrid between <i>D. punctata</i> and a yellow-flowered <i>Diuris</i> species
<i>Diuris purdiei</i> Diels		WA	
<i>Diuris recurva</i> D.L.Jones		WA	
<i>Diuris refracta</i> D.L.Jones & C.J.French		WA	
<i>Diuris secundiflora</i> Fitzg.		NSW	
<i>Diuris segregata</i> D.L.Jones & C.J.French		WA	
<i>Diuris semilunulata</i> Messmer		Vic NSW ACT	
<i>Diuris septentrionalis</i> D.L.Jones & C.J.French		WA	
<i>Diuris setacea</i> R.Br.		WA	
<i>Diuris striata</i> Rupp		NSW	taxon of uncertain identity, possibly a form of <i>Diuris dendrobioides</i> with short lateral sepals
<i>Diuris subalpina</i> D.L.Jones		Vic NSW ACT	
<i>Diuris sulphurea</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Diuris systema</i> D.L.Jones & L.M.Copel.		NSW	
<i>Diuris tinkeri</i> D.L.Jones & C.J.French		WA	
<i>Diuris tricolor</i> Fitzg.		Vic NSW Qld	
<i>Diuris unica</i> D.L.Jones		NSW Qld	
<i>Diuris venosa</i> Rupp		NSW	
<i>Diuris</i> species 'Augusta'		WA	late-flowering aff. <i>laxiflora</i> with bright yellow flowers with a broader labellum mid-lobe; amongst sedges in seasonally damp habitats, flowering best after fire; only near Augusta (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Byfield'		Qld	no information available (Queensland Herbarium record)
<i>Diuris</i> species 'Carnarvon'		Qld	aff. <i>punctata</i> ; sparse woodland on sandstone at Mt Moffat & Carnarvon Range; AVH records listed as <i>D. punctata</i> but almost certainly not this species; <i>Diuris parvipetala</i> is also recorded from the region
<i>Diuris</i> species 'Clarencetown'		NSW	aff. <i>alba</i> with smaller more uniformly coloured flowers with erect dorsal sepals; shrubby open forest near Clarencetown (Bishop 2000)
<i>Diuris</i> species 'Colo River'		NSW	aff. <i>punctata</i> with smaller flowers with short lateral sepals and reflexed petals and creamy yellow between labellum callus ridges; swamp margins in heath, heathy & grassy woodland Mellong Range to Hunter Valley (Bishop 2000)

Australian Species	Alternate Genus	Distribution	Notes
<i>Diuris</i> species 'Darling scarp'		WA	late spring flowering, tall aff. <i>amplissima</i> with large brownish-yellow and mauve flowers (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Ebor'		NSW	tall aff. <i>dendrobioides</i> with larger, darker, more uniformly coloured flowers with more erect dorsal sepals; montane grassland and grassy woodland near Ebor (Copeland 2005)
<i>Diuris</i> species 'Gippsland plains'		Vic	small aff. <i>dendrobioides</i> with blotched flowers; plains grasslands in central Gippsland (Jeanes & Backhouse 2006)
<i>Diuris</i> species 'Grafton'		NSW	early spring-flowering aff. <i>dendrobioides</i> with smaller paler flowers with shorter lateral sepals; grassy open forest near Grafton (L. Copeland unpublished)
<i>Diuris</i> species 'Green Range'		WA	winter flowering aff. <i>corymbosa</i> with colourful flowers; heaths and heathy woodlands on south coast between Denmark and Esperance (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Guyra'		NSW	small aff. <i>dendrobioides</i> with mainly white flowers; montane grasslands above 1,200 m on New England tableland (Copeland 2005)
<i>Diuris</i> species 'Hunter Valley'		NSW	very small aff. <i>dendrobioides</i> with rounded petals not tapering into the petiole; grassy open forest in the central Hunter Valley (Bishop 2000)
<i>Diuris</i> species 'Jollys Falls'		Qld	no information available (Qld Herbarium record)
<i>Diuris</i> species 'Kemerton'		WA	aff. <i>corymbosa</i> with yellow and red-brown flowers; coastal woodland on sandy soils between Lake Clifton and Capel (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'mid-north'		WA	small, often clumping aff. <i>corymbosa</i> with pale flowers with recurved dorsal sepal and crossed, recurved lateral sepals; heath and woodland between Moora and Mingenew (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Monaro'		NSW	small late spring-early summer flowering aff. <i>dendrobioides</i> with indistinct blotches and yellow between callus ridges; native grasslands on the Monaro tableland (Bishop 2000)
<i>Diuris</i> species 'Murchison River'		WA	tall aff. <i>corymbosa</i> with mainly yellow flowers with recurved petals and sepals; moist sedgeland near Murchison River (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'New England escarpment'		NSW	spring flowering aff. <i>alba</i> with colourful flowers with broad, reflexed petals and yellow tubers; heathy woodland and open forest on granite hills on the New England Tableland (Bishop 2000)
<i>Diuris</i> species 'New England swamps'		NSW	summer flowering aff. <i>alba</i> with recurved petals; swamps on New England Tableland (Bishop 2000)
<i>Diuris</i> species 'NSW North Coast'		NSW	short, robust aff. <i>punctata</i> with large paler flowers with broad dorsal sepal, petals & labellum, white/cream between labellum callus ridges; heaths and grassy woodland on coastal headlands from Kempsey to Yamba (Bishop 2000)
<i>Diuris</i> species 'Perth swamps'		WA	small, short aff. <i>corymbosa</i> with widely spaced flowers with a short labellum midlobe; swamps just south of Perth (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Red Gum waterholes'		SA	short aff. <i>behrii</i> with small pale sometimes brown-streaked flowers with a depressed dorsal sepal and short labellum; damp, sometimes swampy sites in grassy woodland in the lower south-east (Bates 2011)
<i>Diuris</i> species 'sandhills'		SA Vic	aff. <i>chryseopsis</i> with narrower petals, labellum & narrow black-striped dorsal sepal; heathy woodland on grey sandy rises in the lower south-east of SA & far western Vic (Bates 2011)
<i>Diuris</i> species 'sandplain'		WA	tall, colourful aff. <i>corymbosa</i> ; sandplain habitats between Mandurah and Bunbury (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Scaddan'		WA	small, winter flowering aff. <i>corymbosa</i> ; woodland on rocky outcrops between Hyden and Scaddan; included in Brown <i>et al.</i> 2008 but not in later books
<i>Diuris</i> species 'south coast'		WA	winter-flowering aff. <i>corymbosa</i> with small, mainly yellow flowers; coastal heaths and heathy woodland between Munglinup and Denmark (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Tenterfield/Stanthorpe'		NSW Qld	aff. <i>punctata</i> with pale mottled flowers with relatively short lateral sepals and white between the labellum callus ridges; open forest in the granite ranges around Tenterfield NSW and Stanthorpe Qld
<i>Diuris</i> species 'thick sepals'		Vic	late flowering aff. <i>pardina</i> with shorter, thicker, less curved to straight lateral sepals; dry woodland and open forest in western Vic & probably SA (Jeanes & Backhouse 2006); may be same as <i>D. calcicola</i> R.J. Bates
<i>Diuris</i> species 'western wheatbelt'		WA	short aff. <i>corymbosa</i> with pale yellow flowers; woodlands and rocky areas between York and Ravensthorpe (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Williams'		WA	small aff. <i>corymbosa</i> ; <i>Allocasuarina</i> woodland thickets near Williams; included in Brown <i>et al.</i> 2008 but not in later books
<i>Diuris</i> species 'Wyalkatchem'		WA	early spring-flowering aff. <i>corymbosa</i> with yellow flowers with small red patches; inland woodlands on sandy soils between Bencubbin and Wyalkatchmen (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Yalgorup'		WA	tall colourful aff. <i>corymbosa</i> ; woodland and open forest between and Mandurah and Bunbury (Brown <i>et al.</i> 2013)



Australian Species	Alternate Genus	Distribution	Notes
<i>Diuris</i> species 'Zuytdorp cliffs'		WA	smallish, compact aff. <i>corymbosa</i> with mainly yellow flowers with short lateral sepals; coastal sedges on limestone sands around Kalbarri (Brown <i>et al.</i> 2013)
<b><i>Drakaea</i> Lindl</b>			
<i>Drakaea andrewsiae</i> Hopper & A.P.Br.		WA	
<i>Drakaea concolor</i> Hopper & A.P.Br.		WA	
<i>Drakaea confluens</i> Hopper & A.P.Br.		WA	
<i>Drakaea elastica</i> Lindl.		WA	
<i>Drakaea glyptodon</i> Fitzg.		WA	
<i>Drakaea gracilis</i> Hopper & A.P.Br.		WA	
<i>Drakaea isolata</i> Hopper & A.P.Br.		WA	
<i>Drakaea livida</i> J.Drummond		WA	
<i>Drakaea micrantha</i> Hopper & A.P.Br.		WA	
<i>Drakaea thynniphila</i> A.S.George		WA	
<b><i>Drymoanthus</i> Nicholls</b>			
<i>Drymoanthus minutus</i> Nicholls		Qld	
<b><i>Elythranthera</i> (Endl.) A.S.George</b>			
<i>Elythranthera brunonis</i> (Endl.) A.S.George	<i>Caladenia</i>	WA	
<i>Elythranthera emarginata</i> (Lindl.) A.S.George	<i>Caladenia</i>	WA	
<b><i>Empusa</i> Lindl</b>			
<i>Empusa habenarina</i> (F.Muell.) Benth.	<i>Liparis</i>	WA NSW Qld NT	
<b><i>Epiblema</i> R.Br.</b>			
<i>Epiblema grandiflorum</i> R.Br.		WA	
<b><i>Epipogium</i> J.F.Gmel. ex Borkh.</b>			
<i>Epipogium roseum</i> (D.Don) Lindl.		NSW Qld	
<b><i>Ericksonella</i> Hopper &amp; A.P.Br.</b>			
<i>Ericksonella saccharata</i> (Rchb.f.) Hopper & A.P.Br.	<i>Caladenia</i>	WA	
<b><i>Eriochilus</i> R.Br.</b>			
<i>Eriochilus collinus</i> D.L. Jones & R.J.Bates		SA	
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		SA Vic Tas NSW ACT Qld	<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>brevifolius</i> (Benth.) Hopper & A.P.Brown		WA	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>dilatatus</i>		WA	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>magnus</i> Hopper & A.P.Br.		WA	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>multiflorus</i> Hopper & A.P.Br.		WA	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>orientalis</i> Lindl.		WA	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>undulatus</i> Hopper & A.P.Br.		WA	
<i>Eriochilus helonomos</i> Hopper & A.P.Br.		WA	
<i>Eriochilus magenteus</i> D.L.Jones		Vic NSW ACT	
<i>Eriochilus paludosus</i> D.L. Jones & R.J.Bates		SA	
<i>Eriochilus petricola</i> D.L.Jones & M.A.Clem.		NSW Qld	
<i>Eriochilus pulchellus</i> Hopper & A.P.Br.		WA	
<i>Eriochilus scaber</i> Lindl. subspecies <i>orbifolia</i> Hopper & A.P.Br.		WA	
<i>Eriochilus scaber</i> Lindl. subspecies <i>scaber</i>		WA	
<i>Eriochilus tenuis</i> Lindl.		WA	
<i>Eriochilus valens</i> Hopper & A.P.Br.		WA	
<i>Eriochilus</i> species 'dry plains'		SA	aff. <i>cucullatus</i> with soft deciduous hairs and minor ribbing on leaf, which is mostly green above, paler below; limestone plains of SA from west coast to Vic border (R. Bates unpublished)
<i>Eriochilus</i> species 'smooth leaf'		SA Vic NSW ACT Qld	aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<b><i>Erythrorchis</i> Blume</b>			
<i>Erythrorchis cassythoides</i> (A.Cunn. ex Lindl.) Garay		NSW Qld	

Australian Species	Alternate Genus	Distribution	Notes
<b>Eulophia</b> R.Br.			
<i>Eulophia agrostophylla</i> Benth.		Qld NT	may be the same as and therefore a synonym of <i>Eulophia venosa</i>
<i>Eulophia bicallosa</i> (D.Don) P.F.Hunt & Summerh.		WA Qld	
<i>Eulophia pelorica</i> D.L.Jones & M.A.Clem.		Qld	
<i>Eulophia venosa</i> (F.Muell.) Benth.		Qld NT	
<i>Eulophia zollingeri</i> (Rchb.f.) J.J.Sm.		Qld	
<b>Gastrodia</b> R.Br.			
<i>Gastrodia crebriflora</i> D.L.Jones		Qld	
<i>Gastrodia entomogama</i> D.L.Jones		NSW ACT	
<i>Gastrodia lacista</i> D.L.Jones		WA	
<i>Gastrodia procera</i> G.W.Carr		SA Vic Tas NSW	
<i>Gastrodia sesamoides</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Gastrodia surcula</i> D.L.Jones		NSW ACT	
<i>Gastrodia urceolata</i> D.L.Jones		Qld	
<i>Gastrodia vescula</i> D.L.Jones		SA	
<b>Genoplesium</b> R.Br.			
<i>Genoplesium baueri</i> R.Br.		NSW	
<b>Geodorum</b> Jacks.			
<i>Geodorum densiflorum</i> (Lam.) Schltr.		WA NSW Qld NT	also called <i>Geodorum terrestre</i> (L.) Garay
<i>Geodorum neocaledonicum</i> Kraenzl.		WA	record most likely false, probably resulting from an incorrect determination of <i>Eulophia bicallosa</i>
<b>Glossodia</b> R.Br.			
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	SA Vic Tas NSW ACT Qld	
<i>Glossodia minor</i> R.Br.	<i>Caladenia</i>	Vic NSW Qld	alternative name <i>Caladenia minorata</i> M.A. Clem.
<b>Goodyera</b> R.Br.			
<i>Goodyera grandis</i> (Blume) Blume		Qld	
<i>Goodyera viridiflora</i> (Blume) Blume		Qld	also called <i>Eucosia umbrosa</i> D.L.Jones & M.A.Clem.
<b>Habenaria</b> Willd.			
<i>Habenaria chlorosepala</i> D.L.Jones		Qld	
<i>Habenaria elongata</i> R.Br.		WA Qld NT	
<i>Habenaria euryloba</i> D.L.Jones		Qld	
<i>Habenaria exilis</i> D.L.Jones		Qld	
<i>Habenaria ferdinandii</i> Schltr.		NT	
<i>Habenaria fuscina</i> D.L.Jones		Qld	
<i>Habenaria halata</i> D.L.Jones		NT	
<i>Habenaria harroldii</i> D.L.Jones		Qld	
<i>Habenaria hymenophylla</i> Schltr.		WA Qld NT	
<i>Habenaria macraithii</i> Lavarack		Qld	
<i>Habenaria ochroleuca</i> R.Br.		WA NT	
<i>Habenaria praecox</i> Lavarack & Dockrill		Qld	
<i>Habenaria propinquior</i> Rchb.f.		Qld	
<i>Habenaria rumphii</i> (Brogn.) Lindl.		Qld NT	
<i>Habenaria triplonema</i> Schltr.		Qld NT	
<i>Habenaria vatia</i> D.L.Jones		AIT	Moa Island, Torres Strait
<i>Habenaria xanthantha</i> F.Muell.		Qld	
<i>Habenaria</i> species 'Beverley Springs Station'		WA	aff. <i>praecox</i> with a bluntly pointed, tri-lobed labellum; damp sites in woodland in the Kimberley region; <i>joesmithii</i> ms (Brown <i>et al.</i> 2013)
<b>Hetaeria</b> Blume			
<i>Hetaeria oblongifolia</i> Blume		Qld	
<b>Leporella</b> A.S.George			
<i>Leporella fimbriata</i> (Lindl.) A.S.George		WA SA Vic	
<b>Leptoceras</b> (R.Br.) Lindl.			
<i>Leptoceras menziesii</i> (R.Br.) Lindl.	<i>Caladenia</i>	WA SA Vic Tas	
<b>Luisia</b> Gaudich.			
<i>Luisia atacta</i> D.L.Jones		Qld	called <i>Luisia teretifolia</i> in Qld
<i>Luisia corrugata</i> D.L.Jones		NT	

Australian Species	Alternate Genus	Distribution	Notes
<b>Lyperanthus</b> R.Br.			
<i>Lyperanthus serratus</i> Lindl.		WA	
<i>Lyperanthus suaveolens</i> R.Br.		Vic Tas NSW ACT Qld	
<b>Micropera</b> Lindl.			
<i>Micropera fasciculata</i> (Lindl.) Garay		Qld	
<b>Microtis</b> R.Br.			
<i>Microtis alba</i> R.Br.		WA	
<i>Microtis alboviridis</i> R.J.Bates		WA	
<i>Microtis angusii</i> D.L.Jones		NSW	
<i>Microtis arenaria</i> Lindl.		SA Vic Tas	
<i>Microtis atrata</i> Lindl.	<i>Microtidium</i>	WA SA Vic Tas	
<i>Microtis brownii</i> Rchb.f.		WA	
<i>Microtis cupularis</i> (D.L.Jones & G.Brockman) A.P.Br.	<i>Hydrorchis</i>	WA	
<i>Microtis eremaea</i> R.J.Bates		WA SA	
<i>Microtis eremicola</i> (R.J.Bates) D.L.Jones & M.A.Clem.		WA	
<i>Microtis familiaris</i> R.J.Bates		WA	
<i>Microtis frutetorum</i> Schldl.		SA	
<i>Microtis globula</i> R.J.Bates		WA	
<i>Microtis gracilentia</i> (R.J.Bates) R.J.Bates		SA	
<i>Microtis graniticola</i> R.J.Bates		WA	
<i>Microtis media</i> R.Br. subspecies <i>densiflora</i> R.J.Bates		WA	
<i>Microtis media</i> R.Br. subspecies <i>media</i>		WA	
<i>Microtis oblonga</i> R.S.Rogers		SA Vic Tas NSW ACT	
<i>Microtis orbicularis</i> R.S.Rogers	<i>Hydrorchis</i>	WA SA Vic Tas	
<i>Microtis parviflora</i> R.Br.		SA Vic Tas NSW ACT Qld	also on Lord Howe Island
<i>Microtis pulchella</i> R.Br.		WA	
<i>Microtis quadrata</i> (R.J.Bates) D.L.Jones & M.A.Clem.		WA	
<i>Microtis rara</i> R.Br.		SA NSW Qld	identity and distribution uncertain, thought to be only NSW & Qld (Jones 2006), although also claimed to be in Sth Aust (Bates 2011)
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		SA Vic Tas NSW ACT Qld AIT	also on Lord Howe Island, Norfolk Island
<i>Microtis</i> species 'alpine'		Vic NSW ACT	aff. <i>unifolia</i> with fewer smaller spaced flowers with a smaller labellum; wet mountain meadows and woodland (Jones <i>et al.</i> 2008)
<i>Microtis</i> species 'curved lip'		SA	aff. <i>unifolia</i> with a blunt dorsal sepal and curved labellum; swampy habitats in central SA, flowering after fire; <i>curvichila</i> ms (Bates 2011)
<i>Microtis</i> species 'dense spikes'		SA	aff. <i>parviflora</i> with a dense spike of numerous flowers with a tiny labellum; swampy habitats in sth Mt Lofty Range; <i>brevichila</i> ms (Bates 2011); probably only a subspecies of <i>parviflora</i>
<i>Microtis</i> species 'Flinders Ranges'		SA	aff. <i>eremaea</i> with short slender spike of tiny congested blue-green flowers with fat rounded ovary; dry ridges of Flinders Ranges; <i>flindersica</i> ms (Bates 2011)
<i>Microtis</i> species 'fragrant'		SA Vic NSW	aff. <i>arenaria</i> with sparser, scented flowers with a longer labellum with a raised central ridge; widespread in many habitats in south-eastern Australis; <i>fragrans</i> ms (Bates 2011)
<i>Microtis</i> species 'hanging bogs'		SA	aff. <i>oblonga</i> with tall spike of well-spaced flowers with a broad dorsal sepal; peat bogs in the Adelaide Hills SA (Bates 2011); probably only a form of <i>oblonga</i>
<i>Microtis</i> species 'late flowering'		SA	late flowering aff. <i>unifolia</i> with glossy flowers with a quadrate, unnotched labellum; woodland in higher parts of the Adelaide Hills; <i>postremus</i> ms (Bates 2011)
<i>Microtis</i> species 'narrow lip'		SA Vic	aff. <i>oblonga</i> with a long, narrow labellum constricted in the middle; tea-tree swamps on black alkaline mud in sth-east SA, also Vic?; <i>leptochila</i> ms (Bates 2011)
<i>Microtis</i> species 'subalpine grasslands'		ACT	short, summer flowering aff. <i>parviflora</i> with fewer smaller flowers that often lack the labellum apical callus; damp subalpine grasslands (Jones <i>et al.</i> 2008)

Australian Species	Alternate Genus	Distribution	Notes
<i>Microtis</i> species 'short leaf'		SA	small slender aff. <i>unifolia</i> with short thick leaf & well spaced flowers with a long slender labellum; widespread; <i>brevis</i> ms (Bates 2011)
<i>Microtis</i> species 'Topperwein'		SA Vic	aff. <i>oblonga</i> with tall spike of larger, well-spaced fleshy flowers; River Red Gum swamps in sth-east SA & west Vic, flowering without fire (Bates 2011)
<i>Microtis</i> species 'Windy Harbour'	<i>Hydrorchis</i>	WA	summer flowering aff. <i>orbicularis</i> with smaller, more crowded reddish flowers with cupped labellum; seasonal swamps between Northcliff and Windy Harbour (Brown <i>et al.</i> 2013)
<b>Mobilabium</b> Rupp			
<i>Mobilabium hamatum</i> Rupp		Qld	
<b>Nervilia</b> Comm. ex Gaudich.			
<i>Nervilia aragoana</i> Gaud.		Qld NT	
<i>Nervilia crociformis</i> (Zoll. & Mor.) Seidenf.		Qld NT	
<i>Nervilia holochila</i> (F.Muell.) Schltr.		WA Qld NT	
<i>Nervilia peltata</i> B.Gray & D.L.Jones		Qld NT	
<i>Nervilia plicata</i> (Andrews) Schltr.		Qld NT	also called <i>Nervilia dallachyana</i> (F.Muell. ex. Benth.)
<i>Nervilia uniflora</i> (F.Muell.) Schltr.		Qld NT	
<b>Oberonia</b> Lindl.			
<i>Oberonia attenuata</i> Dockrill		Qld	
<i>Oberonia carnosa</i> Lavarack		Qld	
<i>Oberonia complanata</i> (A.Cunn.) M.A.Clem. & D.L.Jones		NSW Qld	
<i>Oberonia crateriformis</i> D.L.Jones & M.A.Clem.		Qld	
<i>Oberonia flavescens</i> D.L.Jones & M.A.Clem.		Qld	
<i>Oberonia palmicola</i> F.Muell.		NSW Qld	also included in <i>Oberonia titania</i>
<i>Oberonia rimachila</i> D.L.Jones & M.A.Clem.		Qld	
<i>Oberonia titania</i> Lindl.		AIT	Norfolk Island
<b>Octarrhena</b> Thwaites			
<i>Octarrhena pusilla</i> (F.M.Bailey) M.A.Clem. & D.L.Jones		Qld	
<b>Orthoceras</b> R.Br.			
<i>Orthoceras strictum</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Orthoceras</i> species 'short bracts'		Vic	aff. <i>strictum</i> with short flower bracts, rounded labellum apex; heathy woodland Otway Ranges; resembles New Zealand species <i>O. novae-zeelandiae</i> (Jeanes & Backhouse 2006); grows with and probably only a variant of typical <i>O. strictum</i>
<b>Pachystoma</b> Blume			
<i>Pachystoma pubescens</i> Blume		Qld NT	
<b>Papillilabium</b> Dockrill			
<i>Papillilabium beckeri</i> (F.Muell. ex Benth.) Dockrill		NSW Qld	
<b>Paracaleana</b> Blaxell			
<i>Paracaleana alcockii</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana brockmanii</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana disjuncta</i> D.L.Jones	<i>Caleana</i>	WA SA Vic	
<i>Paracaleana dixonii</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana gracilicordata</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana granitica</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana hortiorum</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana lyonsii</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	SA Vic Tas NSW ACT Qld	
<i>Paracaleana nigrita</i> J. Drummond ex Lindl.	<i>Caleana</i>	WA	
<i>Paracaleana parvula</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana terminalis</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana triens</i> Hopper & A.P.Br.	<i>Caleana</i>	WA	
<i>Paracaleana</i> species 'Darling Range'	<i>Caleana</i>	WA	aff. <i>nigrita</i> with leaf still green at flowering and flat labellum with calli covering apical half; grows with <i>P. brockmanii</i> but flowers earlier and has smaller flower; woodland on lateritic soils near Kelmscott (Brown <i>et al.</i> 2013)
<b>Peristeranthus</b> T.E.Hunt			
<i>Peristeranthus hillii</i> (F.Muell.) T.E.Hunt		NSW Qld	



Australian Species	Alternate Genus	Distribution	Notes
<b>Peristylus</b> Blume			
<i>Peristylus banfieldii</i> (F.M.Bailey) Lavarack		Qld	
<i>Peristylus candidus</i> J.J.Sm.		Qld	
<i>Peristylus chlorandrellus</i> D.L.Jones & M.A.Clem.		Qld	
<i>Peristylus papuanus</i> (Kraenzl.) J.J.Sm.		Qld	
<b>Phaius</b> Lour.			
<i>Phaius amboinensis</i> Blume		NT	
<i>Phaius australis</i> F.Muell.		NSW Qld	
<i>Phaius bernaysii</i> Rowland ex Rchb.f.		Qld	
<i>Phaius pictus</i> T.E.Hunt		Qld	
<b>Phalaenopsis</b>			
<i>Phalaenopsis rosenstromii</i> F.M.Bailey		Qld	also called <i>P. amabilis</i> subspecies <i>rosenstromii</i>
<b>Pheladenia</b> D.L.Jones & M.A.Clem.			
<i>Pheladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	WA SA Vic Tas NSW	
<b>Pholidota</b> Lindl. ex Hook.			
<i>Pholidota imbricata</i> Hook	<i>Colelogyne</i>	Qld	
<b>Phreatia</b> Lindl.			
<i>Phreatia paleata</i> (Rchb.f.) Rchb.f.		AIT	Norfolk Island
<b>Pinalia</b> Lindl.			
<i>Pinalia fitzalanii</i> (F.Muell.) Kuntze	<i>Eria, Hymeneria</i>	Qld	
<i>Pinalia kingii</i> (F.Muell.) Kuntze	<i>Eria, Hymeneria</i>	Qld	
<b>Plectorrhiza</b> Dockrill			
<i>Plectorrhiza brevilabris</i> (F.Muell.) Dockrill		Qld	
<i>Plectorrhiza erecta</i> (Fitzg.) Dockrill		NSW	Lord Howe Island only
<i>Plectorrhiza tridentata</i> (Lindl.) Dockrill		Vic NSW Qld	
<b>Plexaure</b> Endl.			
<i>Plexaure crassiuscula</i> (Nicholls) M.A.Clem. & D.L.Jones		Qld	
<i>Plexaure limenophylax</i> Endl.		AIT	Norfolk Island
<i>Plexaure listeri</i> (Rolfe) M.A.Clem. & D.L.Jones		AIT	Christmas Island
<b>Pomatocalpa</b> Breda, Kuhlew. & Hasselt			
<i>Pomatocalpa macphersonii</i> (F.Muell.) T.E.Hunt		Qld	
<i>Pomatocalpa marsupiale</i> (Kraenzl.) J.J.Sm.		Qld	
<b>Praecoxanthus</b> Hopper & A.P.Br.			
<i>Praecoxanthus aphyllus</i> (Benth.) Hopper & A.P.Br.		WA	
<b>Prasophyllum</b> R.Br.			
<i>Prasophyllum affine</i> Lindl.		NSW	
<i>Prasophyllum album</i> R.S.Rogers		SA	
<i>Prasophyllum alpestre</i> D.L.Jones		Vic NSW ACT	
<i>Prasophyllum alpinum</i> R.Br.		Tas	
<i>Prasophyllum amoenum</i> D.L.Jones		Tas	
<i>Prasophyllum anticum</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum apoxychilum</i> D.L.Jones		Tas	
<i>Prasophyllum appendiculatum</i> Nicholls		Vic NSW	
<i>Prasophyllum asinantum</i> R.J.Bates		SA	
<i>Prasophyllum atratum</i> D.L.Jones		Tas	
<i>Prasophyllum australe</i> R.Br.		SA Vic Tas NSW Qld	
<i>Prasophyllum bagoense</i> D.L.Jones		NSW	
<i>Prasophyllum barnettii</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum beatrix</i> D.L.Jones & D.T.Rouse		Vic NSW	
<i>Prasophyllum brevilabre</i> (Lindl.) Hook.f.		Vic Tas NSW ACT Qld	
<i>Prasophyllum brownii</i> Rchb.f.		WA	
<i>Prasophyllum calcicola</i> R.J.Bates		WA SA	
<i>Prasophyllum campestre</i> R.J.Bates & D.L.Jones		NSW Qld	
<i>Prasophyllum canaliculatum</i> D.L.Jones		NSW ACT	
<i>Prasophyllum candidum</i> R.J.Bates & D.L.Jones		NSW	
<i>Prasophyllum caricetum</i> D.L.Jones		NSW	

Australian Species	Alternate Genus	Distribution	Notes
<i>Prasophyllum castaneum</i> D.L.Jones		Tas	
<i>Prasophyllum catenemum</i> D.L.Jones		SA	
<i>Prasophyllum caudiculum</i> D.L.Jones		NSW	
<i>Prasophyllum chasmogamum</i> R.J.Bates & D.L.Jones		Vic	
<i>Prasophyllum colemaniae</i> R.S.Rogers		Vic	probably extinct
<i>Prasophyllum collinum</i> D.L.Jones		SA	
<i>Prasophyllum concinnum</i> Nicholls		Tas	
<i>Prasophyllum constrictum</i> R.S.Rogers		SA	
<i>Prasophyllum correctum</i> D.L.Jones		Vic	
<i>Prasophyllum crebriflorum</i> D.L.Jones		Tas	
<i>Prasophyllum cucullatum</i> Rchb.f.	<i>Chiloterus</i>	WA	
<i>Prasophyllum cyphochilum</i> Benth.		WA	
<i>Prasophyllum diversiflorum</i> Nicholls		Vic	
<i>Prasophyllum dossenium</i> R.J.Bates & D.L.Jones		NSW	
<i>Prasophyllum drummondii</i> Rchb.f.		WA	
<i>Prasophyllum elatum</i> R.Br.		WA SA Vic Tas NSW Qld	
<i>Prasophyllum erythrocommum</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum exilis</i> D.L.Jones & R.J.Bates		NSW Qld	
<i>Prasophyllum favonium</i> D.L.Jones		Tas	
<i>Prasophyllum fecundum</i> R.J.Bates		SA	
<i>Prasophyllum fimbria</i> Rchb.f.		WA	
<i>Prasophyllum fitzgeraldii</i> R.S.Rogers & Maiden		SA	
<i>Prasophyllum flavum</i> R.Br.		Vic Tas NSW Qld	
<i>Prasophyllum fosteri</i> D.L.Jones		Vic	
<i>Prasophyllum frenchii</i> F.Muell.		SA Vic	
<i>Prasophyllum fuscum</i> R.Br.		NSW	
<i>Prasophyllum gibbosum</i> R.Br.	<i>Chiloterus</i>	WA	
<i>Prasophyllum giganteum</i> Lindl.		WA	
<i>Prasophyllum gilgai</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum goldsackii</i> J.Z. Weber & R.J.Bates		SA	
<i>Prasophyllum gracile</i> Lindl.		WA	
<i>Prasophyllum helophilum</i> D.L.Jones & D.T.Rouse		NSW	
<i>Prasophyllum hians</i> Rchb.f.		WA	
<i>Prasophyllum hygrophilum</i> D.L.Jones & D.Rouse		Vic	
<i>Prasophyllum incompositum</i> D.L.Jones		Qld	
<i>Prasophyllum incorrectum</i> D.L.Jones		Tas	
<i>Prasophyllum incurvum</i> D.L.Jones		Tas	
<i>Prasophyllum innubum</i> D.L.Jones		NSW ACT	
<i>Prasophyllum keltonii</i> D.L.Jones		NSW	
<i>Prasophyllum laxum</i> R.J.Bates		SA	
<i>Prasophyllum limnetes</i> D.L.Jones		Tas	
<i>Prasophyllum lindleyanum</i> Rchb.f.		Vic Tas	
<i>Prasophyllum litorale</i> R.J.Bates		SA Vic	
<i>Prasophyllum maccannii</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum macrostachyum</i> R.Br.		WA	
<i>Prasophyllum macrotys</i> Lindl.		WA	
<i>Prasophyllum milfordense</i> D.L.Jones		Tas	
<i>Prasophyllum mimulum</i> D.L.Jones		Tas	
<i>Prasophyllum montanum</i> R.J.Bates & D.L.Jones		NSW ACT	also recorded for Tasmania but these records represent an undescribed species <i>Prasophyllum</i> species 'Arthurs Lake'
<i>Prasophyllum morganii</i> Nicholls		Vic	probably extinct
<i>Prasophyllum murfetii</i> D.L.Jones		SA	
<i>Prasophyllum niphopedium</i> D.L.Jones		Vic	
<i>Prasophyllum occidentale</i> R.S.Rogers		SA Vic	
<i>Prasophyllum occultans</i> R.J.Bates		SA Vic	
<i>Prasophyllum odoratissimum</i> D.L.Jones		WA	
<i>Prasophyllum odoratum</i> R.S.Rogers		SA Vic	

Australian Species	Alternate Genus	Distribution	Notes
<i>Prasophyllum olidum</i> D.L.Jones		Tas	
<i>Prasophyllum ovale</i> Lindl.		WA	
<i>Prasophyllum pallens</i> D.L.Jones		NSW	
<i>Prasophyllum pallidum</i> Nicholls		SA	
<i>Prasophyllum parviflorum</i> Nicholls		Vic	
<i>Prasophyllum parvifolium</i> Lindl.	<i>Mecopodum</i>	WA	
<i>Prasophyllum patens</i> R.Br.		Vic NSW	
<i>Prasophyllum paulineae</i> D.L.Jones & M.A.Clem.		WA	
<i>Prasophyllum perangustum</i> D.L.Jones		Tas	
<i>Prasophyllum petilum</i> D.L.Jones & R.J.Bates		NSW ACT	
<i>Prasophyllum plumiforme</i> Fitzg.		WA	
<i>Prasophyllum praecox</i> D.L.Jones		SA	
<i>Prasophyllum pruinatum</i> R.S.Rogers		SA	
<i>Prasophyllum pulchellum</i> D.L.Jones		Tas	
<i>Prasophyllum pyriforme</i> E.Coleman		Vic	
<i>Prasophyllum readii</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum regium</i> R.S.Rogers		WA	
<i>Prasophyllum retroflexum</i> D.L.Jones		Vic NSW	
<i>Prasophyllum robustum</i> (Nicholls) M.A.Clem. & D.L.Jones		Tas	
<i>Prasophyllum rogersii</i> Rupp		NSW	
<i>Prasophyllum rostratum</i> Lindl.		Tas	
<i>Prasophyllum rotundiflorum</i> R.S.Rogers		SA	
<i>Prasophyllum sargentii</i> (Nicholls) A.S.George		WA	
<i>Prasophyllum secutum</i> D.L.Jones		Tas	
<i>Prasophyllum solstitium</i> D.L.Jones		NSW	
<i>Prasophyllum sphacelatum</i> D.L.Jones		Vic Tas NSW ACT	
<i>Prasophyllum spicatum</i> R.J.Bates & D.L.Jones		SA Vic	
<i>Prasophyllum stellatum</i> D.L.Jones		Tas	
<i>Prasophyllum striatum</i> R.Br.	<i>Mecopodum</i>	NSW	
<i>Prasophyllum suaveolens</i> D.L.Jones & R.J.Bates		Vic	
<i>Prasophyllum subbisectum</i> Nicholls		Vic	
<i>Prasophyllum suttonii</i> R.S.Rogers & B.Rees		Vic	probably extinct
<i>Prasophyllum sylvestre</i> R.J.Bates & D.L.Jones		Vic NSW	
<i>Prasophyllum tadgellianum</i> R.S.Rogers		Vic Tas NSW ACT	
<i>Prasophyllum taphanyx</i> D.L.Jones		Tas	
<i>Prasophyllum triangulare</i> Fitzg.		WA	
<i>Prasophyllum truncatum</i> Lindl.		Tas	
<i>Prasophyllum tunbridgense</i> D.L.Jones		Tas	
<i>Prasophyllum uvidulum</i> D.L.Jones & D.T.Rouse		Vic	
<i>Prasophyllum validum</i> R.S.Rogers		SA	
<i>Prasophyllum venustum</i> D.L.Jones & D.T.Rouse		NSW ACT	
<i>Prasophyllum vesuvium</i> R.J.Bates		SA	
<i>Prasophyllum viretrum</i> D.L.Jones & D.T.Rouse		SA Vic	
<i>Prasophyllum viriosum</i> D.L.Jones & D.T.Rouse		NSW ACT	
<i>Prasophyllum wallum</i> R.J.Bates & D.L.Jones		Qld	
<i>Prasophyllum wilkinsoniorum</i> D.L.Jones		NSW	
<i>Prasophyllum</i> species 'ACT 1'		ACT	late spring-summer flowering, scented aff. <i>patens</i> with extensively white petals and strongly recurved labellum with short green callus plate; montane forest between 700 and 900 m (Jones <i>et al.</i> 2008)
<i>Prasophyllum</i> species 'ACT 2'		ACT	late spring-summer flowering, scented aff. <i>patens</i> with white-margined brownish petals and more upright labellum with short green callus plate; montane forest between 700 and 900 m (Jones <i>et al.</i> 2008)
<i>Prasophyllum</i> species 'Alligator Gorge'		SA	aff. <i>fitzgeraldii</i> with slender flowers with a pale pink labellum; woodland in southern Flinders Ranges; probably extinct (Bates 2011)
<i>Prasophyllum</i> species 'Anglesea'		Vic	late spring flowering aff. <i>odoratum</i> with fewer, smaller flowers with a shorter labellum; heathy woodland at Anglesea (Jeanes & Backhouse 2006; = aff. <i>odoratum</i> E Rouse 2002)

Australian Species	Alternate Genus	Distribution	Notes
<i>Prasophyllum</i> species 'Arthurs Lake'		Tas	aff. <i>montanum</i> ; grows in the ecotone between wet grassland and woodland on rocky ground above 700 m information available; previously called <i>P. montanum</i> in Tasmania (Jones <i>et al.</i> 1999)
<i>Prasophyllum</i> species 'Barrington Tops'		NSW	aff. <i>fuscum</i> from Barrington Tops; known only from a single collection, no other information available
<i>Prasophyllum</i> species 'basalt plains'		Vic	spring-flowering, robust, clump-forming aff. <i>suaveolens</i> with larger flowers; western basalt plains grasslands (Jeanes & Backhouse 2006; = aff. <i>occidentale</i> E Rouse 2002)
<i>Prasophyllum</i> species 'black soil plains'		NSW	early spring-flowering aff. <i>odoratum</i> with large flowers with relatively short lateral sepals; black soil plains near Inverell (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'Boorhaman'		Vic	spring-flowering aff. <i>petilum</i> with very narrow labellum and narrow petals; native grasslands near Boorhaman (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Boyd Plateau'		NSW	tall aff. <i>alpestre</i> with narrow flowers with smaller, less ruffled labellum; wet heaths on Boyd Plateau and Mt Werong (Bishop 2000)
<i>Prasophyllum</i> species 'bright shiny maroon'		SA	aff. <i>fitzgeraldii</i> with bright shiny maroon flowers; woodland and mallee scrub on sandy soils between Keith and Loxton; <i>leiochilum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'Brookton Highway'	<i>Mecopodum</i>	WA	late winter-early spring flowering aff. <i>parvifolium</i> with few small poorly opening self-pollinating flowers; swamps and moist margins of granite outcrops from Aramadale to Brookton (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'brown lip'		SA Vic	aff. <i>fitzgeraldii</i> with brown lip with narrow white margin; heathy woodland in lower south-east of SA and far western Vic (Bates 2011; Jeanes & Backhouse 2006); <i>kafechilum</i> ms
<i>Prasophyllum</i> species 'Bunyip'		Vic	tall, fairly slender aff. <i>odoratum</i> with widely spaced flowers with narrowly spreading to incurved petals; open forest in west & central Gippsland, seen mostly after fire (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'bushfires'		SA	slender aff. <i>constrictum</i> with variably coloured, strongly scented flowers with a long, slender, constricted labellum; woodland on rocky ridges in the Mount Lofty Ranges, flowering only after fire (Bates 2011); <i>pyrophilum</i> ms
<i>Prasophyllum</i> species 'coast sandhills'		SA	coastal clumping aff. <i>odoratum</i> with upright labellum and thick-textured flowers; widely distributed in coastal heaths and grasslands (Bates 2011); <i>carnosum</i> ms
<i>Prasophyllum</i> species 'crowded'		WA	aff. <i>elatum</i> with crowded spike of poorly opening, self-pollinating flowers; woodland on sandy rises above seasonally wet flats between Perth and Ravensthorpe, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'Dandenong Ranges'		Vic	aff. <i>odoratum</i> with tall spike of well-spaced recurved flowers; open forest in the Dandenong Ranges (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> G Rouse 2002)
<i>Prasophyllum</i> species 'Deep Lead'		Vic	aff. <i>fitzgeraldii</i> with short-lived flowers with brown labellum lamina; heathy woodland on clay soils at Deep Lead (Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> B Rouse 2002)
<i>Prasophyllum</i> species 'Delegate'		NSW	late spring/early summer flowering, small aff. <i>petilum</i> ; grasslands on the Monaro Tablelands near Delegate
<i>Prasophyllum</i> species 'desert'		SA NSW	robust aff. <i>odoratum</i> with long narrow petals; semi-arid sandy and rocky grassy shrublands, often with <i>Triodia</i> ; widespread across central SA and far western NSW (Bates 2011); <i>eremaemum</i> ms
<i>Prasophyllum</i> species 'Digby'		Vic	aff. <i>fitzgeraldii</i> with green labellum lamina and pink callus; dense heathy woodland near Digby, flowering after fire (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'early'	<i>Mecopodum</i>	WA	autumn flowering aff. <i>parvifolium</i> with scented green and white flowers lacking red stripes; heath, woodland, forest and granite outcrops between Bunbury and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'eastern highlands'		Vic NSW	summer flowering aff. <i>odoratum</i> with tall, moderately crowded spike of large flowers; subalpine woodland of eastern Vic and southern NSW above 800 m (=aff. <i>odoratum</i> L Rouse 2002)
<i>Prasophyllum</i> species 'eastern wheatbelt'		WA	aff. <i>elatum</i> with smaller, dark flowers with a less ruffled labellum margin; flowers after fire and on lightly disturbed ground (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'enigma'		SA	aff. <i>fitzgeraldii</i> with partially opening, often contorted flowers with a green labellum callus and purple lamina; hilly woodland from central Eyre Peninsula to the southern Mt Lofty Ranges (Bates 2011); <i>aenigmum</i> ms
<i>Prasophyllum</i> species 'foothills'		Vic	small, early spring-flowering aff. <i>patens</i> with slender spike of widely spaced flowers with long, narrow pointed labellum; foothill open forest from Moondarra (Vic) to Yass (NSW) (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Gilbert River'		SA	aff. <i>frenchii</i> with larger green and maroon flowers; moist grasslands on heavy clay soils in the Gilbert River valley in the northern Mt Lofty Ranges; last seen in 1990 and almost certainly extinct (Bates 2011)

Australian Species	Alternate Genus	Distribution	Notes
<i>Prasophyllum</i> species 'grassy woodlands'		Vic	spring-flowering aff. <i>diversiflorum</i> with shorter spike of smaller, more crowded flowers; damp sites in grassy woodland in the north-east between Nagambie and Chesney (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Inglewood'		Vic	small aff. <i>pyriforme</i> with a raised, broadly and deeply grooved labellum callus; dry woodland at Inglewood & Arnold West (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'inland woodlands'		Vic	aff. <i>validum</i> with large, relatively broad dull brownish-green flowers; widespread across drier inland woodlands Chiltern to Stawell (Jeanes & Backhouse 2006; =aff. <i>pyriforme</i> 5 Rouse 2002)
<i>Prasophyllum</i> species 'Innes'		SA	aff. <i>occidentale</i> with larger, more numerous flowers with a broad pinkish labellum lamina; coastal sedges and short open scrub on the tip of the Yorke Peninsula; <i>ventosum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'Inverleigh'		Vic	aff. <i>validum</i> with dull green flowers with short, broad column arms; sparsely shrubby woodland on fertile sandy loam soils at Inverleigh (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Jip Jip'		SA	aff. <i>fitzgeraldii</i> with greenish white labellum; dry woodland on sandy soils between Keith and Padthaway, almost certainly extinct (Bates 2011)
<i>Prasophyllum</i> species 'Lara'		Vic	aff. <i>odoratum</i> with long narrow green callus plate; basalt plains grasslands near Lara; presumed extinct (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> M Rouse)
<i>Prasophyllum</i> species 'late coastal dunes'		SA	robust, late flowering aff. <i>odoratum</i> with large fleshy flowers and senescent leaf; coastal scrubs on calcareous sand dunes and flats (Bates 2011)
<i>Prasophyllum</i> species 'late summer'		NSW	late summer-flowering aff. <i>dossenum</i> with a prominent kink in the lateral sepals; montane grassland and grassy woodland on the Barrington Tops (L. Copeland unpublished)
<i>Prasophyllum</i> species 'late swamps'		WA	late flowering aff. <i>drummondii</i> with smaller, darker, more widely spaced flowers; on seasonally wet flats between Augusta and Israelite Bay, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'limestone heath'		Vic	small aff. <i>parviflorum</i> with smaller, paler flowers; coastal heath on limestone between Portland and Nelson (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'lower south-east'		SA	aff. <i>fitzgeraldii</i> with fleshy bright green flowers with a green labellum callus and off white lamina; winter wet heath on silty clays in the lower south-east, flowering after fire; <i>crassum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'Mallacoota'		Vic	robust aff. <i>spicatum</i> with smaller, less crowded, fragrant flowers with widely spreading petals; coastal heaths, heathy woodland in East Gippsland, mostly seen after fire (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> K Rouse 2002)
<i>Prasophyllum</i> species 'mallee sands'		SA Vic	aff. <i>odoratum</i> with narrow overlapping green and white flowers; mallee heath on sand dunes in the Big Desert (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> B Rouse)
<i>Prasophyllum</i> species 'Midlands'		Vic	tall aff. <i>odoratum</i> with widely spaced, somewhat narrow flowers; woodland on rocky hills from Chewton to St Arnaud (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Mokota'		SA	short, slender aff. <i>constrictum</i> with small green flowers; montane grassland on bare hills above 500 m in the Mt Lofy Ranges; <i>arvense</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'montane'		Vic	aff. <i>montanum</i> with narrower labellum callus plate, broader lamina and longer, narrower sepals; foothills and mountains of eastern Victoria from Alexandra to Wulgulmerang (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Mortlake'		Vic	tall, robust aff. <i>correctum</i> with larger, more numerous and more crowded flowers; basalt plains grassland at Mortlake (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Mount Kaputar'		NSW	aff. <i>odoratum</i> with green and white flowers with long, thin petals and lateral sepals; open forest in Nandawar Range (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'mountain bogs'		Vic NSW	relatively tall, late summer-flowering aff. <i>alpestre</i> with narrow petals and sharply curved labellum; mountain swamps above 700 m, Tamboritha (Vic) to Adaminaby (NSW) (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Murchison'		Vic	aff. <i>petilum</i> with narrower, more brownish to greenish labellum; native grasslands near Murchison (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'needle points'		SA	tall, robust aff. <i>occidentale</i> with pinkish toned flowers with long fine points to the sepals and reddish column wings; dry woodland under she-oaks <i>Allocasuarina</i> spp. on the eastern Eyre Peninsula; <i>aciculare</i> ms (Bates 2011)
<i>Prasophyllum</i> species 'New England batholith'		NSW Qld	late spring-summer flowering aff. <i>odoratum</i> with crowded flowers with large, deeply ruffled labellum; open forest in the granite ranges above 900 m on New England Tableland NSW (L. Copeland unpubl.) & near Stanthorpe Qld



Australian Species	Alternate Genus	Distribution	Notes
<i>Prasophyllum</i> species 'Noorinbee'		Vic	slender early spring-flowering aff. <i>odoratum</i> with few widely spaced flowers; shrubby open forest in East Gippsland, seen mostly after fires (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> I Rouse 2002)
<i>Prasophyllum</i> species 'northern montane'		NSW	small, summer flowering aff. <i>odoratum</i> with widely spaced brown and white flowers; moist grassy woodland and open forest above 1,000, Barrington Tops, Ebor and Dorrigo (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'Nowra'		NSW	small aff. <i>appendiculatum</i> with more brownish flowers with shorter, broader petals and sepals and a broader, flatter labellum; heathy woodland on the Shoalhaven escarpment (Stephenson 2011)
<i>Prasophyllum</i> species 'Nunniong'		Vic	larger, mid-late summer flowering aff. <i>sphacelatum</i> with broader lip and growing in damper sites; alpine plains and woodland in the Nunniong and Mt Cobberas regions (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Old Bar'		NSW	aff. <i>fuscum</i> with greenish brown flowers with white labellum lamina and short broad lateral sepals; coastal heath at Old Bar (Copeland unpubl.)
<i>Prasophyllum</i> species 'Otway coast'		Vic	aff. <i>odoratum</i> with short, crowded spike; coastal heath from Anglesea to Port Campbell (=aff. <i>odoratum</i> E/F Rouse 2002)
<i>Prasophyllum</i> species 'Otway Ranges'		Vic	tall, slender aff. <i>odoratum</i> with small, widely spaced flowers; shrubby open forest in the Otway Ranges, seen mostly after fire (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> H Rouse 2002)
<i>Prasophyllum</i> species 'Piccaninnie Ponds'		SA	aff. <i>parviflorum</i> with short broad petals and sepals and a broad deeply grooved labellum; swampy sites in far south-east (Bates 2011)
<i>Prasophyllum</i> species 'Pilliga Scrub'		NSW	robust aff. <i>odoratum</i> with pale green, white and brown flowers with long, narrow, spreading lateral sepals, narrow petals and long labellum with short callus plate and broad lamina; dry woodland in Pilliga Scrub (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'pink lip'		SA Vic	aff. <i>fitzgeraldii</i> with pink labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> A Rouse); <i>rosaechilum</i> ms
<i>Prasophyllum</i> species 'purple lip'		SA Vic	aff. <i>fitzgeraldii</i> with purple labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> E Rouse); <i>violescens</i> ms
<i>Prasophyllum</i> species 'Riverina grasslands'		Vic NSW	tall, early flowering aff. <i>suaveolens</i> with larger flowers with a long narrowly triangular labellum; Riverina plains grasslands from Hunter (Vic) to Moama (NSW) (Jeanes & Backhouse 2006; =aff. <i>occidentale</i> D Rouse 2002)
<i>Prasophyllum</i> species 'sedge swamps'		SA Vic	short aff. <i>odoratum</i> with few widely spaced flowers; heath and sedge swamps, often flowering in water (= sp. 'Drajurk' Backhouse <i>et al.</i> 2016; sp. 'Topperwein' Bates 2011)
<i>Prasophyllum</i> species 'Smokers Gap'		Vic ACT	large, late-summer flowering aff. <i>sphacelatum</i> ; alpine meadows in Namadgi National Park; mentioned in notes on <i>P. spacelatum</i> in Jones <i>et al.</i> (2008); possibly the same as <i>P. species</i> 'Nunniong'?
<i>Prasophyllum</i> species 'southern'		WA	short, late spring-early summer flowering aff. <i>fimbria</i> with smaller darker flowers; seasonally wet flats between Augusta and Israelite Bay, flowering after summer fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'striped'	<i>Chiloterus</i>	WA	aff. <i>gibbosum</i> with green and white flowers with humped base to the lateral sepals; seasonally wet flats between Perth and Busselton, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'Swan region'		WA	aff. <i>elatum</i> with smaller paler flowers with a less ruffled labellum margin; woodland in rocky hills of the Darling Scarp between Greenmount and Armadale, flowering only after summer fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'Teralba'		NSW	aff. <i>exilis</i> with ovary held away from the stem; coastal <i>Angophora</i> forest near Teralba (NSW Herbarium information)
<i>Prasophyllum</i> species 'Verran Tanks'		SA	aff. <i>validum</i> with fleshy pink to brown-toned flowers; dry scrubs on sand plains and rocky hills on the eastern Eyre Peninsula between Cleve and Cowell (Bates 2011); <i>eyreanum</i> ms
<i>Prasophyllum</i> species 'waterholes'		SA Vic	very late flowering aff. <i>odoratum</i> with very widely spreading lateral sepals and incurved petals; heath and sedge swamps in south-western Vic & south-eastern SA (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> C/D Rouse 2002)
<i>Prasophyllum</i> species 'Wilson's Promontory'		Vic	aff. <i>frenchii</i> with smaller, duller flowers with more ruffled labellum and lower callus; moist coastal heath at Wilson's Promontory (= sp. <i>frenchii</i> 2 Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Wimmera woodlands'		SA Vic	short aff. <i>occidentale</i> with smaller, more widely spaced flowers with a narrower labellum; dry woodlands in narrow band between Murtoa (Vic) and Keith (SA)
<b><i>Pseudovanilla</i> Garay</b>			
<i>Pseudovanilla foliata</i> (F. Muell.) Garay		NSW Qld	

Australian Species	Alternate Genus	Distribution	Notes
<b><i>Pterostylis</i> R.Br.</b>			
<i>Pterostylis abrupta</i> D.L.Jones	<i>Diplodium</i>	NSW	
<i>Pterostylis aciculiformis</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	Vic NSW ACT	
<i>Pterostylis acuminata</i> R.Br.		Vic NSW Qld	
<i>Pterostylis aestiva</i> D.L.Jones	<i>Diplodium</i>	Vic NSW ACT	
<i>Pterostylis agrestis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic	
<i>Pterostylis alata</i> (Labill.) Rchb.f.	<i>Diplodium</i>	Tas	
<i>Pterostylis allantoidea</i> R.S.Rogers	<i>Eremorchis</i>	WA	
<i>Pterostylis alpina</i> R.S.Rogers		Vic NSW ACT	
<i>Pterostylis alveata</i> Garnet	<i>Diplodium</i>	Vic NSW	
<i>Pterostylis amabilis</i> (D.L.Jones & L.M.Copel.) D.L.Jones	<i>Speculantha</i>	NSW	
<i>Pterostylis anaclasta</i> (D.L.Jones) Janes & Duretto		Qld	
<i>Pterostylis anatona</i> D.L.Jones		Qld	
<i>Pterostylis aneba</i> D.L.Jones		Vic NSW ACT	Victorian records in doubt
<i>Pterostylis angusta</i> A.S.George	<i>Diplodium</i>	WA	
<i>Pterostylis antennifera</i> D.L.Jones	<i>Speculantha</i>	NSW Qld	
<i>Pterostylis aphylla</i> Lindl.	<i>Speculantha</i>	Tas	
<i>Pterostylis aquilonia</i> D.L.Jones & B.Gray	<i>Diplodium</i>	Qld	
<i>Pterostylis arenicola</i> M.A.Clem. & J.Stewart	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis aspera</i> D.L.Jones & M.A.Clem.	<i>Diplodium</i>	WA	
<i>Pterostylis atrans</i> D.L.Jones	<i>Diplodium</i>	Vic Tas NSW ACT	
<i>Pterostylis atriola</i> D.L.Jones	<i>Speculantha</i>	Tas	
<i>Pterostylis baptistii</i> Fitzg.		Vic NSW Qld	
<i>Pterostylis barbata</i> Lindl.	<i>Plumatichilos</i>	WA	
<i>Pterostylis barringtonensis</i> D.L.Jones	<i>Bunochilus</i>	NSW	
<i>Pterostylis basaltica</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	Vic	
<i>Pterostylis bicolor</i> M.A.Clem. & D.L.Jones	<i>Hymenochilus</i>	Vic NSW ACT Qld	
<i>Pterostylis bicornis</i> D.L.Jones & M.A.Clem.	<i>Petrorchis</i>	NSW Qld	
<i>Pterostylis biseta</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis boormanii</i> Rupp	<i>Oligochaetochilus</i>	SA Vic NSW	
<i>Pterostylis brevichila</i> D.L.Jones & C.J.French	<i>Diplodium</i>	WA	
<i>Pterostylis brunneola</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis bryophila</i> D.L.Jones	<i>Diplodium</i>	SA	
<i>Pterostylis calceolus</i> M.A.Clem.	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis caligna</i> M.T.Mathieson		Qld	
<i>Pterostylis chaetophora</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis cheraphila</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	Vic	
<i>Pterostylis chlorogramma</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	SA Vic	
<i>Pterostylis chocolatina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis ciliata</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis clavigera</i> Fitzg.	<i>Linguella</i>	NSW	doubtfully distinct from <i>Pterostylis nana</i>
<i>Pterostylis clivicola</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic NSW ACT	
<i>Pterostylis clivosa</i> (D.L.Jones) D.L.Jones	<i>Speculantha</i>	Vic	
<i>Pterostylis cobarensis</i> M.A.Clem.	<i>Oligochaetochilus</i>	SA NSW Qld	
<i>Pterostylis coccinea</i> Fitzg.	<i>Diplodium</i>	Vic NSW ACT	
<i>Pterostylis collina</i> (Rupp) M.A.Clem. & D.L.Jones	<i>Taurantha</i>	NSW	
<i>Pterostylis commutata</i> D.L.Jones	<i>Oligochaetochilus</i>	Tas	
<i>Pterostylis concava</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	WA	
<i>Pterostylis concinna</i> R.Br.	<i>Taurantha</i>	SA Vic Tas NSW	
<i>Pterostylis conferta</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic	
<i>Pterostylis crassa</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	Vic	
<i>Pterostylis crassicaulis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic NSW ACT	
<i>Pterostylis crassichila</i> D.L.Jones	<i>Bunochilus</i>	Qld	
<i>Pterostylis crebra</i> D.L.Jones	<i>Speculantha</i>	NSW	

Australian Species	Alternate Genus	Distribution	Notes
( <i>Pterostylis crebriflorus</i> D.L.Jones & C.J.French)	<i>Urochilus</i>	WA	in describing <i>crebriflorus</i> , Jones & French have taken a narrower interpretation than that of Brown <i>et al.</i> 2013 for <i>P.</i> species 'crowded'; described in <i>Urochilus</i> , new combination in <i>Pterostylis</i> not yet made
<i>Pterostylis cucullata</i> R.Br. subspecies <i>cucullata</i>		SA Vic Tas	extinct in South Australia (Bates 2011)
<i>Pterostylis cucullata</i> R.Br. subspecies <i>sylvicola</i> D.L.Jones		SA Vic	
<i>Pterostylis curta</i> R.Br.		SA Vic Tas NSW ACT Qld	also on Lord Howe Island
<i>Pterostylis cynnocephala</i> Fitzg.	<i>Hymenochilus</i>	SA Vic NSW ACT Qld	
<i>Pterostylis daintreeana</i> Benth.	<i>Pharochilum</i>	NSW Qld	also spelt <i>daintreana</i>
<i>Pterostylis decurva</i> R.S.Rogers	<i>Diplodium</i>	Vic Tas NSW ACT	
<i>Pterostylis depauperata</i> F.M.Bailey	<i>Crangonorchis</i>	Qld	
<i>Pterostylis despectans</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	SA Vic	
<i>Pterostylis dilatata</i> A.S.George	<i>Linguella</i>	WA	
<i>Pterostylis diminuta</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	Vic	
<i>Pterostylis dolichochila</i> D.L.Jones & M.A.Clem.	<i>Diplodium</i>	SA Vic	
<i>Pterostylis dubia</i> R.Br.		Tas	
<i>Pterostylis echinulata</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis elegans</i> D.L.Jones	<i>Diplodium</i>	NSW	
<i>Pterostylis elegantissima</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis erecta</i> T.E.Hunt		NSW Qld	
<i>Pterostylis erubescens</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis erythroconcha</i> M.A.Clem. & D.L.Jones	<i>Diplodium</i>	SA	
<i>Pterostylis exalla</i> D.L.Jones	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis excelsa</i> M.A.Clem.	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis exquisita</i> D.L.Jones	<i>Speculantha</i>	Qld	
<i>Pterostylis exserta</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis extranea</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	Qld	
<i>Pterostylis falcata</i> R.S.Rogers		SA Vic Tas NSW ACT	
<i>Pterostylis ferruginea</i> (D.L.Jones) G.N.Backh.	<i>Oligochaetochilus</i>	SA Vic	
<i>Pterostylis fischii</i> Nicholls	<i>Diplodium</i>	Vic NSW ACT Qld	
<i>Pterostylis flavovirens</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	SA	
<i>Pterostylis foliata</i> Hook.f.		SA Vic Tas NSW	
<i>Pterostylis frenchii</i> (D.L.Jones) A.P.Br.	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis furcata</i> Lindl.		Tas	
<i>Pterostylis furva</i> D.L.Jones	<i>Speculantha</i>	NSW	
<i>Pterostylis gibbosa</i> R.Br.	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis glebosa</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis glyphida</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	NSW	
( <i>Pterostylis gracilens</i> (D.L.Jones)	<i>Plumatichilos</i>	SA	described in <i>Plumatichilos</i> , new combination in <i>Pterostylis</i> not yet made
<i>Pterostylis grandiflora</i> R.Br.	<i>Diplodium</i>	Vic Tas NSW Qld	
<i>Pterostylis hadra</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis hamata</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	Vic NSW ACT	
<i>Pterostylis hamiltonii</i> Nicholls	<i>Diplodium</i>	WA	
<i>Pterostylis hians</i> D.L.Jones	<i>Diplodium</i>	NSW	
<i>Pterostylis hildae</i> Nicholls		NSW Qld	
<i>Pterostylis hispidula</i> Fitzg.		NSW Qld	
<i>Pterostylis incognita</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic	probably extinct
<i>Pterostylis insectifera</i> M.A.Clem.	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis jacksonii</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis jonesii</i> G.N.Backh.	<i>Bunochilus</i>	Vic NSW ACT	originally described as <i>Bunochilus montanus</i> D.L. Jones
<i>Pterostylis karri</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis laxa</i> Blackmore	<i>Diplodium</i>	Vic NSW ACT	
<i>Pterostylis lepida</i> D.L.Jones	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis leptochila</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis lineata</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis lingua</i> M.A.Clem.	<i>Oligochaetochilus</i>	SA Vic NSW	there is some doubt as to the correct identity of Victorian and South Australian records of this species
<i>Pterostylis littorale</i> D.L.Jones	<i>Plumatichilos</i>	NSW	
<i>Pterostylis littoralis</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	SA	
<i>Pterostylis loganii</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	Vic NSW	
<i>Pterostylis longicurva</i> Rupp	<i>Diplodium</i>	NSW Qld	
<i>Pterostylis longifolia</i> R.Br.	<i>Bunochilus</i>	NSW Qld	
<i>Pterostylis longipetala</i> Rupp	<i>Diplodium</i>	Vic NSW	
<i>Pterostylis lortensis</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis lustra</i> D.L.Jones		SA Vic Tas	
<i>Pterostylis macilenta</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	Vic	
<i>Pterostylis macrocalymma</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis macrosceles</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis macrosepala</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis major</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis maxima</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	SA Vic NSW	
<i>Pterostylis melagramma</i> D.L.Jones	<i>Bunochilus</i>	SA Vic Tas NSW	
<i>Pterostylis metcalfei</i> D.L.Jones	<i>Diplodium</i>	NSW	
<i>Pterostylis microglossa</i> D.L.Jones & C.J.French	<i>Diplodium</i>	WA	
<i>Pterostylis microphylla</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis mirabilis</i> D.L.Jones	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis mitchellii</i> Lindl.	<i>Oligochaetochilus</i>	NSW Qld	
<i>Pterostylis monticola</i> D.L.Jones		Vic NSW ACT	
<i>Pterostylis multiflora</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	Vic NSW ACT	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	WA SA Vic Tas NSW ACT Qld	
<i>Pterostylis mystacina</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	Qld	
<i>Pterostylis nana</i> R.Br.	<i>Linguella</i>	SA Vic Tas NSW	
<i>Pterostylis nigricans</i> D.L.Jones & M.A.Clem	<i>Speculantha</i>	NSW Qld	
<i>Pterostylis nutans</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Pterostylis oblonga</i> D.L.Jones		NSW	
<i>Pterostylis obtusa</i> R.Br.	<i>Diplodium</i>	NSW Qld	also on Lord Howe Island
<i>Pterostylis ophioglossa</i> R.Br.	<i>Taurantha</i>	NSW Qld	
<i>Pterostylis oreophila</i> Clemesha		Vic NSW ACT	
<i>Pterostylis ovata</i> M.A.Clem.	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis parca</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis parva</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA SA	
<i>Pterostylis parviflora</i> R.Br.	<i>Speculantha</i>	SA Vic Tas NSW Qld	
<i>Pterostylis pearsonii</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	Qld	
<i>Pterostylis pedina</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis pedoglossa</i> Fitzg.	<i>Crangonorchis</i>	Vic Tas NSW Qld	
<i>Pterostylis pedunculata</i> R.Br.		SA Vic Tas NSW ACT Qld	also on Lord Howe Island
<i>Pterostylis perculata</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis petrosa</i> D.L.Jones & M.A.Clem	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis picta</i> M.A.Clem.	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis planulata</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	SA Vic	
<i>Pterostylis platypetala</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis plumosa</i> Cady	<i>Plumatichilos</i>	Vic Tas NSW	
<i>Pterostylis praetermissa</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	NSW Qld	
<i>Pterostylis prasina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	SA Vic	
<i>Pterostylis pratensis</i> D.L.Jones	<i>Hymenochilus</i>	Tas	

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis procera</i> D.L.Jones & M.A.Clem.		Qld	
<i>Pterostylis psammophila</i> D.L.Jones	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis pulchella</i> Messmer	<i>Diplodium</i>	NSW	
<i>Pterostylis pusilla</i> R.S.Rogers	<i>Oligochaetochilus</i>	SA Vic NSW	
<i>Pterostylis pyramidalis</i> Lindl.	<i>Linguella</i>	WA	
<i>Pterostylis recurva</i> Benth.	<i>Stamnorchis</i>	WA	
<i>Pterostylis reflexa</i> R.Br.	<i>Diplodium</i>	NSW ACT	
<i>Pterostylis revoluta</i> R.Br.	<i>Diplodium</i>	NSW	
<i>Pterostylis riparia</i> D.L.Jones		NSW	
<i>Pterostylis robusta</i> R.S.Rogers	<i>Diplodium</i>	SA Vic	
<i>Pterostylis robusta</i> R.S.Rogers subspecies 'gorges'	<i>Diplodium</i>	SA	distinctive tall form of <i>robusta</i> with dark red and white-striped flowers with longer dorsal sepal; woodland in gorges in Flinders & Mt Lofty Ranges (Bates 2011); subspecies <i>calvicola</i> ms
<i>Pterostylis roensis</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis rogersii</i> E.Coleman	<i>Diplodium</i>	WA	
<i>Pterostylis rubenachii</i> D.L.Jones	<i>Hymenochilus</i>	Tas	
<i>Pterostylis rubescens</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	Vic NSW ACT	
<i>Pterostylis rufa</i> R.Br.	<i>Oligochaetochilus</i>	NSW Qld	
<i>Pterostylis russellii</i> T.E.Hunt	<i>Diplodium</i>	NSW Qld	
<i>Pterostylis sanguinea</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	WA SA Vic Tas	
<i>Pterostylis sargentii</i> C.R.P.Andrews	<i>Ranorchis</i>	WA	
<i>Pterostylis saxicola</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	NSW	
<i>Pterostylis scabra</i> Lindl.	<i>Diplodium</i>	WA	
<i>Pterostylis scabrida</i> Lindl.		Tas	
<i>Pterostylis scitula</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis scoliosa</i> D.L.Jones	<i>Diplodium</i>	Qld	
<i>Pterostylis setifera</i> M.A.Clem., Matthias & D.L.Jones	<i>Oligochaetochilus</i>	SA Vic NSW Qld	
<i>Pterostylis simulans</i> R.J.Bates	<i>Oligochaetochilus</i>	SA	
<i>Pterostylis sinuata</i> (D.L.Jones) Janes & Duretto	<i>Hymenochilus</i>	WA	
<i>Pterostylis smaragdina</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	SA Vic NSW	
<i>Pterostylis spathulata</i> M.A.Clem.	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis spissa</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	Vic	
<i>Pterostylis squamata</i> R.Br.	<i>Oligochaetochilus</i>	Vic Tas NSW ACT	
<i>Pterostylis stenochila</i> D.L.Jones	<i>Bunochilus</i>	Tas	
<i>Pterostylis stenosepala</i> (D.L.Jones)	<i>Bunochilus</i>	NSW	
<i>Pterostylis striata</i> Fitzg.	<i>Diplodium</i>	SA Vic NSW	
<i>Pterostylis stricta</i> Clemesha & B.Gray		Qld	
<i>Pterostylis subtilis</i> D.L.Jones		NSW	
<i>Pterostylis tasmanica</i> D.L.Jones	<i>Plumatichilos</i>	SA Vic Tas NSW	
<i>Pterostylis taurus</i> M.A.Clem. & D.L.Jones	<i>Taurantha</i>	Qld	
<i>Pterostylis telmata</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	
<i>Pterostylis tenuis</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis tenuissima</i> Nicholls	<i>Diplodium</i>	SA Vic	
<i>Pterostylis thulia</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	Qld	
<i>Pterostylis timothyi</i> (D.L.Jones) Janes & Duretto	<i>Linguella</i>	WA	
<i>Pterostylis torquata</i> D.L.Jones	<i>Diplodium</i>	NSW Qld	
<i>Pterostylis truncata</i> Fitzg.	<i>Diplodium</i>	Vic NSW ACT	
<i>Pterostylis tunstallii</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	Vic Tas NSW	
<i>Pterostylis turfosa</i> Endl.	<i>Plumatichilos</i>	WA	
<i>Pterostylis uliginosa</i> D.L.Jones	<i>Speculantha</i>	SA Vic Tas NSW	
<i>Pterostylis umbrina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW ACT	
<i>Pterostylis valida</i> (Nicholls) D.L.Jones	<i>Oligochaetochilus</i>	Vic	
<i>Pterostylis ventricosa</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	NSW	
<i>Pterostylis vernalis</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	NSW	
<i>Pterostylis viriosa</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	SA	
<i>Pterostylis vitrea</i> (D.L.Jones) Bostock	<i>Bunochilus</i>	Qld	
<i>Pterostylis vittata</i> Lindl.	<i>Urochilus</i>	WA	
<i>Pterostylis voigtii</i> D.L.Jones & C.J.French	<i>Linguella</i>	WA	



Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis wapstrarum</i> D.L.Jones	<i>Hymenochilus</i>	Tas	also spelt <i>wapstreorum</i>
<i>Pterostylis williamsonii</i> D.L.Jones	<i>Bunochilus</i>	Tas	
<i>Pterostylis woollsii</i> Fitzg.	<i>Oligochaetochilus</i>	Vic NSW Qld	
<i>Pterostylis xerophila</i> M.A.Clem.	<i>Oligochaetochilus</i>	SA Vic	
<i>Pterostylis zebrina</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	WA	
<i>Pterostylis ziegeleri</i> D.L.Jones	<i>Hymenochilus</i>	Tas	
<i>Pterostylis</i> species 'Abercrombie Caves'	<i>Oligochaetochilus</i>	NSW	large aff. <i>aciculiformis</i> with crowded flowers on short pedicels; grassy woodland in Abercrombie Caves region on southern tablelands (Bishop 2000)
<i>Pterostylis</i> species 'ACT'	<i>Bunochilus</i>	ACT	small, early flowering aff. <i>melagramma</i> with brown sepal tips to the sepals; shrubby open forest in Namadgi National Park (J. Egan & T. Wood pers. comm.)
<i>Pterostylis</i> species 'Adelaide hills'	<i>Diplodium</i>	SA	aff. <i>alata</i> with smaller, narrower flowers; heathy and shrubby woodland in the Mt Lofty Range and Kangaroo Island (Bates 2011)
<i>Pterostylis</i> species 'Albury'	<i>Bunochilus</i>	NSW	aff. <i>tunstallii</i> with strongly striped flowers; woodland on rocky hills near Albury (Jones & Clements 2006)
<i>Pterostylis</i> species 'Anglesea'	<i>Plumatichilos</i>	Vic	large aff. <i>plumosa</i> with long sepal points; coastal heathy woodland at Anglesea (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Angourie'	<i>Oligochaetochilus</i>	NSW	aff. <i>woollsii</i> with strongly recurved dorsal sepal tail, lateral sepal tails curving forwards then drooping and recurved basal lobe on labellum; coast Banksia & tea-tree on moist sandy soils (Bishop 2000)
<i>Pterostylis</i> species 'Arkaroola'	<i>Oligochaetochilus</i>	SA	aff. <i>biseta</i> with smaller flowers with more rounded lateral sepals and a larger, broadly triangular labellum; rocky slopes and ledges in dry woodland in the northern Flinders Ranges (Bates 2011)
<i>Pterostylis</i> species 'Barrington Tops'	<i>Hymenochilus</i>	NSW	stout aff. <i>cyncephala</i> with larger more crowded flowers with a slender stalked labellum appendage; moist grassland around swamps on Barrington Tops (Bishop 2000)
<i>Pterostylis</i> species 'Beechworth'	<i>Oligochaetochilus</i>	Vic	aff. <i>boormanii</i> with less bristly, brown flowers; <i>Callitris</i> spp. woodland on granite hills around Beechworth (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Berrook'	<i>Oligochaetochilus</i>	Vic	small aff. <i>excelsa</i> ; mallee on sandy soils in the south-western Sunset Country near Berrook (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Bimbowrie'	<i>Oligochaetochilus</i>	SA	aff. <i>boormanii</i> with dark brown flowers and a broadly triangular labellum; under low shrubs in sheltered sites on rocky hills north of Olary (Bates 2011)
<i>Pterostylis</i> species 'bloated'	<i>Plumatichilos</i>	WA	late flowering aff. <i>barbata</i> with larger more bloated flowers; damp open forest and seasonally wet flats between Augusta and Mt Barker (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'bloated snail'	<i>Linguella</i>	WA	aff. <i>nana</i> with large 'bloated flowers; open forest on sandy soils between Armadale and Margaret River; included in Brown <i>et al.</i> 2008 but not in later books; possibly just a broad flowered form of <i>P.</i> species 'crinkled leaf'
<i>Pterostylis</i> species 'bluebush plains'	<i>Oligochaetochilus</i>	SA	tall aff. <i>biseta</i> with narrower petals and lateral sepals; widespread across inland bluebush & saltbush plains; <i>thamnophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Border Ranges'	<i>Bunochilus</i>	Qld	aff. <i>melagramma</i> with an obovate labellum; open forest in Border Ranges region (Jones & Clements 2006)
<i>Pterostylis</i> species 'Boyup'	<i>Plumatichilos</i>	WA	early flowering aff. <i>plumosa</i> with light green flowers with a shorter, upturned dorsal sepal tip and a more densely bristled labellum; woodland between Augusta and Tenterden (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'bristly lip'		Vic	small non-colony forming aff. <i>nutans</i> with less nodding flower with a very bristly labellum; forest in south-western Victoria flowering after disturbance such as grading or fire (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'Broughton River'	<i>Oligochaetochilus</i>	SA	small red & green aff. <i>biseta</i> with short lateral sepals and fleshy labellum with a grooved base; woodland on rocky hills in Broughton River gorge and southern Flinders Ranges; probably extinct (Bates 2011)
<i>Pterostylis</i> species 'Buccarumbi'	<i>Diplodium</i>	NSW	aff. <i>revoluta</i> with very long, thin flowers with strongly projecting sinus; open forest around Buccarumbi (L. Copeland unpubl.)
<i>Pterostylis</i> species 'canegrass'	<i>Oligochaetochilus</i>	SA NSW	aff. <i>boormanii</i> with paler flowers with a more sparsely bristled labellum; grows amongst Triodia on red sand dunes on sandy plains between Renmark and Olary (Bates 2011)
<i>Pterostylis</i> species 'Cape Le Grande'	<i>Linguella</i>	WA	aff. <i>nana</i> with broad fleshy leaves in rosette and uniformly thickened lateral sepals; coastal scrubs along moist drainage lines and on granite rock sheets between Esperance and Cape Arid (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'cauline leaves'	<i>Linguella</i>	WA	aff. <i>nana</i> with fleshy rosette, robust habit with short thickened lateral sepals; woodland and open forest on sandy soils between Perth and Walpole (Brown <i>et al.</i> 2013)

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis</i> species 'chocolate'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> type with dark brown flowers; in shallow soil pockets on granite outcrops near Karonia (Brown <i>et al.</i> 2008); not included in later books but still thought to be distinct
<i>Pterostylis</i> species 'Chudalup'	<i>Linguella</i>	WA	aff. <i>nana</i> with sparsely leafy rosette, short stem with narrow flower with short clubbed lateral sepals; moss mats on granite rocks near Northcliff (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'chunky flowers'	<i>Diplodium</i>	SA Vic	aff. <i>dolichochila</i> with a shorter labellum; coastal scrubs from Kingston (SA) to Portland (Vic) (Bates 2011; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Coaldale'	<i>Oligochaetochilus</i>	NSW	aff. <i>boormanii</i> with paler flowers with narrower lateral sepals with more sparsely bristled outer margins and a broader labellum; open forest on sandy soils over sandstone near Coaldale (L. Copeland unpubl.)
<i>Pterostylis</i> species 'coastal'	<i>Urochilus</i>	WA	aff. <i>vitatta</i> with smaller, more widely spaced flowers; scrubs and woodland from Dongara to Bunbury (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'coastal clubbed sepals'	<i>Linguella</i>	WA	aff. <i>nana</i> with short stem and bloated flowers with very short clubbed lateral sepals; widespread on south-west coast, in scrubs, heath and woodland on sandy soils (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'coastal SA'	<i>Linguella</i>	SA	aff. <i>nana</i> with large raised crinkled basal leaves and large yellow-green flowers aging to brownish; coastal scrubs over limestone SA; <i>paractites</i> ms (Bates 2011)
<i>Pterostylis</i> species 'coastal snail'	<i>Linguella</i>	WA	aff. <i>nana</i> with short stem and somewhat bloated flowers with short thickened lateral sepals; coastal scrubs on sand dunes between Bremer Bay and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Coonabarabran'	<i>Oligochaetochilus</i>	NSW	aff. <i>praeternissa</i> with vertical, divergent shallowly concave shortly pointed lateral sepals wider than the galea, labellum with ~12 pairs of marginal hairs; <i>Callitris</i> woodland from Coonabarabran to Warialda (Bishop 2000)
<i>Pterostylis</i> species 'crinkled leaf'	<i>Linguella</i>	WA	aff. <i>nana</i> with crinkled leaves and tall slender flower with pointed galea and pointed tips to the lateral sepals; widely distributed between Perth and Albany, inland to Hyden, in woodland and open forest (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'dainty brown'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with small reddish brown with an extensively translucent hood and short forward-curved tips to the lateral sepals; open woodland on rocky hills and around salt lakes between Kalbarri and Eyre (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Daylesford'	<i>Diplodium</i>	Vic	aff. <i>deflexa</i> with smaller flowers with only gently curved galea, tip just below horizontal, not pointing to ground; foothill forest between Woodend and Ballarat (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'Dorrigo'	<i>Speculantha</i>	NSW	short, small, late spring-early summer flowering aff. <i>rubescens</i> with plump, extensively brown flowers; wet scrub and rainforest margins near Dorrigo (L. Copeland unpublished) <i>rubiginosa</i> ms
<i>Pterostylis</i> species 'Dundonnell'	<i>Hymenochilus</i>	Vic	aff. <i>bicolor</i> with very short spike of densely crowded flowers, basalt plains grasslands near Dundonnell (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'dwarf'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with small dark green flowers with short sepal tips; widespread in shrublands and mallee heath across inland sth-west WA (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'East Gippsland'	<i>Speculantha</i>	Vic	aff. <i>rubescens</i> with slightly larger, plumper flowers; heath and heathy open forest in East Gippsland (AVH record) <i>protera</i> ms
<i>Pterostylis</i> species 'East Kangaloon'	<i>Bunochilus</i>	NSW	aff. <i>smaragdyna</i> with a narrow elongated labellum; open forest at East Kangaloon (Jones & Clements 2006)
<i>Pterostylis</i> species 'Ebor Falls'	<i>Diplodium</i>	NSW	aff. <i>elegans</i> with very small flowers; single location in damp open forest at Ebor (L. Copeland unpublished) ( <i>micranthum</i> ms)
<i>Pterostylis</i> species 'Esperance'	<i>Plumatichilos</i>	WA	short, small aff. <i>plumosa</i> with light green flowers with long galea point and long dense bristles on the labellum; mallee woodland, shrubland and on granite rocks between Munglinup and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Everard Range'	<i>Oligochaetochilus</i>	SA	winter flowering aff. <i>excelsa</i> with extremely long free points to the lateral sepals; dry woodland on rocky hills, almost certainly extinct (Bates 2011)
<i>Pterostylis</i> species 'Eyre'	<i>Urochilus</i>	WA	aff. <i>vitatta</i> with fewer smaller paler flowers; shrubland and woodland on sand dunes, limestone cliffs sth of Cocklebidy (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Eyre Peninsula'	<i>Hymenochilus</i>	SA	aff. <i>cycnocephala</i> with short habit, small striped flowers with prominent labellum appendage & crenulate bracts; mallee and dry woodland on the Eyre Peninsula; <i>cymbella</i> ms (Bates 2011)
<i>Pterostylis</i> species 'fine sepals'	<i>Linguella</i>	WA	aff. <i>nana</i> with small flowers with very long slender lateral sepals; wandoo woodland between Perth and Bindoon (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Flinders Island'	<i>Bunochilus</i>	Tas	aff. <i>chlorogramma</i> with a broadly rectangular labellum; heathy woodland on Flinders Island (Jones & Clements 2006)

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis</i> species 'Flinders'	<i>Oligochaetochilus</i>	SA	tall aff. <i>excelsa</i> with smaller, drooping flowers on long slender pedicels; mallee and <i>Callitris</i> woodland on rocky hills in the Flinders Ranges between Hawker and Narrina (Bates 2011)
<i>Pterostylis</i> species 'Flinders Ranges north 1'	<i>Oligochaetochilus</i>	SA	unknown affiliation, mentioned in NOSSA Journal 39(11) Dec 2015
<i>Pterostylis</i> species 'Flinders Ranges north 2'	<i>Oligochaetochilus</i>	SA	unknown affiliation, mentioned in NOSSA Journal 39(11) Dec 2015
<i>Pterostylis</i> species 'foothill forest'	<i>Diplodium</i>	Vic	autumn flowering aff. <i>alveata</i> with more protruding sinus; foothill forests in East Gippsland east from Bruthen (Jeanes & Backhouse 2006; Turner <i>et al.</i> 2014)
<i>Pterostylis</i> species 'Gammon Range'	<i>Oligochaetochilus</i>	SA	aff. <i>excelsa</i> with larger flowers with a narrower labellum and very long sepal free points; <i>Callitris</i> woodland and dry shrubland in the northern Flinders Ranges from Mt Fitton to Hawker (Bates 2011)
<i>Pterostylis</i> species 'Gippsland'	<i>Plumatichilos</i>	Vic	aff. <i>plumosa</i> with smaller flowers with a green apical knob; heathy woodland and open forest in East Gippsland between Bruthen and Genoa (Turner <i>et al.</i> 2014)
<i>Pterostylis</i> species 'Goulburn'	<i>Diplodium</i>	NSW	small aff. <i>revoluta</i> with a very obtuse sinus; hilly woodland and open forest north of Goulburn (Bishop 2000)
<i>Pterostylis</i> species 'granite bird'	<i>Plumatichilos</i>	WA	aff. <i>plumosa</i> with very long sepal free points and the dorsal sepal tip upcurved; shrubland on granite outcrops between Albany and Shannon (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'greasy'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with smaller, nodding, 'greasy' reddish green to greenish flowers with a short upcurved tip to the dorsal sepal; woodland and shrubland between Wubin and Norseman (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'greater Grampians'	<i>Speculantha</i>	Vic	aff. <i>rubescens</i> ; woodland on gravelly clay soils in the Ararat-Stawell region and northern Grampians; no further information available (AVH record) <i>aurifera</i> ms
<i>Pterostylis</i> species 'green flowers'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with large, fairly crowded, nodding green flowers with a large spoon-shaped labellum; shrubland on granite outcrops between Salmon Gums and Menzies (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Greenbushes'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with large pale flowers constricted & flattened in middle, a broad lower opening & sigmoid-shaped dorsal sepal; woodland and open forest between Dardanup and Augusta (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Griffith'	<i>Oligochaetochilus</i>	NSW	small green to brownish aff. <i>biseta</i> with oblong angular labellum virtually lacking a basal constriction; mallee and dry woodland north of Griffith (Bishop 2000)
<i>Pterostylis</i> species 'Guildford'	<i>Bunochilus</i>	Vic	aff. <i>macrosepala</i> with a distinctly rectangular labellum with a very short midlobe; open forest at Guildford (Jones & Clements 2006)
<i>Pterostylis</i> species 'Gull Rock'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with long narrow flowers with very long sepal free points; heathy woodland and coastal scrubs near Albany (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Gundiah'		Qld	no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Helena River'	<i>Linguella</i>	WA	aff. <i>nana</i> with large broad flowers with extended dorsal sepal and narrowly clubbed lateral sepals; winter-wet sites in woodland and coastal heaths, Perth to Dunsborough (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'hooked sepals'	<i>Oligochaetochilus</i>	NSW Qld	reddish aff. <i>hamata</i> with longer dorsal sepal point and sparser labellum hairs; <i>Callitris</i> woodland on rocky hills in the ranges, tablelands and western slopes of northern NSW (= 'Northern Tablelands' Bishop 2000) and southern Qld
<i>Pterostylis</i> species 'inland'	<i>Linguella</i>	WA	tall aff. <i>nana</i> with a hairy stem; widespread in inland areas in woodland and shrublands (Brown <i>et al.</i> 2013); uncertain how it differs from and is almost certainly the same as <i>P. nana</i>
<i>Pterostylis</i> species 'Joanna'	<i>Diplodium</i>	SA	aff. <i>alata</i> with smaller, narrower flowers with a small opening; heathy woodland near Narracoorte (Bates 2011)
<i>Pterostylis</i> species 'Joiners Ridge'		Qld	no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Lara'	<i>Oligochaetochilus</i>	Vic	aff. <i>biseta</i> with short, thick stem and smallish flowers; basalt plains grasslands near Lara, probably extinct (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'large curved flowers'	<i>Diplodium</i>	Vic NSW ACT Qld	aff. <i>revoluta</i> with larger, more curved flowers; widespread in dry woodland and open forest through inland south-eastern Australia (Bishop 2000; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'large fawn'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with tawny nodding flowers with forward-pointing lateral sepal tips and a broad, ovate labellum; dry woodland and shrubland on rocky soils, also granite outcrops near Karonie (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'large striped flowers'	<i>Oligochaetochilus</i>	SA Vic	aff. <i>biseta</i> with large brownish striped flowers, mallee, <i>Callitris</i> woodland and dry woodland from Charlton (Vic) to Renmark (SA) (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'late flowering'	<i>Linguella</i>	WA	late spring-flowering aff. <i>nana</i> with short stems and clubbed lateral sepals; damp open forest near Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'limestone'	<i>Hymenochilus</i>	SA	aff. <i>cycnocephala</i> with prominent labellum appendage and distinctly striped flowers; woodland and scrubs on clay and sandy soils over limestone on the Yorke Peninsula and Murray mallee region; <i>calpicola</i> ms (Bates 2011)

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis</i> species 'limestone sands'	<i>Linguella</i>	WA	tall aff. <i>nana</i> with large flowers; near coastal woodland on well-drained calcareous sands from Yanchep to Busselton; included in Hoffman & Brown 2011 although not in later books; very similar to <i>P.</i> sp. 'Helena River'
<i>Pterostylis</i> species 'long sepals'	<i>Linguella</i>	WA	short, small aff. <i>nana</i> with long narrow lateral sepals and small, compact rosette of pointed leaves; moss mats on granite outcrops between Denmark and Windy Harbour (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'lower Murray'	<i>Oligochaetochilus</i>	SA	aff. <i>boormanii</i> with darker flowers with forward curved lateral sepals free points and a more rounded labellum; Callitris woodland on rocky areas along the lower Murray River, probably extinct (Bates 2011)
<i>Pterostylis</i> species 'Maldon'	<i>Oligochaetochilus</i>	Vic	aff. <i>biseta</i> with a short stem bearing very large, striped flowers with a bristly lip; sparsely shrubby woodland on granite hills near Maldon (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'mallee heath'	<i>Plumatichilos</i>	SA Vic	smallish glossy aff. <i>plumosa</i> with long pedicel and long dense labellum hairs; mallee heath on sandy to clay loam soils in western Vic & sth-eastern SA (Bates 2011; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'mallee snail'	<i>Linguella</i>	SA Vic	short aff. <i>nana</i> with small squat flowers with short lateral sepals; mallee on sandy to clay loam soils (Backhouse <i>et al.</i> 2016; Bates 2011) <i>tenuis</i> ms
<i>Pterostylis</i> species 'Malua Bay'	<i>Bunochilus</i>	NSW	aff. <i>macrosepala</i> ; heathy woodland at Malua Bay (Jones & Clements 2006)
<i>Pterostylis</i> species 'Mambray Creek'	<i>Bunochilus</i>	SA	aff. <i>chlorogramma</i> with a broadly rectangular labellum; dry woodland at Mambray Creek (Jones & Clements 2006)
<i>Pterostylis</i> species 'Mehinup'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with a very sparsely hairy labellum and widely spreading lateral sepals with upturned tips; on and around granite outcrops, Denmark to Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Mokota'	<i>Hymenochilus</i>	SA	aff. <i>cycnocephala</i> with short, stocky habit, thick fleshy leaves and barely striped flowers; montane grassland near Mt Bryan; <i>anemophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Mount Duval'	<i>Diplodium</i>	NSW	small aff. <i>alveata</i> with decurved petal tips and broad labellum with rounded apex just visible above the sinus; moist eucaypt forest near Armidale and Dorrigo (Bishop 2000)
<i>Pterostylis</i> species 'Mount Geharty'	<i>Oligochaetochilus</i>	SA	aff. <i>excelsa</i> with slender flowers with long free points to the sepals; dry shrubland on rocky outcrops on the Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'Mount Kaputar'	<i>Bunochilus</i>	NSW	aff. <i>lineata</i> with long, wide lateral sepals and a dark brown labellum with short bristles; open forest in the Nandewar Range
<i>Pterostylis</i> species 'Mount Maroon'		Qld	no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Mount Mee'		Qld	no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Mount Moffat'		Qld	no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Mount Olinthus'	<i>Oligochaetochilus</i>	SA	short aff. <i>excelsa</i> with rigid flowers with a small basal rosette and an anvil-shaped labellum; dry shrubland on rocky ridges on the Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'Mount Victoria Uranium Mine'	<i>Oligochaetochilus</i>	SA	dwarf aff. <i>boormanii</i> with smaller green flowers with flatter, hooked lateral sepals; sparse shrublands on granite hills north of Olary (Bates 2011)
<i>Pterostylis</i> species 'Muir's Highway'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with very long sepal free points and labellum bristles in opposite pairs; woodland on sandy soils along the Muir's Highway (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Murchison'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with nodding flowers with bright red lateral sepals with green inner margins and forward curved tips; dry woodland and shrubland on sandy soils between Murchison and Shark Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Narrandera range'	<i>Oligochaetochilus</i>	NSW	large aff. <i>petrosa</i> with narrow lateral sepals with long tips and a more rounded labellum with a prominent basal constriction; dry woodland on rocky hills near Narrandera (Bishop 2000)
<i>Pterostylis</i> species 'New England striped'	<i>Diplodium</i>	NSW	aff. <i>alata</i> with narrow striped flowers with angled basally flat lateral sepals in profile with backswept tips; forest and woodland on the New England Tableland (Bishop 2000)
<i>Pterostylis</i> species 'Newland Head'	<i>Diplodium</i>	SA	aff. <i>dolichochila</i> with a more widely opening flowers, a protruding sinus and shorter labellum; coastal scrubs on Kangaroo Island and the lower south-east (Bates 2011)
<i>Pterostylis</i> species 'Nora Creina'	<i>Diplodium</i>	SA Vic	aff. <i>alata</i> with tricolored flowers, a long fine point on dorsal sepal and a distinct gap between lateral sepals and galea; coastal scrubs in the lower south-east SA and sth-west Vic (Bates 2011)
<i>Pterostylis</i> species 'north-east Vic hills'	<i>Oligochaetochilus</i>	Vic	aff. <i>aciculiformis</i> with long lateral sepals and broader more bristly labellum; dry woodland on rocky hills in the north-east (Jeanes & Backhouse 2006); falls within the variation of and doubtfully distinct from <i>Pterostylis aciculiformis</i>
<i>Pterostylis</i> species 'northern'	<i>Urochilus</i>	WA	short aff. <i>vitatta</i> with fewer smaller flowers; dense shrubland on sandy soils between Catby and Binu (Brown <i>et al.</i> 2013)

Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis</i> species 'northern thick sepals'	<i>Linguella</i>	WA	aff. <i>nana</i> with broad fleshy rosette and short uniformly thickened lateral sepals; wandoo woodland between Perth and Cataby (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'north-west mallee'	<i>Oligochaetochilus</i>	Vic	aff. <i>excelsa</i> with large green flowers; mallee on sandy to clay loam soils in the Big Desert and southern Sunset Country (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'Old Boolcoolmata'	<i>Oligochaetochilus</i>	SA	aff. <i>lingua</i> with larger, dark maroon flowers with a larger labellum; Callitris woodland along drainage lines in rocky hills north of Olary (Bates 2011); <i>lingua</i> subspecies <i>ampla</i> ms
<i>Pterostylis</i> species 'Omeo'	<i>Bunochilus</i>	Vic	aff. <i>smaragdyna</i> with a narrowly obovate labellum; montane open forest near Omeo (Jones & Clements 2006)
<i>Pterostylis</i> species 'Oratan Rock'	<i>Oligochaetochilus</i>	SA	short aff. <i>excelsa</i> with very small flowers with hooked lateral sepals; shrublands on rocky hills near Lillydale (Bates 2011)
<i>Pterostylis</i> species 'Outback'	<i>Oligochaetochilus</i>	SA NSW	aff. <i>biseta</i> with broader, rounded lateral sepals; sparse shrublands on rocky hills between Renmark and Olary (SA) and near Broken Hill (NSW) (Bates 2011)
<i>Pterostylis</i> species 'Paynes Find'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with smallish nodding red-striped flowers with cupped lateral sepals with forward-pointing tips; sparse shrublands on granite outcrops and rocky breakaways between Wubin and Mount Magnet (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'plumed'	<i>Plumatichilos</i>	WA	aff. <i>plumosa</i> with smallish, dark green flowers with very short free point on dorsal sepal; woodland and shrubland between Esperance and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'quartz'	<i>Oligochaetochilus</i>	SA	aff. <i>lingua</i> with thick, fleshy labellum; sparse shrublands with <i>Triodia</i> on rocky quartz ridges near Olary (Bates 2011)
<i>Pterostylis</i> species 'Ranking Springs'	<i>Oligochaetochilus</i>	NSW	small aff. <i>biseta</i> with short tips to the lateral sepals; Callitris woodland on hills around Rankin Springs (Bishop 2000)
<i>Pterostylis</i> species 'Ravensthorpe'	<i>Linguella</i>	WA	aff. <i>nana</i> with moderately tall stem and large bloated flowers with moderately long lateral sepals; open woodland and shrubland between the Stirling Range and Esperance (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'red lip'	<i>Bunochilus</i>	Vic	aff. <i>smaragdyna</i> with large, rectangular red labellum; foothills open forest between Tonimbuk and Mallacoota (Backhouse <i>et al.</i> 2016; Turner <i>et al.</i> 2015)
<i>Pterostylis</i> species 'robust'	<i>Linguella</i>	WA	robust, spring flowering aff. <i>nana</i> with a blunt galea and thickened lateral sepals; damp sites in open forest on heavy clay soils (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'rock ledges'	<i>Oligochaetochilus</i>	SA	aff. <i>aciculiformis</i> with greenish flowers with longer free points to the sepals; open woodland on rocky hills in the upper south-east and Adelaide Hills (Bates 2011)
<i>Pterostylis</i> species 'rocky ranges'	<i>Hymenochilus</i>	SA	aff. <i>mutica</i> with short stocky habit and crowded, barely striped flowers; shrubland and dry woodland on rocky ranges in central SA; <i>pagophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'sandheath'	<i>Speculantha</i>	SA	aff. <i>rubescens</i> from heathy woodland near Glencoe (Bates 2011); possibly the same as <i>Pterostylis rubescens</i> as the differences are unclear
<i>Pterostylis</i> species 'sandstone heath'	<i>Speculantha</i>	NSW	summer flowering short aff. <i>parviflora</i> with crowded green, brown and white flowers with lateral sepals bowed in profile; heathland on shallow sandy soils and moss mats on sandstone, south of Sydney (Bishop 2000)
<i>Pterostylis</i> species 'short sepals'	<i>Linguella</i>	WA	aff. <i>nana</i> with short stem and narrow flowers with short lateral sepals; heath and heathy woodland on sandy soils between Perth and Yanchee (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'skinny'	<i>Linguella</i>	WA	aff. <i>nana</i> with tall, slender habit and tall, narrow flowers with long slender lateral sepals; open forest and woodland between Perth and Mt Barker (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'slender desert'	<i>Oligochaetochilus</i>	SA	small, slender aff. <i>excelsa</i> with pale flowers with short free points to the sepals; sparse shrubland on red soils around granite boulders near Olary (Bates 2011)
<i>Pterostylis</i> species 'small bands'	<i>Urochilus</i>	WA	aff. <i>vitatta</i> with fewer, smaller flowers with rounded lateral sepals; shrublands and woodlands between Reagans Ford and Three Springs (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'south-east Queensland'	<i>Oligochaetochilus</i>	Qld	aff. <i>chaetophora</i> from woodland and open forest on steep rocky hills; Gayndah to Tamborine Mountain; Qld records of <i>P. chaetophora</i> are this taxon
<i>Pterostylis</i> species 'southern Eyre Peninsula'	<i>Oligochaetochilus</i>	SA	aff. <i>excelsa</i> with a raised rosette of fleshy, crinkled strongly veined leaves and darker flowers with a broader labellum; sparse shrublands on rocky outcrops in the southern Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'southern granites'	<i>Linguella</i>	WA	aff. <i>nana</i> with small rosette, short stem and long lateral sepals; moss mats on granite outcrops between Manjimup and Hopetoun (Brown <i>et al.</i> 2013)



Australian Species	Alternate Genus	Distribution	Notes
<i>Pterostylis</i> species 'sparse'	<i>Plumatichilos</i>	WA	aff. <i>barbata</i> with very sparse bristles on labellum; woodland and open forest between Dardanup and Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Spring Ridge'	<i>Oligochaetochilus</i>	NSW	short aff. <i>biseta</i> with small flowers; dry woodland and open forest at Spring Ridge
<i>Pterostylis</i> species 'straight tops'	<i>Oligochaetochilus</i>	WA	aff. <i>rufa</i> with nodding mostly translucent flowers with almost solid red lateral sepals with straight upper margins and moderately long tips; widely distributed in inland woodland and shrublands often on rocky areas (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'tablelands'	<i>Diplodium</i>	NSW	colourful aff. <i>reflexa</i> with semi-circle curved galea with decurved apex and labellum just protruding over the broad protruding sinus; grassy forest from Bungonia to New England (Bishop 2000)
<i>Pterostylis</i> species 'tall mallee snail'	<i>Linguella</i>	SA	aff. <i>nana</i> with tall hairless stem and narrow green flowers with long light brownish sepal tips; inland mallee and dry woodland (Bates 2011); <i>naniflora</i> ms
<i>Pterostylis</i> species 'tall purple'	<i>Diplodium</i>	SA	aff. <i>dolichochila</i> with darker flowers with a narrower labellum; coastal scrubs and heaths in the south-east and possibly Kangaroo Island (Bates 2011)
<i>Pterostylis</i> species 'tall woodland'	<i>Oligochaetochilus</i>	SA	tall aff. <i>biseta</i> with long pedicels, narrow galea, distinctive lateral sepals with a double bend and a much broader labellum; woodland on fertile soils in the Mount Lofty and Flinders Ranges (Bates 2011); <i>sigmoidea</i> ms
<i>Pterostylis</i> species 'The Knoll'	<i>Plumatichilos</i>	WA	late flowering aff. <i>plumosa</i> with very long bristles on labellum; woodland and open forest between Albany and Dardanup (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'tiny flowers'	<i>Hymenochilus</i>	SA	aff. <i>mutica</i> with tall slender habit and small, spaced barely striped flowers; widely distributed in shrublands and woodland; <i>pisinna</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Torrington'	<i>Oligochaetochilus</i>	NSW	small aff. <i>praetermissa</i> with tiny, almost rounded labellum and lateral sepals wider than galea, labellum with ~8 pairs of marginal hairs; tea-tree thickets in northern granite region (Bishop 2000)
<i>Pterostylis</i> species 'upper Spencer Gulf'	<i>Oligochaetochilus</i>	SA	aff. <i>excelsa</i> ; no other information (NOSSA Journal 37(9): 94 (2013))
<i>Pterostylis</i> species 'Urbenville'	<i>Bunochilus</i>	NSW	aff. <i>barringtonensis</i> with a larger labellum with short, stiff hairs; open forest in the far northern volcanic ranges around Urbenville
<i>Pterostylis</i> species 'Warialda'	<i>Oligochaetochilus</i>	NSW	'markedly different' aff. <i>boormanii</i> ; dry woodland near Warialda (Bishop 2000)
<i>Pterostylis</i> species 'Warren snail'	<i>Linguella</i>	SA	aff. <i>nana</i> with pale, fleshy, smooth, ovate leaves, a hairless stem, swollen stem bracts and inflated base to the flower; open woodland in the Adelaide Hills (Bates 2011)
<i>Pterostylis</i> species 'Wauchope'	<i>Speculantha</i>	NSW	autumn flowering, robust aff. <i>rubescens</i> with larger flowers long, narrow, bristly dorsal sepal and smooth petals and lateral sepals; lowland open forest on gravelly soils on the north coast (recta ms)
<i>Pterostylis</i> species 'Werrikimbe'	<i>Speculantha</i>	NSW	robust aff. <i>rubescens</i> with bristly flowers; moist open forest on escarpments in the Barrington Tops and Werrikimbe; <i>oresbia</i> ms
<i>Pterostylis</i> species 'western Victoria'	<i>Oligochaetochilus</i>	Vic	small aff. <i>aciculiformis</i> with strongly backswept lateral sepals; dry woodland and mallee, sometimes in rocky areas, in the west (Bishop 2000)
<i>Pterostylis</i> species 'western woodlands'	<i>Oligochaetochilus</i>	Vic	early summer flowering aff. <i>biseta</i> with smaller, less prominently striped flowers; dry woodlands in western Vic (Melb to Stawell to Portland)
<i>Pterostylis</i> species 'Wilson's Promontory'		Vic	small aff. <i>alpina</i> ; coastal scrubs on sand dunes at Tidal River Wilsons Promontory (Jeanes & Backhouse 2006); probably only a form of <i>P. alpina</i>
<i>Pterostylis</i> species 'wrinkled lateral sepals'	<i>Oligochaetochilus</i>	NSW	aff. <i>praetermissa</i> with wrinkled lateral sepals; woodland and open forest above 500 m; open forest on rocky hills in the Barrington Tops, New England Tableland and Warrumbungles (L. Copeland unpubl.)
<i>Pterostylis</i> species 'woodland'	<i>Plumatichilos</i>	SA	aff. <i>plumosa</i> with ovate, bluntly pointed rosette leaves; widespread in woodland, the common form of <i>plumosa</i> in South Australia (Bates 2011)
<i>Pterostylis</i> species 'woodlands'	<i>Hymenochilus</i>	SA Vic	tall aff. <i>cycnocephala</i> with striped, spaced flowers; widespread in woodlands across SA and in west Vic; <i>nemoralis</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Woohlpooer'	<i>Bunochilus</i>	Vic	aff. <i>melagramma</i> with a thin labellum which has a large basal lobe; grassy woodlands near Woohlpooer (Jones & Clements 2006)
<b><i>Pyrorchis</i> D.L.Jones &amp; M.A.Clem.</b>			
<i>Pyrorchis forrestii</i> (F.Muell.) D.L.Jones & M.A.Clem.		WA	
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		WA SA Vic Tas NSW	

Australian Species	Alternate Genus	Distribution	Notes
<b>Rhinerrhiza</b> Rupp			
<i>Rhinerrhiza divitiflora</i> (F.Muell. ex Benth.) Rupp		NSW Qld	
<i>Rhinerrhizopsis matutina</i> D.L.Jones & M.A.Clem.	<i>Sarcochilus</i>	Qld	formerly <i>Sarcochilus moorei</i>
<b>Rhipidorchis</b> D.L.Jones & M.A.Clem.			
<i>Rhipidorchis micrantha</i> (A.Rich.) D.L.Jones & M.A.Clem.		Qld	
<b>Rhizanthella</b> R.S.Rogers			
<i>Rhizanthella gardneri</i> R.S.Rogers		WA	
<i>Rhizanthella omissa</i> D.L.Jones & M.A.Clem.		Qld	
<i>Rhizanthella slateri</i> (Rupp) M.A.Clem. & P.J.Cribb		NSW	
<i>Rhizanthella</i> species 'Munglinup'		WA	winter flowering aff. <i>gardneri</i> with fewer, larger, paler coloured flowers; under dense <i>Melaleuca</i> scrubs on sandy loam soils near Munglinup; <i>johnstonii</i> MS (Brown <i>et al.</i> 2013)
<b>Rhomboda</b> Lindl.			
<i>Rhomboda polygonoides</i> (F.Muell.) Ormerod	<i>Zeuxine</i>	Qld	
<b>Rimacola</b> Rupp			
<i>Rimacola elliptica</i> (R.Br.) Rupp		NSW	
<b>Robiquetia</b> Gaudich.			
<i>Robiquetia gracilistipes</i> (Schltr.) J.J.Sm.		Qld	
<i>Robiquetia wassellii</i> Dockrill		Qld	
<b>Saccolabiopsis</b> J.J.Sm.			
<i>Saccolabiopsis armitii</i> (F.Muell.) Dockrill		Qld	
<i>Saccolabiopsis rectifolia</i> (Dockrill) Garay		Qld	
<b>Salacistic</b> Rchb.f.			
<i>Salacistic ochroleuca</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Goodyera</i>	Qld	
<b>Sarcanthopsis</b> Garay			
<i>Sarcanthopsis warocqueana</i> (Rolfe) Garay		AIT	Moa Island, Torres Strait
<b>Sarcochilus</b> R.Br.			
<i>Sarcochilus aequalis</i> D.L.Jones & M.A.Clem.		NSW	
<i>Sarcochilus argochilus</i> D.L.Jones & M.A.Clem.		Qld	called <i>S. olivaceous</i> in Queensland
<i>Sarcochilus australis</i> (Lindl.) Rchb.f.		Vic Tas NSW	
<i>Sarcochilus borealis</i> (Nicholls) M.A.Clem.& D.L.Jones		Qld	called <i>S. olivaceous</i> in Queensland
<i>Sarcochilus ceciliae</i> F.Muell.		Qld	
<i>Sarcochilus dilatatus</i> F.Muell.		NSW Qld	
<i>Sarcochilus eriochilus</i> Fitzg.		NSW Qld	
<i>Sarcochilus falcatus</i> R.Br.		Vic NSW Qld	
<i>Sarcochilus fitzgeraldii</i> F.Muell.		NSW Qld	
<i>Sarcochilus hartmannii</i> F.Muell.		NSW Qld	
<i>Sarcochilus hillii</i> (F.Muell.) F.Muell.		NSW Qld	
<i>Sarcochilus hirticalcar</i> (Dockrill) M.A.Clem. & B.J.Wallace		Qld	
<i>Sarcochilus minutiflos</i> F.M.Bailey		Qld	
<i>Sarcochilus parviflorus</i> Lindl.		NSW Qld	called <i>S. olivaceous</i> in Queensland
<i>Sarcochilus roseus</i> (Clemesha) Clemesha		Qld	
<i>Sarcochilus serrulatus</i> D.L.Jones		Qld	
<i>Sarcochilus spathulatus</i> R.S.Rogers		NSW Qld	
<i>Sarcochilus weinthalii</i> F.M.Bailey		NSW Qld	
<i>Sarcochilus</i> species 'Blue Knob'		NSW	aff. <i>hartmannii</i> with erect, dark coloured racemes of all white flowers; sheltered rocky outcrops and cliffs in the northern ranges (Jones 2006)
<i>Sarcochilus</i> species 'brittle leaves'		NSW Qld	late flowering aff. <i>falcatus</i> with larger pale cupped flowers and brittle leaves; wet mountain forest and rainforest from Dorrigo NSW to the Border Ranges Qld
<b>Schistotylus</b> Dockrill			
<i>Schistotylus purpuratus</i> (Rupp) Dockrill		NSW Qld	
<b>Schoenorchis</b> Blume			
<i>Schoenorchis micrantha</i> Reinwardt ex Blume		Qld	
<i>Schoenorchis sarcophylla</i> Schltr.		Qld	
<b>Spathoglottis</b> Blume			
<i>Spathoglottis paulinae</i> F.Muell.		Qld NT	

Australian Species	Alternate Genus	Distribution	Notes
<i>Spathoglottis plicata</i> Blume		Qld	
<b>Spiculaea</b> Lindl.			
<i>Spiculaea ciliata</i> Lindl.		WA	
<b>Spiranthes</b> Rich.			
<i>Spiranthes alticola</i> D.L.Jones		SA Vic Tas NSW ACT	very difficult to distinguish from (and doubtfully distinct from) <i>Spiranthes australis</i>
<i>Spiranthes australis</i> (R.Br.) Lindl.		Vic Tas NSW Qld	
<i>Spiranthes elytra</i> D.L.Jones & R.J.Bates		SA	
<i>Spiranthes sinensis</i> (Pers.) Ames		Qld	uncertain whether this is 'true' <i>sinensis</i> or <i>Spiranthes australis</i>
<b>Taeniophyllum</b> Blume			
<i>Taeniophyllum clementsii</i> (D.L.Jones & B.Gray) Kocyan & Schuit.	<i>Microtatorchis</i>	Qld	
<i>Taeniophyllum confertum</i> B.Gray & D.L.Jones		Qld	
<i>Taeniophyllum cymbiforme</i> T.E.Hunt		Qld	
<i>Taeniophyllum hasseltii</i> Rchb.f.		AIT	Christmas Island
<i>Taeniophyllum lobatum</i> Dockrill		Qld	
<i>Taeniophyllum malianum</i> Schltr.		Qld	
<i>Taeniophyllum muelleri</i> Benth.		NSW Qld	
<i>Taeniophyllum norfolkianum</i> D.L.Jones, B.Gray & M.A.Clem.		AIT	Norfolk Island
<i>Taeniophyllum wilkianum</i> T.E.Hunt		Qld	
<b>Tainia</b> Blume			
<i>Tainia trinervis</i> (Blume) Rchb.f.		Qld	
<b>Thelasis</b> Blume			
<i>Thelasis capitata</i> Blume		AIT	Christmas Island
<i>Thelasis carinata</i> Blume		Qld	
<b>Thelymitra</b> J.R.Forst. & G.Forst.			
<i>Thelymitra adorata</i> Jeanes		NSW	
<i>Thelymitra aggericola</i> D.L.Jones		Tas	
<i>Thelymitra albiflora</i> Jeanes		SA	
<i>Thelymitra alcockii</i> D.L.Jones ex Jeanes		SA Vic	
<i>Thelymitra alpicola</i> Jeanes		Vic NSW	
<i>Thelymitra alpina</i> Jeanes		Vic NSW ACT	
<i>Thelymitra angustifolia</i> R.Br.		NSW Qld	
<i>Thelymitra antennifera</i> (Lindl.) Hook.f.		WA SA Vic Tas	
<i>Thelymitra apiculata</i> (A.S.George) M.A.Clem. & D.L.Jones		WA	
<i>Thelymitra arenaria</i> Lindl.		SA Vic Tas NSW ACT	
<i>Thelymitra aristata</i> Lindl.		SA Vic Tas	
<i>Thelymitra atronitida</i> Jeanes		Vic Tas NSW	
<i>Thelymitra azurea</i> R.S.Rogers		SA Vic	
<i>Thelymitra basaltica</i> Jeanes		Vic	
<i>Thelymitra batesii</i> Jeanes		SA	
<i>Thelymitra benthamiana</i> Rchb.f.		WA SA Vic Tas	
<i>Thelymitra bracteata</i> J.Z.Weber ex Jeanes		SA Vic Tas	
<i>Thelymitra brevifolia</i> Jeanes		SA Vic Tas NSW ACT	
<i>Thelymitra brevifolia</i> Jeanes subspecies 'Porters'		SA	has a paler, non-cleft column apex than <i>brevifolia</i> ; possibly a stabilised <i>T. albiflora</i> x <i>T. brevifolia</i> hybrid? (Bates 2011)
<i>Thelymitra campanulata</i> Lindl.		WA	
<i>Thelymitra canaliculata</i> R.Br.		WA	
<i>Thelymitra carnea</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Thelymitra circumsepta</i> Fitzg.		SA Vic Tas NSW	
<i>Thelymitra cornicina</i> Rchb.f.		WA	
<i>Thelymitra corrugata</i> R.J.Bates		SA	
<i>Thelymitra crenulata</i> R.J.Bates		SA	
<i>Thelymitra crinita</i> Lindl.		WA	
<i>Thelymitra cucullata</i> Rupp		WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Thelymitra cyanapicata</i> Jeanes		SA	
<i>Thelymitra cyanea</i> (Lindl.)		SA Vic Tas NSW ACT	
<i>Thelymitra dedmaniarum</i> R.S.Rogers		WA	
<i>Thelymitra epipactoides</i> F.Muell.		SA Vic	
<i>Thelymitra erosa</i> D.L.Jones & M.A.Clem.		Tas	
<i>Thelymitra exigua</i> Jeanes		SA Vic Tas	
<i>Thelymitra flexuosa</i> Endl.		WA SA Vic Tas	
<i>Thelymitra fragrans</i> D.L.Jones & M.A.Clem.		NSW Qld	
<i>Thelymitra frenchii</i> Jeanes		WA	virtually identical to and very difficult to separate from <i>T. granitora</i>
<i>Thelymitra fuscolutea</i> R.Br.		WA	
<i>Thelymitra glaucophylla</i> Jeanes		SA	
<i>Thelymitra graminea</i> Lindl.		WA	
<i>Thelymitra grandiflora</i> Fitzg. subspecies <i>exposa</i> R.J.Bates		SA	
<i>Thelymitra grandiflora</i> Fitzg. subspecies <i>grandiflora</i>		SA	
<i>Thelymitra granitora</i> D.L.Jones & M.A.Clem.		WA	
<i>Thelymitra gregaria</i> D.L.Jones & M.A.Clem.		Vic	
<i>Thelymitra hiemalis</i> D.L.Jones & M.A.Clem.		Vic	
<i>Thelymitra holmesii</i> Nicholls		SA Vic Tas NSW	
<i>Thelymitra hygrophila</i> R.J.Bates		SA	
<i>Thelymitra imbricata</i> D.L.Jones & M.A.Clem.		Tas	
<i>Thelymitra improcera</i> D.L.Jones & M.A.Clem.		Vic Tas	
<i>Thelymitra incurva</i> Jeanes		Vic NSW	
<i>Thelymitra inflata</i> Jeanes		SA Tas	
<i>Thelymitra ixioides</i> Sw.		SA Vic Tas NSW Qld	
<i>Thelymitra jacksonii</i> Hopper & A.P.Brown ex Jeanes		WA	
<i>Thelymitra jonesii</i> Jeanes		Tas	
<i>Thelymitra juncifolia</i> Lindl.		SA Vic Tas NSW ACT	
<i>Thelymitra kangaloonica</i> Jeanes		NSW	
<i>Thelymitra latifolia</i> R.J.Bates		SA Vic NSW	
<i>Thelymitra latiloba</i> Jeanes		WA	
<i>Thelymitra longifolia</i> J.R.Forst. & G.Forst.		AIT	Norfolk Island
<i>Thelymitra longiloba</i> D.L.Jones & M.A.Clem.		Vic Tas NSW Qld	
<i>Thelymitra lucida</i> Jeanes		SA Vic Tas	
<i>Thelymitra luteocillium</i> Fitzg.		SA Vic	
<i>Thelymitra mackibbinii</i> F.Muell.		Vic	
<i>Thelymitra macrophylla</i> Lindl.		WA	
<i>Thelymitra maculata</i> Jeanes		WA	
<i>Thelymitra magnifica</i> Jeanes		WA	
<i>Thelymitra malvina</i> M.A.Clem.		SA Vic Tas NSW Qld	
<i>Thelymitra matthewsii</i> Cheesem.		SA Vic	
<i>Thelymitra media</i> R.Br.		Vic NSW	
<i>Thelymitra megalyptra</i> Fitzg.		Vic NSW ACT	
<i>Thelymitra mucida</i> Fitzg.		WA SA Vic Tas	
<i>Thelymitra nuda</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Thelymitra occidentalis</i> Jeanes		WA SA	
<i>Thelymitra odora</i> R.J.Bates		SA	
<i>Thelymitra odora</i> R.J.Bates subspecies 'horse manure'		SA	unnamed subsp. of <i>odora</i> with strong horse manure-like odour (Bates 2011)
<i>Thelymitra orientalis</i> R.J.Bates		SA Vic	
<i>Thelymitra pallidiflora</i> Jeanes		Vic	
<i>Thelymitra pallidifructus</i> R.J.Bates		SA Vic	
<i>Thelymitra paludosa</i> Jeanes		WA	

Australian Species	Alternate Genus	Distribution	Notes
<i>Thelymitra pauciflora</i> R.Br.		SA Vic Tas NSW ACT Qld	
<i>Thelymitra peniculata</i> Jeanes		SA Vic Tas NSW ACT	
<i>Thelymitra petrophila</i> Jeanes		WA	
<i>Thelymitra planicola</i> Jeanes		Vic NSW	
<i>Thelymitra polychroma</i> D.L.Jones & M.A.Clem.		SA Tas	
<i>Thelymitra psammophila</i> C.R.P.Andrews		WA	
<i>Thelymitra pulcherrima</i> Jeanes		WA	
<i>Thelymitra purpurata</i> Rupp		NSW Qld	
<i>Thelymitra queenslandica</i> Jeanes		Qld	
<i>Thelymitra reflexa</i> Jeanes		Vic	
<i>Thelymitra rubra</i> Fitzg.		SA Vic Tas NSW ACT	
<i>Thelymitra rubricaulis</i> R.J.Bates		SA	
<i>Thelymitra sargentii</i> R.S.Rogers		WA	
<i>Thelymitra silena</i> D.L.Jones		Tas	
<i>Thelymitra simulata</i> D.L.Jones & M.A.Clem.		Vic Tas NSW ACT	
<i>Thelymitra spadicea</i> D.L.Jones & M.A.Clem.		Tas	
<i>Thelymitra sparsa</i> D.L.Jones & M.A.Clem.		Tas	
<i>Thelymitra speciosa</i> Jeanes		WA	
<i>Thelymitra spiralis</i> (Lindl.) F.Muell.		WA	
<i>Thelymitra stellata</i> Lindl.		WA	
<i>Thelymitra tigrina</i> R.Br.		WA	
<i>Thelymitra uliginosa</i> Jeanes		WA	
<i>Thelymitra variegata</i> (Lindl.) F.Muell.		WA	
<i>Thelymitra venosa</i> R.Br.		NSW	
<i>Thelymitra versicolor</i> Lindl.		Tas	often treated as a synonym of <i>T. nuda</i> , although considered a distinct species by Jones & Clements (1998) and Jones <i>et al.</i> (1999)
<i>Thelymitra villosa</i> Lindl.		WA	
<i>Thelymitra viridis</i> Jeanes		Tas	
<i>Thelymitra vulgaris</i> Jeanes		WA	
<i>Thelymitra xanthotricha</i> Jeanes		WA	
<i>Thelymitra yorkensis</i> Jeanes		WA	
<i>Thelymitra</i> species 'blue column'		NSW	aff. <i>pauciflora</i> with poorly opening purple-blue flowers with a purple-blue column apex; heathy woodland near Guyra (L. Copeland unpublished)
<i>Thelymitra</i> species 'central coast'		NSW	affinity uncertain; recognised by the purple violet flowers with similar colour hair tufts and the narrow horseshoe-shaped yellow apex to the post-anther lobe; heath and heathy woodland around Lake Macquarie
<i>Thelymitra</i> species 'Claire'		SA	short aff. <i>pauciflora</i> with two stem bracts and dark brown column apex with a black collar; woodlands in the Adelaide hills (Bates 2011)
<i>Thelymitra</i> species 'fine spots'		NSW ACT	aff. <i>ixioides</i> with self-pollinating flowers with few very fine spots on dorsal sepal and petals and irregularly glandular post-anther lobe; damp mountain forest, swampy and damp grassy areas above 800 m (Jones <i>et al.</i> 2008)
<i>Thelymitra</i> species 'Gibraltar Falls'		NSW ACT	no information available (Australian National Herbarium record)
<i>Thelymitra</i> species 'Glen Shera'		SA	tiny aff. <i>polychroma</i> with a purplish truncate post-anther lobe, sparse spots and tiny hair tufts; damp sites in heathland on acid sands on the Fleurieu Peninsula and lower South East (Bates 2011)
<i>Thelymitra</i> species 'insipid'		SA Vic	very slender, late flowered aff. <i>pauciflora</i> with long narrow sheathing leaf, rarely opening dull blue flowers and flat red and yellow post-anther lobe; widely distributed in many habitats in cooler, higher rainfall areas (Bates 2011)
<i>Thelymitra</i> species 'Meadows'		SA	aff. <i>pauciflora</i> with scabrid leaves and very rounded flowers with a truncate post-anther lobe; grassy woodland on steep rocky slopes in the southern Mount Lofty Ranges (Bates 2011)
<i>Thelymitra</i> species 'Murchison'		WA	aff. <i>macrophylla</i> with clumping habit and smaller, paler flowers; seasonally moist heathlands between Geraldton and Nerrin Nerrin (Brown <i>et al.</i> 2013)
<i>Thelymitra</i> species 'northern swamps'		WA	short, early-flowering aff. <i>paludosa</i> ; swamps between Perth and Bunbury (Brown <i>et al.</i> 2008 but not in later books); may just be a northern form of <i>T. paludosa</i>



Australian Species	Alternate Genus	Distribution	Notes
<i>Thelymitra</i> species 'Ongerup'		WA	aff. <i>yorkensis</i> with solid bicoloured flowers and more ornate column wings; heathy woodland near Ongerup (Brown <i>et al.</i> 2013)
<i>Thelymitra</i> species 'ornate flowers'		NSW ACT	small self-pollinating aff. <i>simulata</i> with freely opening flowers with shortly hooded deeply notched post-anther lobe; open forest in mountains above 750 m; resembles the New Zealand species <i>T. decora</i> (Jones <i>et al.</i> 2008)
<i>Thelymitra</i> species 'papillose'		SA	small aff. <i>ixioides</i> with papillose post-anther lobe and mop-like hair tufts; heath on damp peaty sands in the lower south-east (Bates 2011)
<i>Thelymitra</i> species 'red gum swamps'		SA Vic Tas	aff. <i>inflata</i> with more circular and narrower column apex; seasonal red gum swamps (SA, Vic) or wet forest (Tas) (Bates 2011)
<i>Thelymitra</i> species 'reed sun orchid'		WA	large aff. <i>petrophila</i> with a longer leaf; called aff. <i>macrophylla</i> in Hoffman & Brown 2011; grows amongst sedges not reeds; possibly just a robust form of <i>T. petrophila</i>
<i>Thelymitra</i> species 'Topperwein'		SA Vic	aff. <i>holmesii</i> with horseshoe-shaped, rounded column apex and yellow hair tufts; sedge swamps in River Red Gum woodland in the lower south-east of SA and western Vic (Bates 2011)
<i>Thelymitra</i> species 'Toy Creek'		Qld	no information available (Qld Herbarium record)
<b><i>Thrixspermum</i> Lour.</b>			
<i>Thrixspermum carinatifolium</i> (Ridl.) Schltr.		AIT	Christmas Island
<i>Thrixspermum congestum</i> (F.M.Bailey) Dockrill		Qld NT	
<i>Thrixspermum platystachys</i> (F.M.Bailey) Schltr.		Qld	
<b><i>Thynninorchis</i> D.L.Jones &amp; M.A.Clem.</b>			
<i>Thynninorchis huntianus</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	Vic Tas NSW ACT	
<i>Thynninorchis nothofagicola</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	Tas	
<b><i>Townsonia</i> Cheeseman</b>			
<i>Townsonia viridis</i> (Hook.f.) Schltr.		Tas	
<b><i>Trachoma</i> Garay</b>			
<i>Trachoma speciosum</i> D.L.Jones, B.Gray, M.A.Clem. & J.J.Wood		Qld	
<i>Trachoma stellatum</i> M.A.Clem., D.L.Jones, B.Gray & J.J.Wood		Qld	
<i>Trachoma subluteum</i> (Rupp) Garay		Qld	
<b><i>Trichoglottis</i> Blume</b>			
<i>Trichoglottis australiensis</i> Dockrill		Qld	
<b><i>Tropidia</i> Lindl.</b>			
<i>Tropidia territorialis</i> D.L.Jones & M.A.Clem.		NT	
<i>Tropidia viridifusca</i> Kraenzl.		AIT	Norfolk Island
<b><i>Vanda</i> Jones ex R.Br.</b>			
<i>Vanda hindsii</i> Lindl.		Qld	
<b><i>Vrydagzynea</i> Blume</b>			
<i>Vrydagzynea grayi</i> D.L.Jones & M.A.Clem.		Qld	
<b><i>Zeuxine</i> Lindl.</b>			
<i>Zeuxine exilis</i> Ridl.		AIT	Christmas Island
<i>Zeuxine oblonga</i> R.S.Rogers & C.T.White		WA NSW Qld NT	

**Total: 1,869 taxa (1,794 species and 116 subspecies and varieties)**

**Non-native (exotic or weed) orchids established in the wild in Australia**

<i>Arundina graminifolia</i> (D.Don) Hochr.	Qld	non-native species from south-east Asia, naturalised in Qld
<i>Disa bracteata</i> Sw.	WA SA Vic Tas	non-native species from southern African now established across southern Australia
<i>Epidendrum</i> species	NSW Qld	non-native species from Central & South America, widely cultivated, established in eastern Australia; variously identified as <i>E. elongatum</i> , <i>E. secundum</i> , <i>E. radicans</i> x <i>E. secundum</i> or <i>E. xobrienianum</i> ( <i>evectum</i> x <i>radicans</i> )
<i>Eulophia graminea</i> Lindl.	NT	non-native species from Asia grown as a garden plant and established around Darwin
<i>Serapias lingua</i> L.	SA	non-native species from southern Europe in cultivation in Australia and established in the Adelaide Hills South Australia (Bates 2011); eradication commenced in 2015
<i>Vanilla planifolia</i> Jacks. ex Andrews	Qld	non-native species from Central & South America; widely cultivated for production of vanilla essence; established in north Queensland (Jones 2006), although almost certainly by deliberate plantings rather than by seed spread

**Total: 6 species**

## Hybrid orchids recorded in Australia

<i>Caladenia ampla (reticulata) X cardiochila</i>		Vic	Backhouse 2011
<i>Caladenia ampla (reticulata) X clavigera</i>		Vic	G. Backhouse personal observation
<i>Caladenia ampla (reticulata) X fulva</i>		Vic	Backhouse 2011
<i>Caladenia ampla (reticulata) X grampiana</i>		Vic	Backhouse 2011
<i>Caladenia ancylosa X peisleyi</i>		Vic	Backhouse 2011
<i>Caladenia applanata X attingens</i>		WA	A. Brown personal observation
<i>Caladenia applanata X infundibularis</i>		WA	A. Brown personal observation
<i>Caladenia applanata X nivalis</i>		WA	A. Brown personal observation
<i>Caladenia arenaria X concinna</i>		NSW	Backhouse 2011
<i>Caladenia arenaria X rileyi</i>		NSW	Backhouse 2011
<i>Caladenia arenaria X stellata</i>		NSW	Backhouse 2011
<i>Caladenia arenaria X species 'inland NSW'</i>		NSW	Backhouse 2011
<i>Caladenia arenicola X huegelii</i>		WA	A. Brown personal observation
<i>Caladenia arenicola X longicauda</i>		WA	A. Brown personal observation
<i>Caladenia arenicola X lorea</i>		WA	Jones 2006
<i>Caladenia arenicola X swartziorum</i>		WA	Brown & Brockman 2015
<i>Caladenia argocalla X tentaculata</i>		SA	Bates 2011
<i>Caladenia attingens X barbarossa</i>		WA	Jones 2006
<i>Caladenia attingens X exstans</i>		WA	Jones 2006
<i>Caladenia attingens X georgei</i>		WA	Backhouse 2011
<i>Caladenia attingens X horistes</i>		WA	Brown & Brockman 2015
<i>Caladenia attingens X infundibularis</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia attingens X longicauda</i>		WA	Brown & Brockman 2015
<i>Caladenia attingens X procera</i>		WA	Jones 2006
<i>Caladenia attingens X thincicola</i>		WA	Jones 2006
<i>Caladenia aurulenta X clavula</i>		SA	Backhouse 2011
<i>Caladenia australis X venusta</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia barbarossa X caesarea</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia barbarossa X falcata</i>		WA	Backhouse 2011
<i>Caladenia barbarossa X lobata</i>		WA	A. Brown personal observation
<i>Caladenia barbarossa X longicauda</i>	<i>xenigma</i> Hopper & A.P. Br.	WA	
<i>Caladenia barbarossa X radiata</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia behrii X reticulata</i>		SA	Bates 2011
<i>Caladenia behrii X rigida</i>		SA	Bates 2011
<i>Caladenia bicallata X capillata</i>		SA	G. Backhouse personal observation
<i>Caladenia brevisura X cruscula</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia brevisura X doutchiae</i>		WA	Jones 2006
<i>Caladenia brevisura X horistes</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia brevisura X polychroma</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia brevisura X voigtii</i>		WA	Jones 2006
<i>Caladenia brownii X ferruginea</i>		WA	A. Brown personal observation
<i>Caladenia brownii X longicauda</i>		WA	Backhouse 2011
<i>Caladenia brownii X pectinata</i>		WA	A. Brown personal observation
<i>Caladenia brownii X serotina</i>		WA	G. Backhouse personal observation
<i>Caladenia brumalis X capillata</i>		SA	Bates 2011
<i>Caladenia brumalis X cardiochila</i>		SA	Bates 2011
<i>Caladenia brumalis X conferta</i>		SA	Bates 2011
<i>Caladenia brumalis X fuliginosa</i>		SA	Bates 2011
<i>Caladenia brumalis X latifolia</i>		SA	Bates 2011
<i>Caladenia brumalis X macroclavia</i>		SA	Bates 2011
<i>Caladenia brumalis X stricta</i>		SA	Bates 2011
<i>Caladenia busselliana X paludosa</i>		WA	Backhouse 2011
<i>Caladenia busselliana X procera</i>		WA	A. Brown personal observation
<i>Caladenia caesarea X polychroma</i>		WA	A. Brown personal observation
<i>Caladenia cairnsiana X dimidia</i>	<i>xericksoniae</i> Nicholls	WA	
<i>Caladenia cairnsiana X filifera</i>		WA	Jones 2006
<i>Caladenia cairnsiana X polychroma</i>		WA	A. Brown personal observation

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Caladenia cairnsiana</i> X <i>pulchra</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia cairnsiana</i> X <i>vulgata</i>		WA	Jones 2006
<i>Caladenia calcicola</i> X <i>cardiochila</i>		SA	Bates 2011
<i>Caladenia calcicola</i> X <i>colorata</i>		SA	Bates 2011
<i>Caladenia capillata</i> X <i>fuscata</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia capillata</i> X <i>toxochila</i>		SA	Bates 2011
<i>Caladenia capillata</i> X <i>zephyra</i>		SA	G. Backhouse personal observation
<i>Caladenia cardiochila</i> X <i>colorata</i>		SA	Backhouse 2011
<i>Caladenia cardiochila</i> X <i>clavigera</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia cardiochila</i> X <i>formosa</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia cardiochila</i> X <i>lowanensis</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia cardiochila</i> X <i>reticulata</i>		SA	plants on Kangaroo Island recognised in Bates (2011) as a species
<i>Caladenia cardiochila</i> X <i>septuosa</i>		SA	G. Backhouse personal observation
<i>Caladenia cardiochila</i> X <i>stricta</i>		SA Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia cardiochila</i> X <i>strigosa</i>		SA	Bates 2011
<i>Caladenia cardiochila</i> X <i>venusta</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia cardiochila</i> X <i>xanthochila</i>		SA	Backhouse 2011
<i>Caladenia cardiochila</i> X <i>zephyra</i>		SA	Bates 2011
<i>Caladenia cardiochila</i> X species 'mallee heath'		SA Vic	Bates 2011
<i>Caladenia carnea</i> X <i>catenata</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia carnea</i> X <i>cucullata</i>		Vic	Kuiter 2010
<i>Caladenia carnea</i> X <i>fuscata</i>		SA Vic	Jones <i>et al.</i> 1999
<i>Caladenia carnea</i> X <i>latifolia</i>		SA	Bates 2011
<i>Caladenia carnea</i> X <i>maritima</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia carnea</i> X <i>vulgaris</i>		SA Vic	Bates 2011
<i>Caladenia catenata</i> X <i>clarkiae</i>		Vic	Backhouse 2011
<i>Caladenia caudata</i> X <i>saggicola</i>		Tas	Backhouse 2011
<i>Caladenia chapmanii</i> X <i>longicauda</i>	× <i>eludens</i> Hopper & A.P. Br.	WA	
<i>Caladenia chapmanii</i> X <i>polychroma</i>		WA	Jones 2006
<i>Caladenia christineae</i> X <i>harringtoniae</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia christineae</i> X <i>latifolia</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia citrina</i> X <i>rhomboidiformis</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia clarkiae</i> X <i>hildae</i>		Vic	Jones 2006
<i>Caladenia clavigera</i> X <i>montana</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia clavigera</i> X <i>oenochila</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia clavigera</i> X <i>parva</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia clavigera</i> X <i>reticulata</i>		SA	G. Backhouse personal observation
<i>Caladenia clavula</i> X <i>stricta</i>		SA	G. Backhouse personal observation
<i>Caladenia clavula</i> X <i>zephyra</i>		SA	Backhouse 2011
<i>Caladenia clavula</i> X species 'short yellow clubs'		SA	Backhouse 2011
<i>Caladenia colorata</i> X <i>formosa</i>		Vic	Backhouse 2011
<i>Caladenia colorata</i> X <i>strigosa</i>		SA	Bates 2011
<i>Caladenia colorata</i> X <i>tensa</i>		SA	Bates 2011
<i>Caladenia colorata</i> X <i>verrucosa</i>		SA	Bates 2011
<i>Caladenia colorata</i> X species 'mallee heath'		SA Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia colorata</i> X species 'Murray mallee'		SA Vic	Bates 2011
<i>Caladenia conferta</i> X <i>macroclavia</i>		SA	Bates 2011
<i>Caladenia congesta</i> X <i>hildae</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia corynephora</i> X <i>serotina</i>	× <i>aestantha</i> Hopper & A.P. Br.	WA	
<i>Caladenia crebra</i> X <i>longicauda</i>	× <i>coactescens</i> Hopper & A.P. Br.	WA	
<i>Caladenia cretacea</i> X <i>cruciformis</i>		Vic	Backhouse 2011
<i>Caladenia cretacea</i> X <i>douglasiorum</i>		Vic	Backhouse 2011
<i>Caladenia cretacea</i> X <i>tentaculata</i>		Vic	Backhouse 2011
<i>Caladenia cristata</i> X <i>denticulata</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia cristata</i> X <i>vulgata</i>		WA	Jones 2006
<i>Caladenia cucullata</i> X <i>ustulata</i>		ACT	J. Egan & T. Wood pers. comm.
<i>Caladenia decora</i> X <i>longicauda</i>		WA	Backhouse 2011
<i>Caladenia denticulata</i> X <i>filifera</i>		WA	A. Brown personal observation

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Caladenia denticulata</i> X <i>footeana</i>		WA	Brown & Brockman 2015
<i>Caladenia dimidia</i> X <i>doutchiae</i>		WA	Backhouse 2011
<i>Caladenia dimidia</i> X <i>erythronema</i>		WA	A. Brown personal observation
<i>Caladenia dimidia</i> X <i>filifera</i>		WA	A. Brown personal observation
<i>Caladenia dimidia</i> X <i>footeana</i>		WA	A. Brown personal observation
<i>Caladenia dimidia</i> X <i>hirta</i>		WA	A. Brown personal observation
<i>Caladenia dimidia</i> X <i>melanema</i>		WA	A. Brown personal observation
<i>Caladenia dimidia</i> X <i>pachychila</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia dimidia</i> X <i>paradoxa</i>		WA	Jones 2006
<i>Caladenia dimidia</i> X <i>polychroma</i>		WA	Jones 2006
<i>Caladenia dimidia</i> X <i>radialis</i>		WA	Backhouse 2011
<i>Caladenia dimidia</i> X <i>roei</i>	× <i>xoleta</i> Hopper & A.P. Br.	WA	
<i>Caladenia discoidea</i> X <i>hirta</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia dorrienii</i> X <i>vulgata</i>		WA	A. Brown personal observation
<i>Caladenia doutchiae</i> X <i>exilis</i>		WA	Jones 2006
<i>Caladenia doutchiae</i> X <i>incrassata</i>		WA	Jones 2006
<i>Caladenia doutchiae</i> X <i>polychroma</i>		WA	Jones 2006
<i>Caladenia doutchiae</i> X <i>pulchra</i>		WA	Jones 2006
<i>Caladenia doutchiae</i> X <i>radialis</i>		WA	Backhouse 2011
<i>Caladenia doutchiae</i> X <i>vulgata</i>		WA	A. Brown personal observation
<i>Caladenia drakeoides</i> X <i>exilis</i> subsp. <i>exilis</i>	× <i>hopperi</i> J.M.H.Shaw	WA	the original name × <i>ornata</i> Hopper & A.P.Br. was invalid as <i>Caladenia ornata</i> was already named
<i>Caladenia drakeoides</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia drakeoides</i> X <i>vulgata</i>		WA	Jones 2006
<i>Caladenia dundasiae</i> X <i>vulgata</i>		WA	A. Brown personal observation
<i>Caladenia elegans</i> X <i>nobilis</i>		WA	Jones 2006
<i>Caladenia elegans</i> X <i>vulgata</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia erythrochila</i> X <i>straminichila</i>		WA	A. Brown personal observation
<i>Caladenia erythronema</i> X <i>polychroma</i>		WA	A. Brown personal observation
<i>Caladenia erythronema</i> X <i>vulgata</i>		WA	A. Brown personal observation
<i>Caladenia excelsa</i> X <i>longicauda</i>		WA	Jones 2006
<i>Caladenia exilis</i> X <i>roei</i>		WA	Jones 2006
<i>Caladenia exstans</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia falcata</i> X <i>integra</i>		WA	A. Brown personal observation
<i>Caladenia falcata</i> X <i>longicauda</i>	× <i>cala</i> Hopper & A.P. Br.	WA	
<i>Caladenia ferruginea</i> X <i>leucochila</i>		WA	Brown & Brockman 2015
<i>Caladenia ferruginea</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia ferruginea</i> X <i>pectinata</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia ferruginea</i> X <i>rhomboidiformis</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia filamentosa</i> X <i>montana</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia filifera</i> X <i>hiemalis</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia flava</i> X <i>hirta</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia flava</i> X <i>latifolia</i>	× <i>spectabilis</i> Hopper & A.P. Br.	WA	
<i>Caladenia flava</i> X <i>longicauda</i>	× <i>triangularis</i> R.S. Rogers	WA	
<i>Caladenia flava</i> X <i>marginata</i>	× <i>erminae</i> Hopper & A.P. Br.	WA	
<i>Caladenia flava</i> X <i>nana</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia flava</i> subspecies <i>flava</i> X <i>reptans</i> subspecies <i>reptans</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia flava</i> subspecies <i>maculata</i> X <i>reptans</i> subspecies <i>impensa</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia flavovirens</i> X <i>parva</i>		Vic	Backhouse 2011
<i>Caladenia flindersica</i> X <i>gladiolata</i>	× <i>ensigera</i> D.L. Jones	SA	
<i>Caladenia flindersica</i> X <i>saxatilis</i>		SA	Backhouse 2011
<i>Caladenia flindersica</i> X <i>tentaculata</i>		SA	Backhouse 2011
<i>Caladenia flindersica</i> X <i>toxochila</i>		SA	Bates 2011
<i>Caladenia fluvialis</i> X <i>footeana</i>		WA	A. Brown personal observation
<i>Caladenia footeana</i> X <i>hiemalis</i>		WA	Jones 2006
<i>Caladenia footeana</i> X <i>horistes</i>		WA	A. Brown personal observation
<i>Caladenia footeana</i> X <i>incensa</i>		WA	A. Brown personal observation



Australian Hybrids	Name and Authority	Distribution	Notes
<i>Caladenia footeana</i> X <i>nobilis</i>		WA	Jones 2006
<i>Caladenia footeana</i> X <i>remota</i>		WA	A. Brown personal observation
<i>Caladenia formosa</i> X <i>parva</i>		SA Vic	G. Backhouse personal observation
<i>Caladenia formosa</i> X <i>tentaculata</i>		Vic	Backhouse 2011
<i>Caladenia formosa</i> X <i>venusta</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia fuliginosa</i> X <i>stricta</i>		SA	Bates 2011
<i>Caladenia fulva</i> X species 'little clowns'		SA	Bates 2011
<i>Caladenia gardneri</i> X <i>interjacens</i>		WA	A. Brown personal observation
<i>Caladenia gardneri</i> X <i>latifolia</i>	× <i>idiastes</i> Hopper & A.P. Br.	WA	
<i>Caladenia georgei</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia georgei</i> X <i>speciosa</i>		WA	Jones 2006
<i>Caladenia georgei</i> X <i>swartziorum</i>		WA	Backhouse 2011
<i>Caladenia gladiolata</i> X <i>saxatilis</i>		SA	Bates 2011
<i>Caladenia gladiolata</i> X <i>viriosa</i>		SA	Bates 2011
<i>Caladenia gracilis</i> X <i>hildae</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia grampiana</i> X <i>parva</i>		Vic	Backhouse 2011
<i>Caladenia harringtoniae</i> X <i>longicauda</i>		WA	Backhouse 2011
<i>Caladenia heberleana</i> X <i>longicauda</i>		WA	Backhouse 2011
<i>Caladenia heberleana</i> X <i>pectinata</i>		WA	A. Brown personal observation
<i>Caladenia hiemalis</i> X <i>polychroma</i>		WA	Jones 2006
<i>Caladenia hiemalis</i> X <i>xantha</i>		WA	Jones 2006
<i>Caladenia hildae</i> X <i>moschata</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia hirta</i> X <i>longicauda</i>	× <i>suffusa</i> Hopper & A.P. Br.	WA	
<i>Caladenia hirta</i> X <i>pendens</i>		WA	A. Brown personal observation
<i>Caladenia hirta</i> X <i>remota</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia hirta</i> X <i>saxicola</i>		WA	A. Brown personal observation
<i>Caladenia hoffmanii</i> X <i>longicauda</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia hopperiana</i> X <i>longicauda</i>		WA	Brown & Brockman 2015
<i>Caladenia horistes</i> X <i>multiclavia</i>		WA	incorrectly named as × <i>resupina</i> , the type of which is <i>polychroma</i> X <i>multiclavia</i>
<i>Caladenia horistes</i> X <i>pachychila</i>		WA	Jones 2006
<i>Caladenia huegelii</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia huegelii</i> X <i>paludosa</i>		WA	A. Brown personal observation
<i>Caladenia huegelii</i> X <i>speciosa</i>		WA	Jones 2006
<i>Caladenia huegelii</i> X <i>swartziorum</i>		WA	Brown & Brockman 2015
<i>Caladenia incensa</i> X <i>incrassata</i>		WA	A. Brown personal observation
<i>Caladenia incensa</i> X <i>pachychila</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia incensa</i> X <i>radialis</i>		WA	A. Brown personal observation
<i>Caladenia incensa</i> X <i>roei</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia incensa</i> X <i>vulgata</i>		WA	Jones 2006
<i>Caladenia incrassata</i> X <i>pluvialis</i>		WA	A. Brown personal observation
<i>Caladenia infundibularis</i> X <i>lodgeana</i>		WA	A. Brown personal observation
<i>Caladenia integra</i> X <i>longicauda</i>		WA	A. Brown personal observation
<i>Caladenia interanea</i> X <i>zephyra</i>		SA	G. Backhouse personal observation
<i>Caladenia latifolia</i> X <i>longicauda</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia latifolia</i> X <i>nana</i>		WA	Jones 2006
<i>Caladenia leptochila</i> X <i>reticulata</i>		SA	Bates 2011
<i>Caladenia leucochila</i> X <i>longicauda</i>		WA	Brown & Brockman 2015
<i>Caladenia lobata</i> X <i>longicauda</i>	× <i>hypata</i> Hopper & A.P. Br.	WA	
<i>Caladenia lobata</i> X <i>macrostylis</i>		WA	A. Brown personal observation
<i>Caladenia lodgeana</i> X <i>serotina</i>		WA	A. Brown personal observation
<i>Caladenia longicauda</i> X <i>lorea</i>		WA	Backhouse 2011
<i>Caladenia longicauda</i> X <i>macrostylis</i>		WA	Jones 2006
<i>Caladenia longicauda</i> X <i>magniclavata</i>		WA	Backhouse 2011
<i>Caladenia longicauda</i> X <i>nobilis</i>		WA	Backhouse 2011
<i>Caladenia longicauda</i> X <i>pectinata</i>		WA	Backhouse 2011
<i>Caladenia longicauda</i> X <i>splendens</i>		WA	Hopper & Brown 2001
<i>Caladenia longicauda</i> X <i>stareorum</i>		WA	Backhouse 2011
<i>Caladenia longicauda</i> X <i>thinicola</i>		WA	Backhouse 2011

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Caladenia longicauda</i> X <i>uliginosa</i>	× <i>exserta</i> Hopper & A.P. Br.	WA	
<i>Caladenia longicauda</i> X <i>winfieldii</i>		WA	Backhouse 2011
<i>Caladenia lowanensis</i> X <i>stricta</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia lowanensis</i> X <i>tensa</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia luteola</i> X <i>pendens</i>		WA	A. Brown personal observation
<i>Caladenia luteola</i> X <i>polychroma</i>		WA	A. Brown personal observation
<i>Caladenia macroclavia</i> X <i>zephyra</i>		SA	G. Backhouse personal observation
<i>Caladenia macrotylis</i> X <i>pectinata</i>		WA	Jones 2006
<i>Caladenia marginata</i> X <i>nana</i>		WA	Jones 2006
<i>Caladenia marginata</i> X <i>reptans</i>		WA	Jones 2006
<i>Caladenia melanema</i> X <i>vulgata</i>		WA	A. Brown personal observation
<i>Caladenia microchila</i> X <i>sigmoidea</i>	× <i>tryphera</i> Hopper & A.P. Br.	WA	
<i>Caladenia multiclavia</i> X <i>polychroma</i>	× <i>resupina</i> Hopper & A.P. Br.	WA	
<i>Caladenia multiclavia</i> X <i>pulchra</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia nana</i> X <i>reptans</i>		WA	Jones 2006
<i>Caladenia necrophylla</i> X <i>venusta</i>		SA	Bates 2011
<i>Caladenia nobilis</i> X <i>occidentalis</i>		WA	Jones 2006
<i>Caladenia nobilis</i> X <i>vulgata</i>		WA	Hopper & Brown 2001
<i>Caladenia occidentalis</i> X <i>pendens</i>		WA	A. Brown personal observation
<i>Caladenia oenochila</i> X <i>rosella</i>		Vic	Backhouse 2011
<i>Caladenia oenochila</i> X <i>tentaculata</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia orientalis</i> X <i>tessellata</i>	× <i>variabilis</i> Nicholls	Vic	
<i>Caladenia osmera</i> X <i>peisleyi</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia pachychila</i> X <i>paradoxa</i>		WA	A. Brown personal observation
<i>Caladenia pachychila</i> X <i>vulgata</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia paludosa</i> X <i>speciosa</i>		WA	Jones 2006
<i>Caladenia paradoxa</i> X <i>pulchra</i>		WA	Jones 2006
<i>Caladenia paradoxa</i> X <i>radialis</i>		WA	A. Brown personal observation
<i>Caladenia paradoxa</i> X <i>roei</i>		WA	A. Brown personal observation
<i>Caladenia parva</i> X <i>stricta</i>		SA	Backhouse 2011
<i>Caladenia parva</i> X <i>venusta</i>		Vic	Jeanes & Backhouse 22006
<i>Caladenia peisleyi</i> X <i>tessellata</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia pendens</i> X <i>postea</i>		WA	A. Brown personal observation
<i>Caladenia pholcoidea</i> X <i>serotina</i>		WA	Jones 2006
<i>Caladenia pluvialis</i> X <i>radialis</i>		WA	Backhouse 2011
<i>Caladenia pluvialis</i> X <i>roei</i>		WA	A. Brown personal observation
<i>Caladenia polychroma</i> X <i>pulchra</i>		WA	Jones 2006
<i>Caladenia polychroma</i> X <i>roei</i>		WA	G. Backhouse personal observation
<i>Caladenia polychroma</i> X <i>xantha</i>		WA	Jones 2006
<i>Caladenia radialis</i> X <i>remota</i>		WA	A. Brown personal observation
<i>Caladenia radialis</i> X <i>roei</i>		WA	A. Brown personal observation
<i>Caladenia radiata</i> X <i>serotina</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia remota</i> X <i>roei</i>		WA	Brown <i>et al.</i> 2008
<i>Caladenia reticulata</i> X <i>rigida</i>		SA	Bates 2011
<i>Caladenia roei</i> X <i>voigtii</i>		WA	A. Brown personal observation
<i>Caladenia roei</i> X <i>vulgata</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia roei</i> X <i>wanosa</i>		WA	A. Brown personal observation
<i>Caladenia saxatilis</i> X <i>woolcockiorum</i>		SA	Bates 2011
<i>Caladenia septuosa</i> X <i>zephyra</i>		SA	Bates 2012
<i>Caladenia stricta</i> X <i>verrucosa</i>		Vic	Jeanes & Backhouse 2006
<i>Caladenia stricta</i> X <i>zephyra</i>		SA	Bates 2011
<i>Caladenia stricta</i> X species 'little clowns'		SA	Bates 2011
<i>Caladenia tentaculata</i> X species 'Raymond Island'		Vic	Backhouse 2011
<i>Caladenia tentaculata</i> X <i>venusta</i>		SA Vic	Jeanes & backhouse 2006
<i>Caladenia toxochila</i> X <i>zephyra</i>		SA	Bates 2011
<i>Caladenia toxochila</i> X <i>viriosa</i>		SA	Bates 2011
<i>Caladenia venusta</i> X <i>villosissima</i>		Vic	Backhouse 2011
<i>Caladenia verrucosa</i> X species 'little clowns'		SA	Bates 2011
<i>Caladenia verrucosa</i> X <i>viriosa</i>		SA	Bates 2011

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Caladenia vulgata</i> X <i>wanosa</i>		WA	Backhouse 2011
<i>Caladenia woolcockiorum</i> X <i>Xviriosa</i>		SA	Bates 2011
<i>Caladenia zephyra</i> X <i>viriosa</i>		SA	Backhouse 2011
<i>Caladenia zephyra</i> X species 'short yellow clubs'		SA	Bates 2011
<i>Caladenia carnea</i> X <i>Cyanicula caerulea</i>		ACT	ACT Orchid Group blog
<i>Caladenia capillata</i> X <i>Pheladenia deformis</i>		SA	Bates 2011
<i>Caladenia carnea</i> X <i>Pheladenia deformis</i>		SA	Bates 2011
<i>Caladenia coactilis</i> X <i>Pheladenia deformis</i>		SA	Bates 2011
<i>Caladenia dimidia</i> X <i>Pheladenia deformis</i>		WA	Brown <i>et al.</i> 2013
<i>Caladenia fuscata</i> X <i>Pheladenia deformis</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Caladenia hirta</i> X <i>Pheladenia deformis</i>		WA	A. Brown personal observation
<i>Caladenia latifolia</i> X <i>Pheladenia deformis</i>		SA	Bates 2011
<i>Caladenia radialis</i> X <i>Pheladenia deformis</i>		WA	Jones 2006
<i>Calochilus paludosus</i> X <i>robertsonii</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Chiloglottis cornuta</i> X <i>valida</i>		Vic	Jeanes & Backhouse 2006
<i>Chiloglottis trapeziformis</i> X <i>valida</i>	× <i>pescottiana</i> R.S. Rogers	Vic NSW ACT	
<i>Corunastylis archeri</i> X <i>despectans</i>		Vic	Jeanes & Backhouse 2006
<i>Corunastylis arrecta</i> X <i>despectans</i>		Vic	Jeanes & Backhouse 2006
<i>Corunastylis ciliata</i> X <i>despectans</i>		SA	Bates 2011
<i>Corunastylis citriodora</i> X <i>fimbriata</i>		NSW	Jones 2006
<i>Corunastylis despectans</i> X <i>morrisii</i>		Vic	Bates 2011
<i>Corunastylis filiformis</i> X <i>fimbriata</i>		NSW	Jones 2006
<i>Corybas despectans</i> X <i>diemenicus</i>		SA	Bates 2011
<i>Corybas despectans</i> X <i>expansus</i>		SA	Bates 2011
<i>Corybas despectans</i> X <i>incurvus</i>		SA	Bates 2011
<i>Corybas dilatatus</i> X <i>incurvus</i>	× <i>miscellus</i> D.L. Jones	SA	
<i>Cyanicula gemmata</i> X <i>sericea</i>		WA	A. Brown personal observation
<i>Cyanicula caerulea</i> X <i>Glossodia major</i>		Vic NSW	
<i>Cyanicula caerulea</i> X <i>Glossodia minor</i>		Vic NSW	Backhouse 2011
<i>Cyanicula amplexans</i> X <i>Pheladenia deformis</i>		WA	Brown <i>et al.</i> 2013
<i>Cyanicula caerulea</i> X <i>Pheladenia deformis</i>		Vic	Brown <i>et al.</i> 2013
<i>Cyanicula sericea</i> X <i>Elythranthera brunonis</i>	× <i>Cyanthera glossodioides</i> Hopper & A.P. Br.	WA	
<i>Dendrobium aemulum</i> X <i>kingianum</i>		NSW	Jones 2006
<i>Dendrobium antennatum</i> X <i>discolor</i>		Qld	Jones 2006
<i>Dendrobium antennatum</i> X <i>nindii</i>		Qld	Jones 2006
<i>Dendrobium banksii</i> X <i>pugioniforme</i>		NSW	Jones 2006
<i>Dendrobium biggibum</i> X <i>discolor</i>	× <i>superbiens</i> Rchb.f.	Qld	
<i>Dendrobium biggibum</i> X × <i>superbiens</i>	× <i>lavarackianum</i> M.A. Clem.	Qld	
<i>Dendrobium bowmanii</i> X <i>cucumerinum</i>		NSW	Jones 2006
<i>Dendrobium calamiformis</i> X <i>nugentii</i>	× <i>grimesii</i> C.T. White & Summerh.	Qld	
<i>Dendrobium calamiformis</i> X <i>rigidum</i>	× <i>foederata</i> St. Cloud	Qld	
<i>Dendrobium canaliculatum</i> X <i>johannis</i>		Qld	
<i>Dendrobium canaliculatum</i> X <i>semifusum</i>		Qld	
<i>Dendrobium canaliculatum</i> X <i>discolor</i>		Qld	
<i>Dendrobium cucumerinum</i> X <i>linguiformis</i>		NSW	Jones 2006
<i>Dendrobium discolor</i> X <i>johannis</i>		Qld	Jones 2006
<i>Dendrobium discolor</i> X <i>lithicola</i>	× <i>vinicolor</i> St. Cloud	Qld	only marginally distinct from and probably the same as <i>D. ×superbiens</i>
<i>Dendrobium discolor</i> X <i>nindii</i>		Qld	
<i>Dendrobium discolor</i> X <i>semifusum</i>		Qld	
<i>Dendrobium dolichophyllum</i> X <i>schoeninum</i>		NSW	Jones 2006
<i>Dendrobium fairfaxii</i> X <i>pugioniforme</i>		NSW	Jones 2006
<i>Dendrobium falcorostrum</i> X <i>gracilicaule</i>		NSW	Jones 2006
<i>Dendrobium falcorostrum</i> X <i>kingianum</i>		NSW	Jones 2006
<i>Dendrobium falcorostrum</i> X <i>speciosum</i>		NSW	Jones 2006
<i>Dendrobium gracilicaule</i> X <i>kingianum</i>	× <i>suffusum</i> Cady	NSW	
<i>Dendrobium gracilicaule</i> X <i>jonesii</i>		Qld	
<i>Dendrobium gracilicaule</i> X <i>speciosum</i>	× <i>gracillimum</i> (Rupp) Leaney	NSW Qld	
<i>Dendrobium jonesii</i> X <i>nitidum</i>		Qld	Jones 2006

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Dendrobium jonesii</i> X <i>speciosum</i>	× <i>rupiosum</i> Clemesha	Qld	
<i>Dendrobium kingianum</i> subsp. <i>kingianum</i> X <i>speciosum</i> subsp. <i>speciosum</i>	× <i>delicatum</i> (F.M. Bailey) F.M. Bailey	NSW Qld	
<i>Dendrobium kingianum</i> var. <i>pulcherrimum</i> X <i>speciosum</i> subsp. <i>speciosum</i>	× <i>kestevenii</i> Rupp	NSW	
<i>Dendrobium linguiformis</i> X <i>pugioniforme</i>		NSW	Jones 2006
<i>Dendrobium linguiformis</i> X <i>teretifolium</i>		NSW	Jones 2006
<i>Dendrobium mortii</i> X <i>pugioniforme</i>		NSW	Jones 2006
<i>Dendrobium mortii</i> X <i>schoeninum</i>		NSW	Jones 2006
<i>Dendrobium mortii</i> X <i>teretifolium</i>		NSW	Jones 2006
<i>Dendrobium nugentii</i> X <i>racemosum</i>		Qld	Jones 2006
<i>Dendrobium pugioniforme</i> X <i>schoeninum</i>		NSW	Jones 2006
<i>Dendrobium pugioniforme</i> X <i>striolatum</i>		NSW	Jones 2006
<i>Dendrobium striolatum</i> X <i>teretifolium</i>		NSW	Jones 2006
<i>Dipodium pardalinum</i> X <i>roseum</i>		SA Vic	Bates 2011; Backhouse <i>et al.</i> 2016
<i>Diuria alba</i> X <i>chrysantha</i>		Qld	
<i>Diuris abbreviata</i> X <i>eborensis</i>		NSW	
<i>Diuris abbreviata</i> X <i>goonooensis</i>		NSW	Jones 2006
<i>Diuris abbreviata</i> X <i>minor</i>		NSW	
<i>Diuris abbreviata</i> X <i>pedunculata</i>		NSW	
<i>Diuris abbreviata</i> X <i>venosa</i>		NSW	
<i>Diuris aurea</i> X <i>punctata</i>	× <i>nebulosa</i> D.L. Jones	NSW	
<i>Diuris basaltica</i> X <i>palustris</i>	× <i>fastidiosa</i> R.S. Rogers	Vic	
<i>Diuris behrii</i> X <i>orientis</i>		SA Vic	Bates 2011
<i>Diuris behrii</i> X <i>palustris</i>		SA	Bates 2011
<i>Diuris behrii</i> X <i>pardina</i>	× <i>palachila</i> R.S. Rogers	SA Vic Qld	Bates 2011; Jeanes & Backhouse 2006; Qld record from Mt Moffat; neither parent occurs within several hundred km of location so identification doubtful
<i>Diuris behrii</i> X <i>punctata</i>		Vic NSW	Backhouse, <i>et al.</i> 2016; Jones 2006
<i>Diuris behrii</i> X <i>sulphurea</i>		SA	Bates 2011
<i>Diuris behrii</i> X <i>tricolor</i>		NSW	G. Backhouse personal observation
<i>Diuris behrii</i> X species 'thick sepals'		Vic	
<i>Diuris brevifolia</i> X <i>orientis</i>		SA	Bates 2011
<i>Diuris brevifolia</i> X <i>pardina</i>		SA	Bates 2011
<i>Diuris calcicola</i> X <i>orientis</i>		SA	Bates 2016
<i>Diuris calcicola</i> X <i>palustris</i>		SA	Bates 2016
<i>Diuris chrysantha</i> X <i>minor</i>		NSW	Jones 2006
<i>Diuris chrysiopsis</i> X <i>nigromontana</i>		ACT	J. Egan & T. Wood pers. comm.
<i>Diuris chrysiopsis</i> X <i>orientis</i>		Vic	Jeanes & Backhouse 2006
<i>Diuris chrysiopsis</i> X <i>pardina</i>		SA Vic NSW ACT	Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Diuris chrysiopsis</i> X <i>palustris</i>		Vic	Jeanes & Backhouse 2006
<i>Diuris chrysiopsis</i> X <i>platichila</i>	× <i>polymorpha</i> Messmer	NSW	
<i>Diuris chrysiopsis</i> X <i>punctata</i>		Vic	Jeanes & Backhouse 2006
<i>Diuris corymbosa</i> X <i>laxiflora</i>		WA	Brown <i>et al.</i> 2013
<i>Diuris eborensis</i> X species 'New England swamps'		NSW	
<i>Diuris goonooensis</i> X <i>tricolor</i>		NSW	Jones 2006
<i>Diuris gregaria</i> X <i>palustris</i>		Vic	Jeanes & Backhouse 2006
<i>Diuris gregaria</i> X <i>pardina</i>		Vic	
<i>Diuris laxiflora</i> X <i>purdei</i>		WA	Brown <i>et al.</i> 2013
<i>Diuris monticola</i> X <i>pardina</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Diuris monticola</i> X <i>semilunulata</i>		ACT	J. Egan & T. Wood pers. comm.
<i>Diuris nigromontana</i> X <i>sulphurea</i>		ACT	J. Egan & T. Wood pers. comm.
<i>Diuris orientis</i> X <i>palustris</i>		SA	Bates 2011
<i>Diuris orientis</i> X <i>pardina</i>		SA Vic	Bates 2011; Jeanes & Backhouse 2006
<i>Diuris orientis</i> X <i>sulphurea</i>		Vic	Jeanes & Backhouse 2006
<i>Diuris palustris</i> X <i>pardina</i>		SA	Bates 2011
<i>Diuris pardina</i> X <i>subalpina</i>		Vic NSW	
<i>Diuris pardina</i> X <i>sulphurea</i>		Vic	Backhouse <i>et al.</i> 2016

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Diuris punctata</i> X <i>sulphurea</i>		NSW	
<i>Diuris punctata</i> X species 'Gippsland plains'		Vic	Backhouse <i>et al.</i> 2016
<i>Diuris septentrionalis</i> X <i>tinkeri</i>		WA	A. Brown personal observation
<i>Diuris septentrionalis</i> X species 'Zuytdorp cliffs'		WA	
<i>Diuris</i> species 'sandplains' X species 'Yalgorup'		WA	Brown <i>et al.</i> 2013
<i>Drakaea glyptodon</i> X <i>livida</i>		WA	
<i>Drakaea gracilis</i> X <i>livida</i>		WA	Jones 2006
<i>Drakaea livida</i> X <i>thynniphila</i>		WA	A. Brown personal observation
<i>Elythranthera brunonis</i> X <i>emarginata</i>	× <i>intermedia</i> (Fitzg.) M.A. Clem.	WA	
<i>Ericksonella saccharata</i> X <i>Pheladenia deformis</i>		WA	Brown <i>et al.</i> 2013
<i>Glossodia major</i> X <i>minor</i>		Vic NSW	Jeanes & Backhouse 2006
<i>Glossodia major</i> X <i>Pheladenia deformis</i>	× <i>Glossadenia tutelata</i> Kavulak	SA Vic Tas	
<i>Microtis alba</i> X <i>media</i>		WA	A. Brown personal observation
<i>Prasophyllum alpestre</i> X <i>sphacelatum</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Prasophyllum alpestre</i> X <i>tadgellianum</i>		Vic	Jeanes & Backhouse 2006
<i>Prasophyllum brownii</i> X <i>regium</i>		WA	Brown <i>et al.</i> 2013
<i>Prasophyllum chasmogamum</i> X <i>frenchii</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Prasophyllum fimbria</i> X <i>giganteum</i>		WA	Brown <i>et al.</i> 2013
<i>Prasophyllum niphopedium</i> X <i>sphacelatum</i>		Vic	Jeanes & Backhouse 2006
<i>Prasophyllum viretrum</i> X species 'Mortlake'		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis aciculiformis</i> X species 'Beechworth'		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis aestiva</i> X <i>atrans</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis aestiva</i> X <i>decurva</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis allantoidea</i> X species 'inland'		WA	A. Brown personal observation
<i>Pterostylis alpina</i> X <i>cucullata</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis alpina</i> X <i>foliata</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis alveata</i> X <i>ophioglossa</i>	× <i>furcillata</i> Rupp	NSW	
<i>Pterostylis atrans</i> X <i>decurva</i>		Vic NSW ACT	Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Pterostylis baptistii</i> X <i>falcata</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis baptistii</i> X <i>nutans</i>		Qld	
<i>Pterostylis boormanii</i> X <i>setifera</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis cheraphila</i> X <i>ferruginea</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>fischii</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>nutans</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>obtusata</i>		NSW	
<i>Pterostylis concinna</i> X <i>ophioglossa</i>	× <i>conoglossa</i> Upton	NSW	
<i>Pterostylis concinna</i> X <i>striata</i>	× <i>toveyana</i> Ewart & Sharman	Vic Tas	
<i>Pterostylis cucullata</i> X <i>falcata</i>	× <i>aenigma</i> D.L. Jones & M.A. Clem.	Vic	
<i>Pterostylis cucullata</i> X <i>foliata</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis cucullata</i> X <i>nutans</i>		SA Vic	Bates 2011
<i>Pterostylis curta</i> X <i>lustra</i>		SA	Bates 2011
<i>Pterostylis curta</i> X <i>nutans</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis curta</i> X <i>pedunculata</i>		SA Vic	Bates 2011; Jeanes & Backhouse 2006
<i>Pterostylis dolichochila</i> X <i>erythroconcha</i>		SA	Bates 2011
<i>Pterostylis dubia</i> X <i>scabrida</i>		Tas	Jones 2006
<i>Pterostylis falcata</i> X <i>nutans</i>	× <i>ingens</i> (Rupp) D.L. Jones	SA Vic Tas NSW	
<i>Pterostylis ferruginea</i> X <i>planulata</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis fischii</i> X <i>laxa</i>		Vic	Jeanes & Backhouse 2006
<i>Pterostylis lingua</i> X species 'large striped flowers'		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis lustra</i> X <i>nutans</i>		SA	Bates 2011
<i>Pterostylis nutans</i> X <i>pedunculata</i>		SA	Bates 2011
<i>Pterostylis parviflora</i> X <i>rubescens</i>		Vic	
<i>Pterostylis robusta</i> X <i>striata</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis robusta</i> X species 'Adelaide hills'		SA	Bates 2011
<i>Pterostylis truncata</i> X species 'large curved flowers'		Vic	Backhouse <i>et al.</i> 2016
<i>Pterostylis</i> species 'limestone' X species 'woodlands'		SA	Bates 2011 ( <i>callicola</i> ms X <i>nemoralis</i> ms)



Australian Hybrids	Name and Authority	Distribution	Notes
<i>Sarcochilus fitzgeraldii</i> X <i>hartmanii</i>		Qld	Jones 2006
<i>Thelymitra albiflora</i> X <i>antennifera</i>		SA	Jeanes 2004; Bates 2011
<i>Thelymitra albiflora</i> X <i>bracteata</i>		SA	Bates 2011
<i>Thelymitra albiflora</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra albiflora</i> X <i>pallidifructus</i>		SA	Bates 2011
<i>Thelymitra albiflora</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra alcockiae</i> X <i>antennifera</i>		SA Vic	Backhouse <i>et al.</i> 2016
<i>Thelymitra alcockiae</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra alpina</i> X <i>simulata</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Thelymitra antennifera</i> X <i>gregaria</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>bracteata</i>		SA	Bates 2011
<i>Thelymitra antennifera</i> X <i>graminea</i>		WA	Brown <i>et al.</i> 2013
<i>Thelymitra antennifera</i> X <i>inflata</i>		SA	Bates 2011
<i>Thelymitra antennifera</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra antennifera</i> X <i>luteocilium</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>macrophylla</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>maculata</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>petrophila</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>vulgaris</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>megcalyptera</i>		SA Vic	Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>nuda</i>	<i>xmacmillanii</i> F. Muell.	SA Vic	Bates 2011
<i>Thelymitra antennifera</i> X <i>pauciflora</i>		SA	Bates 2011
<i>Thelymitra antennifera</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra antennifera</i> X <i>vulgaris</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra apiculata</i> X <i>pulcherrima</i>		WA	
<i>Thelymitra arenaria</i> X <i>ixioides</i>		Vic	
<i>Thelymitra arenaria</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra arenaria</i> X <i>nuda</i>		Vic	Jeanes 2004
<i>Thelymitra aristata</i> X <i>ixioides</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra aristata</i> X <i>media</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra azurea</i> X <i>nuda</i>		SA	Jones 2006
<i>Thelymitra benthamiana</i> X <i>macrophylla</i>		WA	Brown <i>et al.</i> 2008
<i>Thelymitra benthamiana</i> X <i>villosa</i>		WA	Brown <i>et al.</i> 2013
<i>Thelymitra bracteata</i> X <i>ixioides</i>		SA	Bates 2011
<i>Thelymitra bracteata</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra bracteata</i> X <i>nuda</i>		SA	Bates 2011
<i>Thelymitra brevifolia</i> X <i>ixioides</i>		Vic	
<i>Thelymitra brevifolia</i> X <i>juncifolia</i>		SA NSW ACT	Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra brevifolia</i> X <i>ixioides</i>		SA	Bates 2011
<i>Thelymitra brevifolia</i> X <i>nuda</i>		SA	Bates 2011
<i>Thelymitra brevifolia</i> X <i>pauciflora</i>		SA	Bates 2011
<i>Thelymitra campanulata</i> X <i>spiralis</i>		WA	
<i>Thelymitra carnea</i> X <i>flexuosa</i>		NSW	Jones 2006
<i>Thelymitra carnea</i> X <i>ixioides</i>	<i>xirregularis</i> Nicholls	Vic Tas NSW	
<i>Thelymitra carnea</i> X <i>peniculata</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra carnea</i> X <i>rubra</i>		SA NSW	Bates 2011
<i>Thelymitra cornicina</i> X <i>crinita</i>		WA	Jones 2006
<i>Thelymitra corrugata</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra crenulata</i> X <i>ixioides</i>		SA	Bates 2011
<i>Thelymitra crenulata</i> X <i>malvina</i>		SA	Bates 2011
<i>Thelymitra crenulata</i> X <i>peniculata</i>		SA	Bates 2011
<i>Thelymitra crinita</i> X <i>macrophylla</i>		WA	Brown <i>et al.</i> 2013
<i>Thelymitra cyanapicata</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra cyanapicata</i> X <i>nuda</i>		SA	Bates 2011
<i>Thelymitra cyanapicata</i> X <i>pauciflora</i>		SA	Bates 2011
<i>Thelymitra cyanapicata</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra exigua</i> X <i>gregaria</i>		Vic	Jeanes & Backhouse 2006
<i>Thelymitra exigua</i> X <i>ixioides</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Thelymitra flexuosa</i> X <i>macrophylla</i>		WA	Jones 2006

Australian Hybrids	Name and Authority	Distribution	Notes
<i>Thelymitra flexuosa</i> X <i>vulgaris</i>		WA	Brown <i>et al.</i> 2013
<i>Thelymitra glaucopylla</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra holmesii</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra holmesii</i> X <i>pauciflora</i>		SA	Bates 2011
<i>Thelymitra hygrophila</i> X <i>odora</i>		SA	Bates 2011
<i>Thelymitra inflata</i> X <i>juncifolia</i>		SA	Bates 2011
<i>Thelymitra inflata</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra ixiooides</i> X <i>mucida</i>		SA	Jones 2006
<i>Thelymitra ixiooides</i> X <i>nuda</i>		NSW	Jones 2006
<i>Thelymitra ixiooides</i> X <i>pallidifructus</i>		SA	Bates 2010
<i>Thelymitra ixiooides</i> X <i>pauciflora</i>	x <i>truncata</i> R.S. Rogers	SA Vic Tas	
<i>Thelymitra ixiooides</i> X <i>peniculata</i>	x <i>merraniae</i> Nicholls	SA Vic Tas	
<i>Thelymitra ixiooides</i> X <i>rubra</i>		NSW	Jones 2006
<i>Thelymitra juncifolia</i> X <i>mucida</i>		SA	Bates 2011
<i>Thelymitra juncifolia</i> X <i>megcalyptra</i>		Vic	Bates 2011
<i>Thelymitra juncifolia</i> X <i>nuda</i>		SA	Bates 2011
<i>Thelymitra juncifolia</i> X <i>pauciflora</i>		SA NSW ACT	Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra juncifolia</i> X <i>peniculata</i>		ACT	Jones <i>et al.</i> 2008
<i>Thelymitra juncifolia</i> X <i>rubra</i>		SA Vic	Bates 2011
<i>Thelymitra juncifolia</i> X species 'pale buds'		SA	Bates 2011
<i>Thelymitra latiloba</i> X <i>graminea</i>		WA	Brown <i>et al.</i> 2013 as <i>latiloba</i> x <i>petrophila</i>
<i>Thelymitra luteocilium</i> X <i>megcalyptra</i>		SA Vic	Bates 2011; Jeanes & Backhouse 2006
<i>Thelymitra luteocilium</i> X <i>nuda</i>	x <i>chasmogama</i> R.S. Rogers	SA Vic	Bates 2011
<i>Thelymitra luteocilium</i> X <i>rubra</i>		SA Vic	Bates 2011
<i>Thelymitra luteocilium</i> X <i>pauciflora</i>		Vic	Backhouse <i>et al.</i> 2016
<i>Thelymitra macrophylla</i> X <i>vulgaris</i>		WA	Jones 2006
<i>Thelymitra nuda</i> X <i>pallidifructus</i>		SA	Bates 2010
<i>Thelymitra nuda</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra odora</i> subspecies 'horse manure' X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra pauciflora</i> X <i>rubra</i>		SA	Bates 2011
<i>Thelymitra sargentii</i> X species 'Murchison'		WA	A. Brown personal observation
<i>Thelymitra villosa</i> X species 'Murchison'		WA	Brown <i>et al.</i> 2013 as <i>macrophylla</i> x <i>villosa</i>

**Total: 546 hybrids**

Australian Capital Territory Species	Alternate Genus	Notes
<i>Acianthus collinus</i> D.L.Jones		
<i>Acianthus exsertus</i> R.Br.		
<i>Caladenia actensis</i> D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	
<i>Caladenia aestiva</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia alpina</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia armata</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia atrovessa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia cucullata</i> Fitzg.	<i>Stegostyla</i>	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia fitzgeraldii</i> Rupp	<i>Arachnorchis</i>	
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia hildae</i> E.E.Pescott & Nicholls	<i>Stegostyla</i>	
<i>Caladenia mentiensi</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia montana</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia moschata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia phaeoclavia</i> D.L.Jones	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia parva</i>
<i>Caladenia praecox</i> Nicholls	<i>Stegostyla</i>	
<i>Caladenia ustulata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia</i> species 'pointed flowers'	<i>Petalochilus</i>	aff. <i>mentiensi</i> with sharply pointed sepals and petals and incurved dorsal sepal; hilly dry woodland near Gungahlin (J. Egan & T. Wood personal communication)
<i>Calochilus montanus</i> D.L.Jones		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus sandrae</i> D.L.Jones		
<i>Calochilus saprophyticus</i> R.S.Rogers		
<i>Calochilus therophilus</i> D.L.Jones		
<i>Chiloglottis seminuda</i> D.L.Jones		
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmecchila</i>	
<i>Chiloglottis trilabra</i> Fitzg.		
<i>Chiloglottis turfosa</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis valida</i> D.L.Jones	<i>Simpliglottis</i>	'true' <i>valida</i> occurs in the mountains above 500 m but is morphologically indistinguishable from two related cryptic species (Bower 2007; see sp. 'Falls Creek' and sp. 'lowlands'), distribution overlaps with sp. 'Falls Creek'
<i>Chiloglottis</i> species 'highlands'	<i>Simpliglottis</i>	aff. <i>valida</i> with slightly smaller labellum with fewer calli and different wasp pollinator; montane damp forest and woodland above 800 m (Bower 2007; Jones <i>et al.</i> 2008); indistinguishable from <i>C. valida</i>
<i>Corunastylis arrecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis clivicola</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis cornuta</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis ectopa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nuda</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis turfosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corybas grumulus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas hispidus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	
<i>Cyanicula caerulea</i> (R.Br.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyrtostylis reniformi</i> R.Br.		
<i>Dipodium hamiltonianum</i> F.M.Bailey		
<i>Dipodium punctatum</i> (Sm.) R.Br.		
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		
<i>Diuris behrii</i> Schldl.		
<i>Diuris chryseopsis</i> D.L.Jones		
<i>Diuris dendrobioides</i> Fitzg.		
<i>Diuris monticola</i> D.L.Jones		
<i>Diuris nigromontana</i> D.L.Jones		

Australian Capital Territory Species	Alternate Genus	Notes
<i>Diuris pardina</i> Lindl		
<i>Diuris punctata</i> Sm. variety <i>punctata</i>		
<i>Diuris semilunulata</i> Messmer		
<i>Diuris subalpina</i> D.L.Jones		
<i>Diuris sulphurea</i> R.Br.		
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus magenteus</i> D.L.Jones		
<i>Eriochilus</i> species 'smooth leaf'		aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<i>Gastrodia entomogama</i> D.L.Jones		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Gastrodia surcula</i> D.L.Jones		
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Lyperanthus suaveolens</i> R.Br.		
<i>Microtis oblonga</i> R.S.Rogers		
<i>Microtis parviflora</i> R.Br.		
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		
<i>Microtis</i> species 'alpine'		aff. <i>unifolia</i> with fewer smaller spaced flowers with a smaller labellum; wet mountain meadows and woodland (Jones <i>et al.</i> 2008)
<i>Microtis</i> species 'subalpine grasslands'		short, summer flowering aff. <i>parviflora</i> with fewer smaller flowers that often lack the labellum apical callus; damp subalpine grasslands (Jones <i>et al.</i> 2008)
<i>Orthoceras strictum</i> R.Br.		
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Prasophyllum alpestre</i> D.L.Jones		
<i>Prasophyllum brevibracte</i> (Lindl.) Hook.f.		
<i>Prasophyllum canaliculatum</i> D.L.Jones		
<i>Prasophyllum innubum</i> D.L.Jones		
<i>Prasophyllum montanum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum petilum</i> D.L.Jones & R.J.Bates		
<i>Prasophyllum sphacelatum</i> D.L.Jones		
<i>Prasophyllum tadgellianum</i> R.S.Rogers		
<i>Prasophyllum venustum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum viriosum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum</i> species 'ACT 1'		late spring-summer flowering, scented aff. <i>patens</i> with extensively white petals and strongly recurved labellum with short green callus plate; montane forest between 700 and 900 m (Jones <i>et al.</i> 2008)
<i>Prasophyllum</i> species 'ACT 2'		late spring-summer flowering, scented aff. <i>patens</i> with white-margined brownish petals and more upright labellum with short green callus plate; montane forest between 700 and 900 m (Jones <i>et al.</i> 2008)
<i>Prasophyllum</i> species 'Smokers Gap'		large, late-summer flowering aff. <i>sphacelatum</i> ; alpine meadows in Namadgi National Park; mentioned in notes on <i>P. spachelatum</i> in Jones <i>et al.</i> (2008); possibly the same as <i>P.</i> species 'Nunniong'?
<i>Pterostylis aciculiformis</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis aestivalis</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis alpina</i> R.S.Rogers		
<i>Pterostylis aneba</i> D.L.Jones		
<i>Pterostylis atrans</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis bicolor</i> M.A.Clem. & D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis clivicola</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis coccinea</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis crassicaulis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis curta</i> R.Br.		
<i>Pterostylis cycnocephala</i> Fitzg.	<i>Hymenochilus</i>	
<i>Pterostylis decurva</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis falcata</i> R.S.Rogers		
<i>Pterostylis fischii</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis hamata</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	
<i>Pterostylis jonesii</i> G.N.Backh.	<i>Bunochilus</i>	originally described as <i>Bunochilus montanus</i> D.L. Jones
<i>Pterostylis laxa</i> Blackmore	<i>Diplodium</i>	

Australian Capital Territory Species	Alternate Genus	Notes
<i>Pterostylis monticola</i> D.L.Jones		
<i>Pterostylis multiflora</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis oreophila</i> Clemesha		
<i>Pterostylis pedunculata</i> R.Br.		
<i>Pterostylis reflexa</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis rubescens</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis squamata</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis truncata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis umbrina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis</i> species 'ACT'	<i>Bunochilus</i>	small, early flowering aff. <i>melagramma</i> with brown sepal tips to the sepals; shrubby open forest in Namadgi National Park (J. Egan & T. Wood pers. comm.)
<i>Pterostylis</i> species 'large curved flowers'	<i>Diplodium</i>	aff. <i>revoluta</i> with larger, more curved flowers; widespread in dry woodland and open forest through inland south-eastern Australia (Bishop 2000; Jeanes & Backhouse 2006)
<i>Spiranthes alticola</i> D.L.Jones		very difficult to distinguish from (and doubtfully distinct from) <i>Spiranthes australis</i>
<i>Thelymitra alpina</i> Jeanes		
<i>Thelymitra arenaria</i> Lindl.		
<i>Thelymitra brevifolia</i> Jeanes		
<i>Thelymitra carnea</i> R.Br.		
<i>Thelymitra cyanea</i> (Lindl.)		
<i>Thelymitra juncifolia</i> Lindl.		
<i>Thelymitra megcalyptra</i> Fitzg.		
<i>Thelymitra nuda</i> R.Br.		
<i>Thelymitra pauciflora</i> R.Br.		
<i>Thelymitra peniculata</i> Jeanes		
<i>Thelymitra rubra</i> Fitzg.		
<i>Thelymitra simulata</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra</i> species 'fine spots'		aff. <i>ixioides</i> with self-pollinating flowers with few very fine spots on dorsal sepal and petals and irregularly glandular post-anther lobe; damp mountain forest, swampy and damp grassy areas above 800 m (Jones <i>et al.</i> 2008)
<i>Thelymitra</i> species 'Gibraltar Falls'		no information available (Australian National Herbarium record)
<i>Thelymitra</i> species 'ornate flowers'		small self-pollinating aff. <i>simulata</i> with freely opening flowers with shortly hooded deeply notched post-anther lobe; mountain open forest above 750 m; resembles the New Zealand species <i>T. decora</i> (Jones <i>et al.</i> 2008)
<i>Thynniorchis huntianus</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	

**Total: 132 species**



Australian Capital Territory Hybrids	Name and Authority	Notes
<i>Caladenia cucullata</i> X <i>ustulata</i>		J. Egan & T. Wood pers. comm.
<i>Caladenia carnea</i> X <i>Cyanicula caerulea</i>		ACT Orchid Group blog
<i>Chiloglottis trapeziformis</i> X <i>valida</i>	<i>x</i> <i>pescottiana</i> R.S. Rogers	
<i>Diuris chrysiopsis</i> X <i>nigromontana</i>		J. Egan & T. Wood pers. comm.
<i>Diuris chrysiopsis</i> X <i>pardina</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Diuris monticola</i> X <i>semilunulata</i>		J. Egan & T. Wood pers. comm.
<i>Diuris nigromontana</i> X <i>sulphurea</i>		J. Egan & T. Wood pers. comm.
<i>Pterostylis atrans</i> X <i>decurva</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Thelymitra brevifolia</i> X <i>juncifolia</i>		Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra juncifolia</i> X <i>pauciflora</i>		Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra juncifolia</i> X <i>peniculata</i>		Jones <i>et al.</i> 2008

**Total: 11 hybrids**

New South Wales Species	Alternate Genus	Notes
<i>Acianthella amplexicaulis</i> (F.M.Bailey) D.L.Jones & M.A.Clem.	<i>Acianthus</i>	
<i>Acianthus apprimus</i> D.L.Jones		
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	
<i>Acianthus collinus</i> D.L.Jones		
<i>Acianthus exiguus</i> D.L.Jones		
<i>Acianthus exsertus</i> R.Br.		
<i>Acianthus fornicatus</i> R.Br.		
<i>Acianthus pusillus</i> D.L.Jones		
<i>Acianthus saxatilis</i> D.L.Jones & M.A.Clem.		
<i>Acianthus</i> species 'Gibraltar Range'		autumn flowering aff. <i>caudatus</i> with smaller flowers; wet forest in the Gibraltar Range (L. Copeland unpublished)
<i>Acianthus</i> species 'NSW Northern Tablelands'		autumn flowering aff. <i>fornicatus</i> with short dark sepal clubs and no teeth or lobes on lateral sepals; grassy open forest on the New England Tableland (Bishop 2000); records of <i>A. apprimus</i> from northern NSW are probably this taxon
<i>Acianthus</i> species 'Tathra'		tiny aff. <i>fornicatus</i> with labellum <4 mm long; coastal scrubs and heathy woodland on South Coast headlands (Bishop 2000)
<i>Adenochilus nortonii</i> Fitzg.		
<i>Arthrochilus prolixus</i> D.L.Jones		
<i>Bulbophyllum argyropus</i> (Endl.) Rchb.f.	<i>Adelopetalum</i>	also on Lord Howe Island
<i>Bulbophyllum bracteatum</i> Fitzg.	<i>Adelopetalum</i>	
<i>Bulbophyllum elisae</i> F.Muell.	<i>Adelopetalum</i>	
<i>Bulbophyllum exiguum</i> F.Muell.	<i>Adelopetalum</i>	
<i>Bulbophyllum globuliforme</i> Nicholls	<i>Oncophyllum</i>	
<i>Bulbophyllum lamingtonense</i> D.L.Jones	<i>Oxysepala</i>	
<i>Bulbophyllum minutissimum</i> F.Muell.	<i>Oncophyllum</i>	
<i>Bulbophyllum schillerianum</i> Rchb.f. subspecies <i>schillerianum</i>	<i>Oxysepala</i>	
<i>Bulbophyllum shepherdii</i> (F.Muell.) Rchb.f.	<i>Oxysepala</i>	
<i>Bulbophyllum weinthalii</i> R.S.Rogers subspecies <i>weinthalii</i>	<i>Spilorchis</i>	
<i>Burnettia cuneata</i> Lindl.		
<i>Caladenia aestiva</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia alata</i> R.Br.	<i>Petalochilus</i>	
<i>Caladenia alpina</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia amnicola</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia arenaria</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia atrovespa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia attenuata</i> (W.Brinsley) D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia branwhitei</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia leptoclavia</i>
<i>Caladenia cadyi</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia callitrophila</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia catenata</i> (Sm.) Druce	<i>Petalochilus</i>	
<i>Caladenia clarkiae</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia clavigera</i> A.Cunn. ex Lindl.	<i>Arachnorchis</i>	
<i>Caladenia cleistantha</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia concinna</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	
<i>Caladenia concolor</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia cucullata</i> Fitzg.	<i>Stegostyla</i>	
<i>Caladenia curtisepala</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia dimorpha</i> Fitzg.	<i>Stegostyla</i>	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia fitzgeraldii</i> Rupp	<i>Arachnorchis</i>	
<i>Caladenia flaccida</i> D.L.Jones	<i>Jonesiopsis</i>	
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia hildae</i> E.E.Pescott & Nicholls	<i>Stegostyla</i>	
<i>Caladenia hillmanii</i> D.L.Jones	<i>Petalochilus</i>	

New South Wales Species	Alternate Genus	Notes
<i>Caladenia leptoclavia</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia mentiensi</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia minor</i> Hook.f.	<i>Petalochilus</i>	a taxon of highly uncertain identity; Australian records from SA and NSW but generally considered to be endemic to New Zealand
<i>Caladenia montana</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia moschata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia oreophila</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia orestes</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia osmera</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	distinguishable from <i>Caladenia montana</i> only by altitude
<i>Caladenia peisleyi</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia phaeoclavia</i> D.L.Jones	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia parva</i>
<i>Caladenia picta</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia porphyrea</i> D.L.Jones	<i>Petalochilus</i>	doubtfully distinguishable from <i>Caladenia quadrifaria</i>
<i>Caladenia praecox</i> Nicholls	<i>Stegostyla</i>	
<i>Caladenia pusilla</i> W.M.Curtis	<i>Petalochilus</i>	
<i>Caladenia quadrifaria</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia rigens</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia rileyi</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia rosella</i> G.W.Carr	<i>Arachnorchis</i>	almost certainly extinct in NSW, if it ever occurred there
<i>Caladenia stellata</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia subtilis</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia tessellata</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia testacea</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia transitoria</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia ustulata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia whiteheadii</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia</i> species 'central NSW'	<i>Petalochilus</i>	small, self-pollinating aff. <i>vulgaris</i> with colourful flowers with narrow petals and sepals with rounded tips; woodland from Cowra to Armidale (Bishop 2000)
<i>Caladenia</i> species 'crowded calli'	<i>Arachnorchis</i>	aff. <i>concinna</i> with larger flowers with larger, more numerous crowded labellum calli extending almost to the apex, and short sepal clubs; shrubby woodland between Temora and Rankin Springs (Bower 2008; Backhouse 2011)
<i>Caladenia</i> species 'Dorrigo'	<i>Stegostyla</i>	aff. <i>transitoria</i> with internally white sepals and petals; montane scrub on rainforest margin near Dorrigo (Copeland unpubl.; Backhouse 2011) <i>dorrigoensis</i> ms
<i>Caladenia</i> species 'inland NSW'	<i>Arachnorchis</i>	aff. <i>atrovespa</i> with smaller, stiffer flowers; has a different wasp pollinator to <i>atrovespa</i> ; dry woodland on inland plains between Rankin Springs, West Wyalong and Lockhart (Colin Bower personal communication; Backhouse 2011)
<i>Caladenia</i> species 'Orange'	<i>Arachnorchis</i>	aff. <i>patersonii</i> with large pinkish mauve flowers with darker calli and teeth and brownish purple tails; hilly open forest near Orange and the Blue Mountains; last seen in the late 1980s and probably extinct (Bishop 2000)
<i>Caladenia</i> species 'pointed flowers'	<i>Petalochilus</i>	aff. <i>mentiensi</i> with sharply pointed sepals and petals and incurved dorsal sepal; hilly dry woodland near Gungahlin (J. Egan & T. Wood personal communication)
<i>Caladenia</i> species 'Richmond River'	<i>Arachnorchis</i>	aff. <i>patersonii</i> with creamy yellow flowers with long dark glandular tails; heathy open forest near the Richmond River; collected in the late 1800s and not seen since, probably extinct (Bishop 2000)
<i>Caladenia</i> species 'Woronora Plateau'	<i>Petalochilus</i>	aff. <i>alata</i> with calli coalescing into wrinkled ridges and backward pointing large teeth at mid-lob base; heathy woodland on Woronora Plateau (Bishop 2000)
<i>Calanthe australasica</i> D.L.Jones & M.A.Clem.		
<i>Caleana major</i> R.Br.		
<i>Calochilus campestris</i> R.Br.		
<i>Calochilus gracillimus</i> Rupp		
<i>Calochilus grandiflorus</i> (Benth.) Domin		
<i>Calochilus kalaru</i> D.L.Jones		
<i>Calochilus montanus</i> D.L.Jones		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus praeltus</i> D.L.Jones		

New South Wales Species	Alternate Genus	Notes
<i>Calochilus pulchellus</i> D.L.Jones		
<i>Calochilus robertsonii</i> Benth.		
<i>Calochilus russeus</i> D.L.Jones		very similar to, and possibly just a montane form of, <i>Calochilus therophilus</i>
<i>Calochilus sandrae</i> D.L.Jones		
<i>Calochilus saprophyticus</i> R.S.Rogers		
<i>Calochilus therophilus</i> D.L.Jones		
<i>Calochilus</i> species 'Coaldale'		aff. <i>campestris</i> with short flowers with shorter basal plates and broader column wings; heathy woodland on rocky hills near Coaldale (L. Copeland unpublished)
<i>Calochilus</i> species 'Gibraltar Range'		aff. <i>campestris</i> with slim reddish stem with several longish slender bracts held well away from stem and small, narrow flowers with long narrow rostrum; heathy woodland in Gibraltar Range
<i>Calochilus</i> species 'New England'		aff. <i>robertsonii</i> with small flowers with reddish eye-like column glands and a sparsely bristled labellum with short apical bristles and a visible naked tail; basalt grasslands and grassy woodlands between Ebor and Ben Lomond
<i>Cestichis coelognoides</i> (F.Muell.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis reflexa</i> (R.Br.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis swenstonii</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cheirostylis notialis</i> D.L.Jones		
<i>Chiloglottis anatriceps</i> D.L.Jones		
<i>Chiloglottis chlorantha</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis cornuta</i> Hook.f.	<i>Simpliglottis</i>	
<i>Chiloglottis diphylla</i> R.Br.		
<i>Chiloglottis formicifera</i> Fitzg.	<i>Myrmechila</i>	
<i>Chiloglottis palachila</i> D.L.Jones		
<i>Chiloglottis platyptera</i> D.L.Jones	<i>Myrmechila</i>	
<i>Chiloglottis pluricallata</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis reflexa</i> (Labill.) Druce		
<i>Chiloglottis seminuda</i> D.L.Jones		
<i>Chiloglottis sphaerula</i> D.L.Jones		
<i>Chiloglottis sphyrnoides</i> D.L.Jones		
<i>Chiloglottis sylvestris</i> D.L.Jones & M.A.Clem.		
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmechila</i>	
<i>Chiloglottis trilabra</i> Fitzg.		
<i>Chiloglottis turfosa</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis valida</i> D.L.Jones	<i>Simpliglottis</i>	'true' <i>valida</i> occurs in the mountains above 500 m but is morphologically indistinguishable from two related cryptic species (Bower 2007; see sp. 'Falls Creek' and sp. 'lowlands'), distribution overlaps with sp. 'Falls Creek'
<i>Chiloglottis</i> species 'Bald Rock'	<i>Myrmechila</i>	small, spring flowering aff. <i>formicefera</i> with recurved labellum apex, calli extending almost to apex and narrower column wings; grassy open forest on rich loams; border region in Girraween and Bald Rock National Parks (Bishop 2000)
<i>Chiloglottis</i> species 'Barrington Tops'	<i>Simpliglottis</i>	aff. <i>pluricallata</i> with more numerous calli and different wasp pollinator; heathy swamps on Barrington Tops (Bishop 2000)
<i>Chiloglottis</i> species 'Bathurst'		small aff. <i>palachila</i> with yellow clubs on lateral sepals and much smaller calli mass; also has a different wasp pollinator; shrubby open forest and woodland near Bathurst (Bishop 2000)
<i>Chiloglottis</i> species 'Ebor'	<i>Simpliglottis</i>	aff. <i>pluricallata</i> with more numerous calli and different wasp pollinator; wet forest and sphagnum swamps in New England National Park around Point Lookout (Bishop 2000)
<i>Chiloglottis</i> species 'highlands'	<i>Simpliglottis</i>	aff. <i>valida</i> with slightly smaller labellum with fewer calli and different wasp pollinator; montane damp forest and woodland above 800 m (Bower 2007; Jones <i>et al.</i> 2008); indistinguishable from <i>C. valida</i>
<i>Chiloglottis</i> species 'lowlands'	<i>Simpliglottis</i>	aff. <i>valida</i> indistinguishable from <i>valida</i> but with a different wasp pollinator (Bower 2007); widespread in damp shady sites in south-eastern Aust. below 1,000 m, mostly at low altitudes (Jeanes & Backhouse 2006)
<i>Corunastylis acuminata</i> (R.S.Rogers) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis anomala</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis anthracina</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis apostasioides</i> Fitzg.	<i>Genoplesium</i>	
<i>Corunastylis archeri</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	'true' <i>archeri</i> almost certainly does not occur in NSW & Qld, with these plants probably comprising a complex of several different, un-named species
<i>Corunastylis arrecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis bishopii</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	

New South Wales Species	Alternate Genus	Notes
<i>Corunastylis carecta</i> D.L.Jones & L.M.Copel.	<i>Genoplesium</i>	
<i>Corunastylis citriodora</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis clivicola</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis cornuta</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis densa</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis despectans</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis eriochila</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis filiformis</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis fimbriata</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis formosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis insignis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis laminata</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis littoralis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morrisii</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nuda</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis oligantha</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis ostrina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis plumosa</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis pumila</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis rhyolitica</i> (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis rufa</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis ruppii</i> (R.S.Rogers) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis sagittifera</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis simulans</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis stephensonii</i> D.L.Jones	<i>Genoplesium</i>	
<i>Corunastylis superba</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis systema</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis tenella</i> D.L.Jones & L.M.Copel.	<i>Genoplesium</i>	
<i>Corunastylis trifida</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis turfosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis vernalis</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis woollsii</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis</i> species 'Charmhaven'	<i>Genoplesium</i>	aff. <i>archeri</i> with fewer larger flowers in a shorter inflorescence; heathy open forest around Charmhaven
<i>Corunastylis</i> species 'dark flowers'	<i>Genoplesium</i>	aff. <i>rufa</i> with pointed dark reddish black labellum; widespread in dry woodland across inland south-eastern Australia (=C. sp. 'inland' in Backhouse <i>et al.</i> 2016; C. sp. 'dark midge' in Bates 2011)
<i>Corunastylis</i> species 'Mount Kaputar'	<i>Genoplesium</i>	summer-flowering aff. <i>arrecta</i> with numerous crowded flowers with upswept lateral sepals and hairless petals; heathy woodland and rocky outcrops in the Nandewar Range (Bishop 2000)
<i>Corunastylis</i> species 'granite ranges'	<i>Genoplesium</i>	spring-summer flowering, robust aff. <i>archeri</i> with more crowded flowers with strongly upcurved lateral sepals; open forest on granite ranges of north-eastern NSW (L. Copeland unpublished) and south-eastern Qld
<i>Corybas aconitiflorus</i> Salisb.		
<i>Corybas barbarae</i> D.L.Jones		also on Lord Howe Island
<i>Corybas diemenicus</i> (Lindl.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas dowlingii</i> D.L.Jones		doubtfully distinct from, and probably just a form of, <i>Corybas aconitiflorus</i> (Mark Clements CSIRO personal communication)
<i>Corybas fimbriatus</i> (R.Br.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	
<i>Corybas grumulus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas hispidus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	
<i>Corybas pruinosis</i> (A.Cunn.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas undulatus</i> (A.Cunn.) Rupp	<i>Corysanthes</i>	
<i>Corybas unguiculatus</i> (R.Br.) Rchb.f.	<i>Anzybas</i>	
<i>Corybas</i> species 'Gibraltar Range'	<i>Anzybas</i>	aff. <i>unguiculatus</i> with shorter beak on labellum orifice and with calli on the main and lateral veins inside the orifice (D.L. Jones personal communication comm.); damp montane forest in the Gibraltar Range



New South Wales Species	Alternate Genus	Notes
<i>Cryptostylis erecta</i> R.Br.		
<i>Cryptostylis hunteriana</i> Nicholls		
<i>Cryptostylis leptochila</i> Benth.		
<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		
<i>Cyanicula caerulea</i> (R.Br.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cymbidium canaliculatum</i> R.Br.		many forms and varieties have been named (eg. var. <i>sparksii</i> ) although none have any formal taxonomic recognition
<i>Cymbidium madidum</i> Lindl.		
<i>Cymbidium suave</i> R.Br.		
<i>Cyrtostylis reniformi</i> R.Br.		
<i>Danhatchia australis</i> (Hatch) Garay & Christenson	<i>Yoania</i>	
<i>Dendrobium aemulum</i> R.Br.	<i>Tropilis</i>	
( <i>Dendrobium</i> ) <i>angustum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
( <i>Dendrobium</i> ) <i>banksii</i> (D.L.Jones & M.A.Clem.)	<i>Dockrillia</i>	described in <i>Dockrillia</i> , new combination in <i>Dendrobium</i> not yet made; sometimes included in <i>Dendrobium striolatum</i>
<i>Dendrobium bowmanii</i> Benth.	<i>Dockrillia</i>	
<i>Dendrobium cucumerinum</i> MacLeay ex Lindl.	<i>Dockrillia</i>	
<i>Dendrobium dolichophyllum</i> D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	
( <i>Dendrobium</i> ) <i>eburneum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium fairfaxii</i> F.Muell. ex Fitzg.	<i>Dockrillia</i>	
<i>Dendrobium falcorostrum</i> Fitzg.	<i>Thelychiton</i>	
<i>Dendrobium gracilicaule</i> F.Muell.	<i>Thelychiton</i>	
<i>Dendrobium howeanum</i> (Maiden) D.L.Jones	<i>Thelychiton</i>	Lord Howe Island only; variously considered a variety of <i>Dendrobium gracilicaule</i> and <i>Dendrobium macropus</i>
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>kingianum</i>	<i>Thelychiton</i>	several varieties have been named although these are recognised mainly in cultivation and not taxonomically; one variety is recognised here
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>kingianum</i> variety <i>pulcherrimum</i> Rupp	<i>Thelychiton</i>	considered a full species <i>D. pulcherrimum</i> by Jones (2006)
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>huntianum</i> Rupp	<i>Dockrillia</i>	
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>linguiforme</i>	<i>Dockrillia</i>	
<i>Dendrobium melaleucaphilum</i> M.A.Clem. & D.L.Jones	<i>Tetrabaculum</i>	alternative name <i>Dendrobium tetragonum</i> subspecies <i>tetragonum</i> variety <i>melaleucaphilum</i>
<i>Dendrobium monophyllum</i> F.Muell.	<i>Australorchis</i>	
<i>Dendrobium moorei</i> F.Muell.	<i>Thelychiton</i>	Lord Howe Island only
<i>Dendrobium mortii</i> F.Muell.	<i>Dockrillia</i>	
<i>Dendrobium pugioniforme</i> A.Cunn.	<i>Dockrillia</i>	
( <i>Dendrobium</i> ) <i>radiata</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium schneiderae</i> F.M.Bailey variety <i>schneiderae</i>	<i>Australorchis</i>	
<i>Dendrobium schoeninum</i> Lindl.	<i>Dockrillia</i>	
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>hillii</i> Masters	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) tarberi</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>speciosum</i>	<i>Thelychiton</i>	alternative name (in part) <i>Dendrobium (Thelychiton) epiphyticus</i>
<i>Dendrobium striolatum</i> Rchb.f. Rauschert subspecies <i>striolatum</i>	<i>Dockrillia</i>	
<i>Dendrobium teretifolium</i> R.Br.	<i>Dockrillia</i>	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>tetragonum</i> variety <i>tetragonum</i>	<i>Tetrabaculum</i>	
<i>Dipodium atropurpureum</i> D.L.Jones		
<i>Dipodium hamiltonianum</i> F.M.Bailey		
<i>Dipodium pulchellum</i> D.L.Jones & M.A.Clem		
<i>Dipodium punctatum</i> (Sm.) R.Br.		
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		
<i>Dipodium variegatum</i> M.A.Clem. & D.L.Jones		
<i>Diuris abbreviata</i> Benth.		
<i>Diuris aequalis</i> F.Muell. ex Fitzg.		
<i>Diuris alba</i> R.Br.		
<i>Diuris arenaria</i> D.L.Jones		
<i>Diuris aurea</i> Sm.		

New South Wales Species	Alternate Genus	Notes
<i>Diuris behrii</i> Schldl.		
<i>Diuris bracteata</i> Fitzg.		
<i>Diuris brevissima</i> Fitzg. ex Nicholls		included on PlantNET (NSW Flora Online) but status doubtful, usually included in <i>Diuris maculata</i>
<i>Diuris byronensis</i> D.L.Jones		
<i>Diuris callitrophila</i> D.L.Jones		
<i>Diuris chrysantha</i> D.L.Jones & M.A.Clem.		
<i>Diuris chryseopsis</i> D.L.Jones		
<i>Diuris cuneilabris</i> Rupp		included on Atlas of Living Australia but status doubtful, usually included in <i>Diuris platichila</i>
<i>Diuris curta</i> D.L.Jones		
<i>Diuris dendrobioides</i> Fitzg.		
<i>Diuris disposita</i> D.L.Jones		
<i>Diuris eborensis</i> D.L.Jones		
<i>Diuris flavescens</i> D.L.Jones		
<i>Diuris fucosa</i> D.L.Jones		
<i>Diuris goonooensis</i> Rupp		
<i>Diuris maculata</i> Sm.		
<i>Diuris minor</i> (Benth.) D.L.Jones & M.A.Clem.		
<i>Diuris monticola</i> D.L.Jones		
<i>Diuris nigromontana</i> D.L.Jones		
<i>Diuris ochroma</i> D.L.Jones		
<i>Diuris orientis</i> D.L.Jones		
<i>Diuris pardina</i> Lindl.		
<i>Diuris parvipetala</i> (Dockrill) D.L.Jones & M.A.Clem.		
<i>Diuris pedunculata</i> R.Br.		
<i>Diuris platichila</i> Fitzg.		
<i>Diuris praecox</i> D.L.Jones		
<i>Diuris punctata</i> Sm. variety <i>punctata</i>		
<i>Diuris punctata</i> Sm. variety <i>sulfurea</i> Rupp		most likely a hybrid between <i>D. punctata</i> and a yellow-flowered <i>Diuris</i> species
<i>Diuris secundiflora</i> Fitzg.		
<i>Diuris semilunulata</i> Messmer		
<i>Diuris striata</i> Rupp		taxon of uncertain identity, possibly a form of <i>Diuris dendrobioides</i> with short lateral sepals
<i>Diuris subalpina</i> D.L.Jones		
<i>Diuris sulphurea</i> R.Br.		
<i>Diuris systema</i> D.L.Jones & L.M.Copel.		
<i>Diuris tricolor</i> Fitzg.		
<i>Diuris unica</i> D.L.Jones		
<i>Diuris venosa</i> Rupp		
<i>Diuris</i> species 'Clarencetown'		aff. <i>alba</i> with smaller more uniformly coloured flowers with erect dorsal sepals; shrubby open forest near Clarencetown (Bishop 2000)
<i>Diuris</i> species 'Colo River'		aff. <i>punctata</i> with smaller flowers with short lateral sepals, reflexed petals and creamy yellow between labellum callus ridges; swamp margins in heath, heathy & grassy woodland Mellong Range to Hunter Valley (Bishop 2000)
<i>Diuris</i> species 'Ebor'		tall aff. <i>dendrobioides</i> with larger, darker, more uniformly coloured flowers with more erect dorsal sepals; montane grassland and grassy woodland near Ebor (Copeland 2005)
<i>Diuris</i> species 'Grafton'		early spring-flowering aff. <i>dendrobioides</i> with smaller pale flowers with short lateral sepals; grassy open forest near Grafton (L. Copeland unpublished)
<i>Diuris</i> species 'Guyra'		small aff. <i>dendrobioides</i> with mainly white flowers; montane grasslands above 1,200 m on New England tableland (Copeland 2005)
<i>Diuris</i> species 'Hunter Valley'		very small aff. <i>dendrobioides</i> with rounded petals not tapering into the petiole; grassy open forest in the central Hunter Valley (Bishop 2000)
<i>Diuris</i> species 'Monaro'		small late spring-early summer flowering aff. <i>dendrobioides</i> with indistinct blotches and yellow between callus ridges; native grasslands on the Monaro tableland (Bishop 2000)
<i>Diuris</i> species 'New England escarpment'		spring flowering aff. <i>alba</i> with colourful flowers with broad, reflexed petals and yellow tubers; heathy woodland and open forest on granite hills on the New England Tableland (Bishop 2000)
<i>Diuris</i> species 'New England swamps'		summer-flowering aff. <i>alba</i> with recurved petals; swamps on New England Tableland (Bishop 2000)

New South Wales Species	Alternate Genus	Notes
<i>Diuris</i> species 'NSW North Coast'		short, robust aff. <i>punctata</i> with large paler flowers with broad dorsal sepal, petals & labellum, white/cream between labellum callus ridges; heaths and grassy woodland on coastal headlands, Kempsey to Yamba (Bishop 2000)
<i>Diuris</i> species 'Tenterfield/Stanthorpe'		aff. <i>punctata</i> with pale mottled flowers with relatively short lateral sepals and white between the labellum callus ridges; open forest in the granite ranges around Tenterfield NSW and Stanthorpe Qld
<i>Empusa habenarina</i> (F.Muell.) Benth.	<i>Liparis</i>	
<i>Epipogium roseum</i> (D.Don) Lindl.		
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus magenteus</i> D.L.Jones		
<i>Eriochilus petricola</i> D.L.Jones & M.A.Clem.		
<i>Eriochilus</i> species 'smooth leaf'		aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<i>Erythrorchis cassythoides</i> (A.Cunn. ex Lindl.) Garay		
<i>Gastrodia entomogama</i> D.L.Jones		
<i>Gastrodia procera</i> G.W.Carr		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Gastrodia surcula</i> D.L.Jones		
<i>Genoplesium baueri</i> R.Br.		
<i>Geodorum densiflorum</i> (Lam.) Schltr.		also called <i>Geodorum terrestre</i> (L.) Garay
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Glossodia minor</i> R.Br.	<i>Caladenia</i>	alternative name <i>Caladenia minorata</i> M.A. Clem.
<i>Lyperanthus suaveolens</i> R.Br.		
<i>Microtis angusii</i> D.L.Jones		
<i>Microtis oblonga</i> R.S.Rogers		
<i>Microtis parviflora</i> R.Br.		also on Lord Howe Island
<i>Microtis rara</i> R.Br.		identity and distribution uncertain, thought to be only NSW & Qld (Jones 2006), although also claimed to be in Sth Aust (Bates 2011)
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		also on Lord Howe Island
<i>Microtis</i> species 'alpine'		aff. <i>unifolia</i> with fewer smaller spaced flowers with a smaller labellum; wet mountain meadows and woodland (Jones <i>et al.</i> 2008)
<i>Microtis</i> species 'fragrant'		aff. <i>arenaria</i> with sparser, scented flowers with a longer labellum with a raised central ridge; widespread in many habitats in south-eastern Australia; <i>fragrans</i> ms (Bates 2011)
<i>Oberonia complanata</i> (A.Cunn.) M.A.Clem. & D.L.Jones		
<i>Oberonia palmicola</i> F.Muell.		also included in <i>Oberonia titania</i>
<i>Orthoceras strictum</i> R.Br.		
<i>Papillilabium beckeri</i> (F.Muell. ex Benth.) Dockrill		
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Peristeranthus hillii</i> (F.Muell.) T.E.Hunt		
<i>Phaius australis</i> F.Muell.		
<i>Pheladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	
<i>Plectorrhiza erecta</i> (Fitzg.) Dockrill		Lord Howe Island only
<i>Plectorrhiza tridentata</i> (Lindl.) Dockrill		
<i>Prasophyllum affine</i> Lindl.		
<i>Prasophyllum alpestre</i> D.L.Jones		
<i>Prasophyllum appendiculatum</i> Nicholls		
<i>Prasophyllum australe</i> R.Br.		
<i>Prasophyllum bagoense</i> D.L.Jones		
<i>Prasophyllum beatrix</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum brevilabre</i> (Lindl.) Hook.f.		
<i>Prasophyllum campestre</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum canaliculatum</i> D.L.Jones		
<i>Prasophyllum candidum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum caricetum</i> D.L.Jones		
<i>Prasophyllum caudiculum</i> D.L.Jones		
<i>Prasophyllum dossenum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum exilis</i> D.L.Jones & R.J.Bates		

<i>Prasophyllum flavum</i> R.Br.	
<i>Prasophyllum fuscum</i> R.Br.	
<i>Prasophyllum helophilum</i> D.L.Jones & D.T.Rouse	
<i>Prasophyllum innubum</i> D.L.Jones	
<i>Prasophyllum keltonii</i> D.L.Jones	
<i>Prasophyllum montanum</i> R.J.Bates & D.L.Jones	
<i>Prasophyllum pallens</i> D.L.Jones	
<i>Prasophyllum patens</i> R.Br.	
<i>Prasophyllum petilum</i> D.L.Jones & R.J.Bates	
<i>Prasophyllum retroflexum</i> D.L.Jones	
<i>Prasophyllum rogersii</i> Rupp	
<i>Prasophyllum solstitium</i> D.L.Jones	
<i>Prasophyllum sphacelatum</i> D.L.Jones	
<i>Prasophyllum striatum</i> R.Br.	<i>Mecopodum</i>
<i>Prasophyllum sylvestre</i> R.J.Bates & D.L.Jones	
<i>Prasophyllum tadgellianum</i> R.S.Rogers	
<i>Prasophyllum venustum</i> D.L.Jones & D.T.Rouse	
<i>Prasophyllum viriosum</i> D.L.Jones & D.T.Rouse	
<i>Prasophyllum wilkinsoniorum</i> D.L.Jones	
<i>Prasophyllum</i> species 'Barrington Tops'	aff. <i>fuscum</i> from Barrington Tops; known only from a single collection, no other information available
<i>Prasophyllum</i> species 'black soil plains'	early spring-flowering aff. <i>odoratum</i> with large flowers with relatively short lateral sepals; black soil plains near Inverell (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'Boyd Plateau'	tall aff. <i>alpestre</i> with colourful narrowly opening flowers with smaller, less ruffled labellum; wet heaths on Boyd Plateau and Mt Werong (Bishop 2000)
<i>Prasophyllum</i> species 'Delegate'	late spring/early summer flowering, small aff. <i>petilum</i> ; grasslands on the Monaro Tablelands near Delegate
<i>Prasophyllum</i> species 'desert'	robust aff. <i>odoratum</i> with long narrow petals; semi-arid sandy and rocky grassy shrublands, often with <i>Triodia</i> ; widespread across central SA and far western NSW (Bates 2011); <i>eremaemum</i> ms
<i>Prasophyllum</i> species 'eastern highlands'	summer flowering aff. <i>odoratum</i> with tall, moderately crowded spike of large flowers; subalpine woodland of eastern Vic and southern NSW above 800 m (=aff. <i>odoratum</i> L Rouse 2002)
<i>Prasophyllum</i> species 'late summer'	late summer-flowering aff. <i>dossenum</i> with a prominent kink in the lateral sepals; montane grassland and grassy woodland on the Barrington Tops (L. Copeland unpublished)
<i>Prasophyllum</i> species 'Mount Kaputar'	aff. <i>odoratum</i> with green and white flowers with long, thin petals and lateral sepals; open forest in Nandawar Range (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'mountain bogs'	relatively tall, late summer-flowering aff. <i>alpestre</i> with narrow petals and sharply curved labellum; mountain swamps above 700 m, Tamboritha (Vic) to Adaminaby (NSW) (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'New England batholith'	late spring-summer flowering aff. <i>odoratum</i> with crowded flowers with large, deeply ruffled labellum; open forest in the granite ranges above 900 m on New England Tableland NSW (L. Copeland unpubl.) & near Stanthorpe Qld
<i>Prasophyllum</i> species 'northern montane'	small, summer flowering aff. <i>odoratum</i> with widely spaced brown and white flowers; moist grassy woodland and open forest above 1,000, Barrington Tops, Ebor and Dorrigo (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'Nowra'	small aff. <i>appendiculatum</i> with more brownish flowers with shorter, broader petals and sepals and a broader, flatter labellum; heathy woodland on the Shoalhaven escarpment (Stephenson 2011)
<i>Prasophyllum</i> species 'Old Bar'	aff. <i>fuscum</i> with greenish brown flowers with white labellum lamina and short broad lateral sepals; coastal heath at Old Bar (Copeland unpubl.)
<i>Prasophyllum</i> species 'Pilliga Scrub'	robust aff. <i>odoratum</i> with pale green, white and brown flowers with long, narrow, spreading lateral sepals, narrow petals and long labellum with short callus plate and broad lamina; dry woodland in Pilliga Scrub (L. Copeland unpubl.)
<i>Prasophyllum</i> species 'Riverina grasslands'	tall, early flowering aff. <i>suaveolens</i> with larger flowers with a long narrowly triangular labellum; Riverina plains grasslands from Hunter (Vic) to Moama (NSW) (Jeanes & Backhouse 2006; =aff. <i>occidentale</i> D Rouse 2002)
<i>Prasophyllum</i> species 'Teralba'	aff. <i>exilis</i> with ovary held away from the stem; coastal <i>Angophora</i> forest near Teralba (NSW Herbarium information)
<i>Pseudovanilla foliata</i> (F.Muell.) Garay	
<i>Pterostylis abrupta</i> D.L.Jones	<i>Diplodium</i>
<i>Pterostylis aciculiformis</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>

New South Wales Species	Alternate Genus	Notes
<i>Pterostylis acuminata</i> R.Br.		
<i>Pterostylis aestiva</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis alpina</i> R.S.Rogers		
<i>Pterostylis alveata</i> Garnet	<i>Diplodium</i>	
<i>Pterostylis amabilis</i> (D.L.Jones & L.M.Copel.) D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis aneba</i> D.L.Jones		
<i>Pterostylis antennifera</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis atrans</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis baptistii</i> Fitzg.		
<i>Pterostylis barringtonensis</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis bicolor</i> M.A.Clem. & D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis bicornis</i> D.L.Jones & M.A.Clem.	<i>Petrorchis</i>	
<i>Pterostylis boormanii</i> Rupp	<i>Oligochaetochilus</i>	
<i>Pterostylis calceolus</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis chaetophora</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis chocolatina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis clavigera</i> Fitzg.	<i>Linguella</i>	
<i>Pterostylis clivicola</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis cobarensis</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis coccinea</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis collina</i> (Rupp) M.A.Clem. & D.L.Jones	<i>Taurantha</i>	
<i>Pterostylis concinna</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis crassicaulis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis crebra</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis curta</i> R.Br.		also on Lord Howe Island
<i>Pterostylis cycnocephala</i> Fitzg.	<i>Hymenochilus</i>	
<i>Pterostylis daintreeana</i> Benth.	<i>Pharochilum</i>	also spelt <i>daintreana</i>
<i>Pterostylis decurva</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis elegans</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis erecta</i> T.E.Hunt		
<i>Pterostylis falcata</i> R.S.Rogers		
<i>Pterostylis fischii</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis foliata</i> Hook.f.		
<i>Pterostylis furva</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis gibbosa</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis glyphida</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis grandiflora</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis hamata</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	
<i>Pterostylis hians</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis hildae</i> Nicholls		
<i>Pterostylis hispidula</i> Fitzg.		
<i>Pterostylis jonesii</i> G.N.Backh.	<i>Bunochilus</i>	originally described as <i>Bunochilus montanus</i> D.L. Jones
<i>Pterostylis laxa</i> Blackmore	<i>Diplodium</i>	
<i>Pterostylis lineata</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis lingua</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis littorale</i> D.L.Jones	<i>Plumatichilus</i>	
<i>Pterostylis loganii</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis longicurva</i> Rupp	<i>Diplodium</i>	
<i>Pterostylis longifolia</i> R.Br.	<i>Bunochilus</i>	
<i>Pterostylis longipetala</i> Rupp	<i>Diplodium</i>	
<i>Pterostylis macrosepala</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis major</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis maxima</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis melagramma</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis metcalfei</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis mitchellii</i> Lindl.	<i>Oligochaetochilus</i>	
<i>Pterostylis monticola</i> D.L.Jones		
<i>Pterostylis multiflora</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	

New South Wales Species	Alternate Genus	Notes
<i>Pterostylis nana</i> R.Br.	<i>Linguella</i>	
<i>Pterostylis nigricans</i> D.L.Jones & M.A.Clem	<i>Speculantha</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis oblonga</i> D.L.Jones		
<i>Pterostylis obtusa</i> R.Br.	<i>Diplodium</i>	also on Lord Howe Island
<i>Pterostylis ophioglossa</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis oreophila</i> Clemesha		
<i>Pterostylis parca</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis parviflora</i> R.Br.	<i>Speculantha</i>	
<i>Pterostylis pedina</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	
<i>Pterostylis pedoglossa</i> Fitzg.	<i>Crangonorchis</i>	
<i>Pterostylis pedunculata</i> R.Br.		also on Lord Howe Island
<i>Pterostylis petrosa</i> D.L.Jones & M.A.Clem	<i>Oligochaetochilus</i>	
<i>Pterostylis plumosa</i> Cady	<i>Plumatichilos</i>	
<i>Pterostylis praetermissa</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis pulchella</i> Messmer	<i>Diplodium</i>	
<i>Pterostylis pusilla</i> R.S.Rogers	<i>Oligochaetochilus</i>	
<i>Pterostylis reflexa</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis revoluta</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis riparia</i> D.L.Jones		
<i>Pterostylis rubescens</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis rufa</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis russellii</i> T.E.Hunt	<i>Diplodium</i>	
<i>Pterostylis saxicola</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis setifera</i> M.A.Clem., Matthias & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis smaragdina</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis squamata</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis stenosepala</i> (D.L.Jones)	<i>Bunochilus</i>	
<i>Pterostylis striata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis subtilis</i> D.L.Jones		
<i>Pterostylis tasmanica</i> D.L.Jones	<i>Plumatichilos</i>	
<i>Pterostylis tenuis</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis torquata</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis truncata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis tunstallii</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis uliginosa</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis umbrina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis ventricosa</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis vernalis</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis woollsii</i> Fitzg.	<i>Oligochaetochilus</i>	
<i>Pterostylis</i> species 'Abercrombie Caves'	<i>Oligochaetochilus</i>	large aff. <i>aciculiformis</i> with crowded flowers on short pedicels; grassy woodland in Abercrombie Caves region southern tablelands (Bishop 2000)
<i>Pterostylis</i> species 'Albury'	<i>Bunochilus</i>	aff. <i>tunstallii</i> with strongly striped flowers; woodland on rocky hills near Albury (Jones & Clements 2006)
<i>Pterostylis</i> species 'Angourie'	<i>Oligochaetochilus</i>	aff. <i>woollsii</i> with strongly recurved dorsal sepal tail, lateral sepal tails curving forwards then drooping and recurved basal lobe on labellum; coast Banksia & tea-tree on moist sandy soils (Bishop 2000)
<i>Pterostylis</i> species 'Barrington Tops'	<i>Hymenochilus</i>	stout aff. <i>cynocephala</i> with larger more crowded flowers with a slender stalked labellum appendage; moist grassland around swamps on Barrington Tops (Bishop 2000)
<i>Pterostylis</i> species 'Buccarumbi'	<i>Diplodium</i>	aff. <i>revoluta</i> with very long, thin flowers with strongly projecting sinus; open forest around Buccarumbi (L. Copeland unpubl.)
<i>Pterostylis</i> species 'canegrass'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with paler flowers with a more sparsely bristled labellum; grows amongst <i>Triodia</i> on red sand dunes on sandy plains between Renmark and Olary (Bates 2011)
<i>Pterostylis</i> species 'Coaldale'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with paler flowers with narrower lateral sepals with more sparsely bristled outer margins and a broader labellum; open forest on sandy soils over sandstone near Coaldale (L. Copeland unpubl.)
<i>Pterostylis</i> species 'Coonabarabran'	<i>Oligochaetochilus</i>	aff. <i>praetermissa</i> with vertical, divergent shallowly concave shortly pointed lateral sepals wider than the galea, labellum with ~12 pairs of marginal hairs; <i>Callitris</i> woodland from Coonabarabran to Warialda (Bishop 2000)



New South Wales Species	Alternate Genus	Notes
<i>Pterostylis</i> species 'Dorrigo'	<i>Speculantha</i>	short, small, late spring-early summer flowering aff. <i>rubescens</i> with plump, extensively brown flowers; wet scrub and rainforest margins near Dorrigo (L. Copeland unpublished) <i>rubiginosa</i> ms
<i>Pterostylis</i> species 'East Kangaloon'	<i>Bunochilus</i>	aff. <i>smaragdina</i> with a narrow elongated labellum; open forest at East Kangaloon (Jones & Clements 2006)
<i>Pterostylis</i> species 'Ebor Falls'	<i>Diplodium</i>	aff. <i>elegans</i> with very small flowers; single location in damp open forest at Ebor (L. Copeland unpublished) ( <i>micranthum</i> ms)
<i>Pterostylis</i> species 'Goulburn'	<i>Diplodium</i>	small aff. <i>revoluta</i> with a very obtuse sinus; hilly woodland and open forest north of Goulburn (Bishop 2000)
<i>Pterostylis</i> species 'Griffith'	<i>Oligochaetochilus</i>	small green to brownish aff. <i>biseta</i> with oblong angular labellum virtually lacking a basal constriction; mallee and dry woodland north of Griffith (Bishop 2000)
<i>Pterostylis</i> species 'hooked sepals'	<i>Oligochaetochilus</i>	reddish aff. <i>hamata</i> with longer dorsal sepal point and sparser labellum hairs; Callitris woodland on rocky hills in the ranges, tablelands and western slopes of northern NSW (= 'Northern Tablelands' Bishop 2000) and southern Qld
<i>Pterostylis</i> species 'large curved flowers'	<i>Diplodium</i>	aff. <i>revoluta</i> with larger, more curved flowers; widespread in dry woodland and open forest through inland south-eastern Australia (Bishop 2000; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Malua Bay'	<i>Bunochilus</i>	aff. <i>macrosepala</i> ; heathy woodland at Malua Bay (Jones & Clements 2006)
<i>Pterostylis</i> species 'Mount Duval'	<i>Diplodium</i>	small aff. <i>alveata</i> with decurved petal tips and broad labellum with rounded apex just visible above the sinus; moist eucalypt forest near Armidale and Dorrigo (Bishop 2000)
<i>Pterostylis</i> species 'Mount Kaputar'	<i>Bunochilus</i>	aff. <i>lineata</i> with long, wide lateral sepals and a dark brown labellum with short bristles; open forest in the Nandewar Range
<i>Pterostylis</i> species 'Narrandera range'	<i>Oligochaetochilus</i>	large aff. <i>petrosa</i> with narrow lateral sepals with long tips and a more rounded labellum with a prominent basal constriction; dry woodland on rocky hills near Narrandera (Bishop 2000)
<i>Pterostylis</i> species 'New England striped'	<i>Diplodium</i>	aff. <i>alata</i> with narrow striped flowers with angled basally flat lateral sepals in profile with backswept tips; forest and woodland on the New England Tableland (Bishop 2000)
<i>Pterostylis</i> species 'Outback'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with broader, rounded lateral sepals; sparse shrublands on rocky hills between Renmark and Olary (SA) and near Broken Hill (NSW) (Bates 2011)
<i>Pterostylis</i> species 'Ranking Springs'	<i>Oligochaetochilus</i>	small aff. <i>biseta</i> with short tips to the lateral sepals; Callitris woodland on hills around Rankin Springs (Bishop 2000)
<i>Pterostylis</i> species 'sandstone heath'	<i>Speculantha</i>	summer flowering short aff. <i>parviflora</i> with crowded green, brown and white flowers with lateral sepals bowed in profile; heathland on shallow sandy soils and moss mats on sandstone, south of Sydney (Bishop 2000)
<i>Pterostylis</i> species 'Spring Ridge'	<i>Oligochaetochilus</i>	short aff. <i>biseta</i> with small flowers; dry woodland and open forest at Spring Ridge
<i>Pterostylis</i> species 'tablelands'	<i>Diplodium</i>	colourful aff. <i>reflexa</i> with semi-circle curved galea with decurved apex and labellum just protruding over the broad protruding sinus; grassy forest from Bungonia to New England (Bishop 2000)
<i>Pterostylis</i> species 'Torrington'	<i>Oligochaetochilus</i>	small aff. <i>praetermissa</i> with tiny, almost rounded labellum and lateral sepals wider than galea, labellum with ~8 pairs of marginal hairs; tea-tree thickets in northern granite region (Bishop 2000)
<i>Pterostylis</i> species 'Urbenville'	<i>Bunochilus</i>	aff. <i>barringtonensis</i> with a larger labellum with short, stiff hairs; open forest in the far northern volcanic ranges around Urbenville
<i>Pterostylis</i> species 'Warialda'	<i>Oligochaetochilus</i>	'markedly different' aff. <i>boormanii</i> ; dry woodland near Warialda (Bishop 2000)
<i>Pterostylis</i> species 'Wauchope'	<i>Speculantha</i>	autumn flowering, robust aff. <i>rubescens</i> with larger flowers long, narrow, bristly dorsal sepal and smooth petals and lateral sepals; lowland open forest on gravelly soils on the north coast (recta ms)
<i>Pterostylis</i> species 'Werrikimbe'	<i>Speculantha</i>	robust aff. <i>rubescens</i> with bristly flowers; moist open forest on escarpments in the Barrington Tops and Werrikimbe; <i>oresbia</i> ms
<i>Pterostylis</i> species 'wrinkled lateral sepals'	<i>Oligochaetochilus</i>	aff. <i>praetermissa</i> with wrinkled lateral sepals; woodland and open forest above 500 m; open forest on rocky hills in the Barrington Tops, New England Tableland and Warrumbungles (L. Copeland unpubl.)
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		
<i>Rhinerrhiza divitiflora</i> (F.Muell. ex Benth.) Rupp		
<i>Rhizanthella slateri</i> (Rupp) M.A.Clem. & P.J.Cribb		
<i>Rimacola elliptica</i> (R.Br.) Rupp		
<i>Sarcochilus aequalis</i> D.L.Jones & M.A.Clem.		
<i>Sarcochilus australis</i> (Lindl.) Rchb.f.		
<i>Sarcochilus dilatatus</i> F.Muell.		
<i>Sarcochilus eriochilus</i> Fitzg.		

<i>Sarcochilus falcatus</i> R.Br.	
<i>Sarcochilus fitzgeraldii</i> F.Muell.	
<i>Sarcochilus hartmannii</i> F.Muell.	
<i>Sarcochilus hillii</i> (F.Muell.) F.Muell.	
<i>Sarcochilus parviflorus</i> Lindl.	
<i>Sarcochilus spathulatus</i> R.S.Rogers	
<i>Sarcochilus weinthalii</i> F.M.Bailey	
<i>Sarcochilus</i> species 'Blue Knob'	aff. <i>hartmannii</i> with erect, dark coloured racemes of all white flowers; sheltered rocky outcrops and cliffs in the northern ranges (Jones 2006)
<i>Sarcochilus</i> species 'brittle leaves'	late flowering aff. <i>falcatus</i> with larger pale cupped flowers and brittle leaves; mountain moist forest and rainforest from Dorrigo to the Border Ranges
<i>Schistotylus purpuratus</i> (Rupp) Dockrill	
<i>Spiranthes alticola</i> D.L.Jones	very difficult to distinguish from (and doubtfully distinct from) <i>S. australis</i>
<i>Spiranthes australis</i> (R.Br.) Lindl.	
<i>Taeniophyllum muelleri</i> Benth.	
<i>Thelymitra adorata</i> Jeanes	
<i>Thelymitra alpicola</i> Jeanes	
<i>Thelymitra alpina</i> Jeanes	
<i>Thelymitra angustifolia</i> R.Br.	
<i>Thelymitra arenaria</i> Lindl.	
<i>Thelymitra atronitida</i> Jeanes	
<i>Thelymitra brevifolia</i> Jeanes	
<i>Thelymitra carnea</i> R.Br.	
<i>Thelymitra circumsepta</i> Fitzg.	
<i>Thelymitra cyanea</i> (Lindl.)	
<i>Thelymitra fragrans</i> D.L.Jones & M.A.Clem.	
<i>Thelymitra holmesii</i> Nicholls	
<i>Thelymitra incurva</i> Jeanes	
<i>Thelymitra ixioides</i> Sw.	
<i>Thelymitra juncifolia</i> Lindl.	
<i>Thelymitra kangaloonica</i> Jeanes	
<i>Thelymitra latifolia</i> R.J.Bates	
<i>Thelymitra longiloba</i> D.L.Jones & M.A.Clem.	
<i>Thelymitra malvina</i> M.A.Clem.	
<i>Thelymitra media</i> R.Br.	
<i>Thelymitra megalyptra</i> Fitzg.	
<i>Thelymitra nuda</i> R.Br.	
<i>Thelymitra pauciflora</i> R.Br.	
<i>Thelymitra peniculata</i> Jeanes	
<i>Thelymitra planicola</i> Jeanes	
<i>Thelymitra purpurata</i> Rupp	
<i>Thelymitra rubra</i> Fitzg.	
<i>Thelymitra simulata</i> D.L.Jones & M.A.Clem.	
<i>Thelymitra venosa</i> R.Br.	
<i>Thelymitra</i> species 'blue column'	aff. <i>pauciflora</i> with poorly opening purple-blue flowers with a purple-blue column apex; heathy woodland near Guyra (L. Copeland unpublished)
<i>Thelymitra</i> species 'central coast'	affinity uncertain; recognised by the purple violet flowers with similar colour hair tufts and the narrow horseshoe-shaped yellow apex to the post-anther lobe; heath and heathy woodland around Lake Macquarie
<i>Thelymitra</i> species 'fine spots'	aff. <i>ixioides</i> with self-pollinating flowers with few very fine spots on dorsal sepal and petals and irregularly glandular post-anther lobe; damp mountain forest, swampy and damp grassy areas above 800 m (Jones <i>et al.</i> 2008)
<i>Thelymitra</i> species 'Gibraltar Falls'	no information available (Australian National Herbarium record)
<i>Thelymitra</i> species 'ornate flowers'	small self-pollinating aff. <i>simulata</i> with freely opening flowers with shortly hooded deeply notched post-anther lobe; open forest in mountains above 750 m; resembles the New Zealand species <i>T. decora</i> (Jones <i>et al.</i> 2008)
<i>Thynniorchis huntianus</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>
<i>Zeuxine oblonga</i> R.S.Rogers & C.T.White	

**Total: 558 taxa (species, subspecies, varieties)**

<i>Caladenia arenaria</i> X <i>concinna</i>		Backhouse 2011
<i>Caladenia arenaria</i> X <i>rileyi</i>		Backhouse 2011
<i>Caladenia arenaria</i> X <i>stellata</i>		Backhouse 2011
<i>Caladenia arenaria</i> X species 'inland NSW'		Backhouse 2011
<i>Chiloglottis trapeziformis</i> X <i>valida</i>	× <i>pescottiana</i> R.S. Rogers	
<i>Corunastylis citriodora</i> X <i>fimbriata</i>		Jones 2006
<i>Corunastylis filiformis</i> X <i>fimbriata</i>		Jones 2006
<i>Cyanicula caerulea</i> X <i>Glossodia major</i>		
<i>Cyanicula caerulea</i> X <i>Glossodia minor</i>		Backhouse 2011
<i>Dendrobium aemulum</i> X <i>kingianum</i>		Jones 2006
<i>Dendrobium banksii</i> X <i>pugioniforme</i>		Jones 2006
<i>Dendrobium bowmanii</i> X <i>cucumerinum</i>		Jones 2006
<i>Dendrobium cucumerinum</i> X <i>linguiformis</i>		Jones 2006
<i>Dendrobium dolichophyllum</i> X <i>schoeninum</i>		Jones 2006
<i>Dendrobium fairfaxii</i> X <i>pugioniforme</i>		Jones 2006
<i>Dendrobium falcorostrum</i> X <i>gracilicaule</i>		Jones 2006
<i>Dendrobium falcorostrum</i> X <i>kingianum</i>		Jones 2006
<i>Dendrobium falcorostrum</i> X <i>speciosum</i>		Jones 2006
<i>Dendrobium gracilicaule</i> X <i>kingianum</i>	× <i>suffusum</i> Cady	
<i>Dendrobium gracilicaule</i> X <i>speciosum</i>	× <i>gracillimum</i> (Rupp) Leaney	
<i>Dendrobium kingianum</i> subsp. <i>kingianum</i> X <i>speciosum</i> subsp. <i>speciosum</i>	× <i>delicatum</i> (F.M. Bailey) F.M. Bailey	
<i>Dendrobium kingianum</i> var. <i>pulcherrimum</i> X <i>speciosum</i> subsp. <i>speciosum</i>	× <i>kestevenii</i> Rupp	
<i>Dendrobium linguiformis</i> X <i>pugioniforme</i>		Jones 2006
<i>Dendrobium linguiformis</i> X <i>teretifolium</i>		Jones 2006
<i>Dendrobium mortii</i> X <i>pugioniforme</i>		Jones 2006
<i>Dendrobium mortii</i> X <i>schoeninum</i>		Jones 2006
<i>Dendrobium mortii</i> X <i>teretifolium</i>		Jones 2006
<i>Dendrobium pugioniforme</i> X <i>schoeninum</i>		Jones 2006
<i>Dendrobium pugioniforme</i> X <i>striolatum</i>		Jones 2006
<i>Dendrobium striolatum</i> X <i>teretifolium</i>		Jones 2006
<i>Diuris abbreviata</i> X <i>eborensis</i>		
<i>Diuris abbreviata</i> X <i>goonooensis</i>		Jones 2006
<i>Diuris abbreviata</i> X <i>minor</i>		
<i>Diuris abbreviata</i> X <i>pedunculata</i>		
<i>Diuris abbreviata</i> X <i>venosa</i>		
<i>Diuris aurea</i> X <i>punctata</i>	× <i>nebulosa</i> D.L. Jones	
<i>Diuris behrii</i> X <i>punctata</i>		Backhouse, <i>et al.</i> 2016; Jones 2006
<i>Diuris behrii</i> X <i>tricolor</i>		G. Backhouse personal observation
<i>Diuris chrysantha</i> X <i>minor</i>		Jones 2006
<i>Diuris chrysiopsis</i> X <i>pardina</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Diuris chrysiopsis</i> X <i>platichila</i>	× <i>polymorpha</i> Messmer	
<i>Diuris eborensis</i> X species 'New England swamps'		
<i>Diuris goonooensis</i> X <i>tricolor</i>		Jones 2006
<i>Diuris pardina</i> X <i>subalpina</i>		
<i>Diuris punctata</i> X <i>sulphurea</i>		
<i>Glossodia major</i> X <i>minor</i>		Jeanes & Backhouse 2006
<i>Pterostylis alveata</i> X <i>ophioglossa</i>	× <i>furcillata</i> Rupp	
<i>Pterostylis atrans</i> X <i>decurva</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Pterostylis concinna</i> X <i>obtusa</i>		
<i>Pterostylis concinna</i> X <i>ophioglossa</i>	× <i>conoglossa</i> Upton	
<i>Pterostylis falcata</i> X <i>nutans</i>	× <i>ingens</i> (Rupp) D.L. Jones	
<i>Thelymitra brevifolia</i> X <i>juncifolia</i>		Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra carnea</i> X <i>flexuosa</i>		Jones 2006
<i>Thelymitra carnea</i> X <i>ixioides</i>	× <i>irregularis</i> Nicholls	
<i>Thelymitra carnea</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra ixioides</i> X <i>nuda</i>		Jones 2006
<i>Thelymitra ixioides</i> X <i>pauciflora</i>	× <i>truncata</i> R.S. Rogers	
<i>Thelymitra ixioides</i> X <i>peniculata</i>	× <i>merraniae</i> Nicholls	

*Thelymitra ixiooides* X *rubra*

Jones 2006

*Thelymitra juncifolia* X *pauciflora*Bates 2011; Jones *et al.* 2008**Total: 60 hybrids**

Northern Territory Species	Alternate Genus	Notes
<i>Arthrochilus byrnesii</i> Blaxell	<i>Phoringopsis</i>	
<i>Arthrochilus latipes</i> D.L.Jones		
<i>Arthrochilus</i> species 'Bradshaw Station'	<i>Phoringopsis</i>	no information available
<i>Arthrochilus</i> species 'spinifex dweller'	<i>Phoringopsis</i>	no information available; NT Herbarium record
<i>Calochilus caeruleus</i> L.O. Williams		
<i>Calochilus caesius</i> D.L.Jones		
<i>Calochilus holtzei</i> F.Muell.		
<i>Calochilus imperiosus</i> D.L.Jones		
<i>Chiloschista phyllorrhiza</i> (F.Muell.) Schltr.		
<i>Crepidium acuminatum</i> (D.Don) Szlach.	<i>Malaxis</i>	
<i>Crepidium fontinale</i> D.L.Jones & M.A.Clem.	<i>Malaxis</i>	
<i>Crepidium marsupichilum</i> (Upton) Szlach.	<i>Malaxis</i>	
<i>Cymbidium canaliculatum</i> R.Br.		many forms and varieties have been named (eg. var. <i>sparksii</i> ) although none have any formal taxonomic recognition
<i>Dendrobium dicuphum</i> F.Muell.	<i>Vappodes</i>	
<i>Dendrobium foelschei</i> F.Muell.	<i>Cepobaculum</i>	usually treated as a variety of <i>Dendrobium canaliculatum</i>
<i>Dendrobium johannis</i> Rchb.f.	<i>Cepobaculum</i>	
<i>Dendrobium lobbii</i> Teijsm. & Binn.	<i>Conostalix</i>	also called <i>Dendrobium paludica</i> Schltr.
<i>Dendrobium trilamellatum</i> J.J.Sm.	<i>Cepobaculum</i>	
<i>Didymoplexis pallens</i> Griffith		
<i>Dienia montana</i> (Blume) M.A.Clem. & D.L.Jones	<i>Malaxis</i>	
<i>Dipodium stenochilum</i> O.Schwarz		
<i>Empusa habenarina</i> (F.Muell.) Benth.	<i>Liparis</i>	
<i>Eulophia agrostophylla</i> Benth.		may be the same as and therefore a synonym of <i>Eulophia venosa</i>
<i>Eulophia venosa</i> (F.Muell.) Benth.		
<i>Geodorum densiflorum</i> (Lam.) Schltr.		also called <i>Geodorum terrestre</i> (L.) Garay
<i>Habenaria elongata</i> R.Br.		
<i>Habenaria ferdinandii</i> Schltr.		
<i>Habenaria halata</i> D.L.Jones		
<i>Habenaria hymenophylla</i> Schltr.		
<i>Habenaria ochroleuca</i> R.Br.		
<i>Habenaria rumphii</i> (Brogn.) Lindl.		
<i>Habenaria triplonema</i> Schltr.		
<i>Luisia corrugata</i> D.L.Jones		
<i>Nervilia aragoana</i> Gaud.		
<i>Nervilia crocifformis</i> (Zoll. & Mor.) Seidenf.		
<i>Nervilia holochila</i> (F.Muell.) Schltr.		
<i>Nervilia peltata</i> B.Gray & D.L.Jones		
<i>Nervilia plicata</i> (Andrews) Schltr.		also called <i>Nervilia dallachyana</i> (F.Muell. ex. Benth.)
<i>Nervilia uniflora</i> (F.Muell.) Schltr.		
<i>Pachystoma pubescens</i> Blume		
<i>Phaius amboinensis</i> Blume		
<i>Spathoglottis paulinae</i> F.Muell.		
<i>Thrixspermum congestum</i> (F.M.Bailey) Dockrill		
<i>Tropidia territorialis</i> D.L.Jones & M.A.Clem.		
<i>Zeuxine oblonga</i> R.S.Rogers & C.T.White		

**Total: 45 species**

Queensland Species	Alternate Genus	Notes
<i>Acianthella amplexicaulis</i> (F.M.Bailey) D.L.Jones & M.A.Clem.	<i>Acianthus</i>	
<i>Acianthella sublesta</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Acianthus</i>	
<i>Acianthus borealis</i> D.L.Jones		
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	
<i>Acianthus exsertus</i> R.Br.		
<i>Acianthus fornicatus</i> R.Br.		
<i>Acianthus ledwardii</i> Rupp		probably extinct
<i>Acianthus pusillus</i> D.L.Jones		
<i>Acianthus saxatilis</i> D.L.Jones & M.A.Clem.		
<i>Acriopsis emarginata</i> D.L.Jones & M.A.Clem.		
<i>Anoectochilus yatesiae</i> F.M.Bailey		
<i>Aphyllorchis anomala</i> Dockrill		
<i>Aphyllorchis queenslandica</i> Dockrill		
<i>Apostasia wallichii</i> R.Br.		also called <i>A. stylidioides</i> (F.Muell.) Rchb.f.
<i>Appendicula australiensis</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Podochilus</i>	
<i>Arthrochilus apectus</i> D.L.Jones		
<i>Arthrochilus aquilus</i> D.L.Jones		
<i>Arthrochilus corinnae</i> D.L.Jones		
<i>Arthrochilus dockrillii</i> Lavarack	<i>Phoringopsis</i>	
<i>Arthrochilus irritabilis</i> F.Muell.		
<i>Arthrochilus lavarackianus</i> (D.L.Jones) Lavarack	<i>Phoringopsis</i>	
<i>Arthrochilus oreophilus</i> D.L.Jones		
<i>Arthrochilus prolixus</i> D.L.Jones		
<i>Arthrochilus rosulatus</i> D.L.Jones		
<i>Arthrochilus sabulosus</i> D.L.Jones		
<i>Arthrochilus stenophyllus</i> D.L.Jones		
<i>Bromheadia pulchra</i> Schltr.		
<i>Bryobium dischorensis</i> (Schltr.) M.A.Clem. & D.L.Jones	<i>Eria</i>	may be the same as <i>Bryobium intermedium</i>
<i>Bryobium eriaeoides</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Eria</i>	
<i>Bryobium intermedium</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Eria</i>	may be the same as <i>Bryobium dischorensis</i>
<i>Bryobium irukandjianum</i> St.Cloud	<i>Eria</i>	
<i>Bryobium queenslandicum</i> T.E.Hunt	<i>Eria</i>	
<i>Bulbophyllum argyropus</i> (Endl.) Rchb.f.	<i>Adelopetalum</i>	
<i>Bulbophyllum baileyi</i> F.Muell.	<i>Carpamorchis</i>	
<i>Bulbophyllum boonjee</i> B.Gray & D.L.Jones	<i>Adelopetalum</i>	
<i>Bulbophyllum bowkettiae</i> F.M.Bailey	<i>Serpenticaulis</i>	
<i>Bulbophyllum bracteatum</i> Fitzg.	<i>Adelopetalum</i>	
<i>Bulbophyllum elisae</i> F.Muell.	<i>Adelopetalum</i>	
<i>Bulbophyllum evasum</i> T.E.Hunt & Rupp	<i>Kauorchis</i>	
<i>Bulbophyllum exiguum</i> F.Muell.	<i>Adelopetalum</i>	
<i>Bulbophyllum gadgarrense</i> Rupp	<i>Oxysepala</i>	
<i>Bulbophyllum globuliforme</i> Nicholls	<i>Oncophyllum</i>	
<i>Bulbophyllum gracillimum</i> Rolfe	<i>Cirrhopetalum</i>	
<i>Bulbophyllum grandimesense</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	
<i>Bulbophyllum johnsonii</i> T.E.Hunt	<i>Serpenticaulis</i>	
<i>Bulbophyllum kirkwoodiae</i> T.E.Hunt	<i>Serpenticaulis</i>	usually included in <i>Bulbophyllum johnsonii</i>
<i>Bulbophyllum lageniforme</i> F.Muell.	<i>Adelopetalum</i>	
<i>Bulbophyllum lamingtonense</i> D.L.Jones	<i>Oxysepala</i>	
<i>Bulbophyllum lewisense</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	
<i>Bulbophyllum lilianae</i> Rendle	<i>Adelopetalum</i>	
<i>Bulbophyllum longiflorum</i> Thouars	<i>Cirrhopetalum</i>	also called <i>B. clavigerum</i> (Fitzg.) F.M.Bailey
<i>Bulbophyllum macphersonii</i> Rupp	<i>Blepharochilum</i>	
<i>Bulbophyllum maxillare</i> (Lindl.) Rchb.f.	<i>Ehippium</i>	also called <i>B. masdevalliaceum</i> Kraenzl.
<i>Bulbophyllum minutissimum</i> F.Muell.	<i>Oncophyllum</i>	
<i>Bulbophyllum nematopodum</i> F.Muell.	<i>Papulipetalum</i>	



Queensland Species	Alternate Genus	Notes
<i>Bulbophyllum newportii</i> F.M.Bailey	<i>Adelopetalum</i>	
<i>Bulbophyllum radicans</i> F.M.Bailey	<i>Fruticicola</i>	
<i>Bulbophyllum schillerianum</i> Rchb.f. subspecies <i>maritimum</i> F.Muell.	<i>Oxysepala</i>	
<i>Bulbophyllum schillerianum</i> Rchb.f. subspecies <i>schillerianum</i>	<i>Oxysepala</i>	
<i>Bulbophyllum shepherdii</i> (F.Muell.) Rchb.f.	<i>Oxysepala</i>	
<i>Bulbophyllum sladeanum</i> A.D.Hawkes	<i>Blepharochilum</i>	sometimes called <i>Bulbophyllum macphersonii</i> var. <i>spathulatum</i>
<i>Bulbophyllum wadsworthii</i> Dockrill	<i>Oxysepala</i>	
<i>Bulbophyllum weinthalii</i> R.S.Rogers subspecies <i>striatum</i> D.L.Jones	<i>Spilorchis</i>	
<i>Bulbophyllum weinthalii</i> R.S.Rogers subspecies <i>weinthalii</i>	<i>Spilorchis</i>	
<i>Bulbophyllum whitei</i> T.E.Hunt & Rupp	<i>Serpenticaulis</i>	sometimes included in <i>Bulbophyllum johnsonii</i>
<i>Bulbophyllum wilkianum</i> T.E.Hunt	<i>Adelopetalum</i>	
<i>Bulbophyllum windsorens</i> B.Gray & D.L.Jones	<i>Oxysepala</i>	
<i>Bulbophyllum wolfei</i> B.Gray & D.L.Jones	<i>Serpenticaulis</i>	
<i>Caladenia alata</i> R.Br.	<i>Petalochilus</i>	
<i>Caladenia atroclavia</i> D.L.Jones & M.A.Clem.	<i>Arachnorchis</i>	
<i>Caladenia atrovespa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia catenata</i> (Sm.) Druce	<i>Petalochilus</i>	
<i>Caladenia catenata</i> (Sm.) Druce variety <i>gigantea</i> (R.S.Rogers) W.M.Curtis	<i>Petalochilus</i>	taxon of uncertain status
<i>Caladenia chamaephylla</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia flaccida</i> D.L.Jones	<i>Jonesiopsis</i>	South Australian records unlikely to be true <i>Caladenia flaccida</i> ; here listed as an undescribed taxon <i>Caladenia</i> species 'Flinders Ranges red'
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia gracillima</i> (Rupp) D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia picta</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia quadrifaria</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia</i> species 'Kilkoy Creek'	<i>Petalochilus</i>	aff. <i>carnea</i> of uncertain identity; Kilkoy Creek (Qld Herbarium record)
<i>Calanthe australasica</i> D.L.Jones & M.A.Clem.		
<i>Caleana major</i> R.Br.		
<i>Calochilus ammobius</i> D.L.Jones & B.Gray		
<i>Calochilus caeruleus</i> L.O. Williams		
<i>Calochilus campestris</i> R.Br.		
<i>Calochilus cleistanthus</i> D.L.Jones		
<i>Calochilus grandiflorus</i> (Benth.) Domin		
<i>Calochilus imperiosus</i> D.L.Jones		
<i>Calochilus metallicus</i> D.L.Jones		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus psednus</i> D.L.Jones & Lavarack		
<i>Calochilus therophilus</i> D.L.Jones		in Qld recorded as <i>Calochilus gracillimus</i>
<i>Cestichis angustilabris</i> (F.Muell.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis bracteata</i> (T.E.Hunt) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis coelogynoides</i> (F.Muell.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis condylobulbon</i> (Rchb.f.) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis fleckeri</i> (Nicholls) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis nugentiae</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cestichis swenssonii</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Cheirostylis notialis</i> D.L.Jones		
<i>Cheirostylis ovata</i> (F.M.Bailey) Schltr.		
<i>Chiloglottis diphylla</i> R.Br.		
<i>Chiloglottis formicifera</i> Fitzg.	<i>Myrmecchila</i>	
<i>Chiloglottis longiclavata</i> D.L.Jones		
<i>Chiloglottis sphyrnoides</i> D.L.Jones		
<i>Chiloglottis sylvestris</i> D.L.Jones & M.A.Clem.		
<i>Chiloglottis trullata</i> D.L.Jones	<i>Myrmecchila</i>	

Queensland Species	Alternate Genus	Notes
<i>Chiloglottis truncata</i> D.L.Jones & M.A.Clem.	<i>Myrmecchila</i>	
<i>Chiloglottis</i> species 'Bald Rock'	<i>Myrmecchila</i>	small, spring flowering aff. <i>formicefera</i> with recurved labellum apex, calli extending almost to apex and narrower column wings; grassy open forest on rich loams; border region in Girraween and Bald Rock National Parks (Bishop 2000)
<i>Chiloglottis</i> species 'Maryborough'		no information available (Qld. Herbarium record)
<i>Chiloschista phyllorhiza</i> (F.Muell.) Schltr.		
<i>Cooktownia robertsii</i> D.L.Jones		
<i>Corunastylis acuminata</i> (R.S.Rogers) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis alticola</i> (D.L.Jones & B. Gray) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis archeri</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	'true' <i>archeri</i> almost certainly does not occur in NSW & Qld, with these plants probably comprising a complex of several different un-named species
<i>Corunastylis conferta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis cranei</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis filiformis</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis fimbriata</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis parvicalla</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis pedersonii</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis psammophila</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis pumila</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis rufa</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis sagittifera</i> (Rupp) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis sigmaidea</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis tecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis valida</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis</i> species 'Fraser Island'	<i>Genoplesium</i>	no further information (Qld Herbarium record)
<i>Corunastylis</i> species 'granite ranges'	<i>Genoplesium</i>	spring-summer flowering, robust aff. <i>archeri</i> with more crowded flowers with strongly upcurved lateral sepals; open forest on granite ranges of north-eastern NSW (L. Copeland unpublished) and south-eastern Qld
<i>Corunastylis</i> species 'Mount Mee'	<i>Genoplesium</i>	late summer-flowering aff. <i>densa</i> with numerous somewhat spaced flowers with broad lateral sepals and a relatively short leaf ending below the lowest flower; 'clay pan' heath on Mt Mee (David James personal communication)
<i>Corunastylis</i> species 'Baby Bay'	<i>Genoplesium</i>	no further information (Qld Herbarium record)
<i>Corunastylis</i> species 'Wyberba'	<i>Genoplesium</i>	no further information (Qld Herbarium record)
<i>Corybas abellianus</i> Dockrill		
<i>Corybas aconitiflorus</i> Salisb.		
<i>Corybas barbarae</i> D.L.Jones		
<i>Corybas cerasinus</i> D.L.Jones & B.Gray		
<i>Corybas fimbriatus</i> (R.Br.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	
<i>Corybas hispidus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas montanus</i> D.L.Jones	<i>Anzybas</i>	
<i>Corybas undulatus</i> (A.Cunn.) Rupp	<i>Corysanthes</i>	
<i>Corymborkis veratrifolia</i> (Reinw.) Blume		
<i>Crepidium fimbriatum</i> (Lavarack) Szlach.	<i>Malaxis</i>	
<i>Crepidium flavovirens</i> D.L.Jones & M.A.Clem.	<i>Malaxis</i>	
<i>Crepidium lawleri</i> (Lavarack & B.Gray) Szlach.	<i>Malaxis</i>	
<i>Crepidium marsupichilum</i> (Upton) Szlach.	<i>Malaxis</i>	
<i>Cryptostylis erecta</i> R.Br.		
<i>Cryptostylis hunteriana</i> Nicholls		
<i>Cryptostylis leptochila</i> Benth.		
<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		
<i>Cyanicula caerulea</i> (R.Br.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cymbidium canaliculatum</i> R.Br.		many forms and varieties have been named (eg. var. <i>sparksii</i> ) although none have any formal taxonomic recognition
<i>Cymbidium madidum</i> Lindl.		
<i>Cymbidium suave</i> R.Br.		
<i>Cyrtostylis reniformi</i> R.Br.		

Queensland Species	Alternate Genus	Notes
<i>Demorchis queenslandica</i> (Dockrill) D.L.Jones & M.A.Clem.	<i>Gastrodia</i>	
<i>Dendrobium adae</i> F.M.Bailey	<i>Thelychiton</i>	
<i>Dendrobium aemulum</i> R.Br.	<i>Tropilis</i>	
<i>Dendrobium agrostophyllum</i> F.Muell.	<i>Trachyrhizum</i>	
( <i>Dendrobium</i> ) <i>angustum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium antennatum</i> Lindl.	<i>Ceratobium</i>	also called <i>Dendrobium dalbertsii</i> Rchb.f.
<i>Dendrobium aphyllum</i> (Roxb.) C.E.C.Fisch.		occurrence in Australia doubtful
<i>Dendrobium aurantiacopurpureum</i> Nicholls	<i>Davejonesia</i>	doubtfully distinct from <i>Dendrobium lichenastrum</i>
<i>Dendrobium baileyi</i> F.Muell.	<i>Grastidium</i>	
<i>Dendrobium bifalce</i> Lindl.	<i>Leioanthum</i>	
<i>Dendrobium bigibbum</i> Lindl.	<i>Vappodes</i>	
<i>Dendrobium bowmanii</i> Benth.	<i>Dockrillia</i>	
<i>Dendrobium brevicaudum</i> D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	
<i>Dendrobium calamiformis</i> Lodd. ex. Lindl.	<i>Dockrillia</i>	
<i>Dendrobium callitrophilum</i> B.Gray & D.L.Jones	<i>Tropilis</i>	usually treated as a variety of <i>Dendrobium aemulum</i>
<i>Dendrobium canaliculatum</i> R.Br.	<i>Cepobaculum</i>	
<i>Dendrobium cancroides</i> T.E.Hunt	<i>Grastidium</i>	
<i>Dendrobium carrii</i> Rupp & C.T.White	<i>Australorchis</i>	
<i>Dendrobium carronii</i> Lavarack & P.J.Cribb	<i>Cepobaculum</i>	
<i>Dendrobium collinsii</i> (Lavarack) Schuit. & P.B.Adams	<i>Cadetia</i>	
<i>Dendrobium comatum</i> (Blume) Lindl.	<i>Flickingeria</i>	described as <i>Flickingeria clementsii</i>
<i>Dendrobium convexum</i> (Blume) Lindl.	<i>Abaxianthus</i>	
( <i>Dendrobium</i> ) <i>crassum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium cucumerinum</i> MacLeay ex Lindl.	<i>Dockrillia</i>	
<i>Dendrobium discolor</i> R.Br. subspecies <i>discolor</i>	<i>Durabaculum</i>	sometimes called <i>Dendrobium undulatum</i>
<i>Dendrobium discolor</i> R.Br. subspecies <i>incurvata</i> Liddle & P.I. Forst.	<i>Durabaculum</i>	
<i>Dendrobium discolor</i> R.Br. variety <i>broomfieldii</i> Fitzg.	<i>Durabaculum</i>	
<i>Dendrobium discolor</i> R.Br. variety <i>fimbrilabium</i> (Rchb.f.) Dockrill	<i>Durabaculum</i>	
<i>Dendrobium discolor</i> R.Br. variety <i>fuscum</i> (Fitzg.) Dockrill	<i>Durabaculum</i>	
<i>Dendrobium dolichophyllum</i> D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	
( <i>Dendrobium</i> ) <i>eburneum</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
( <i>Dendrobium</i> ) <i>eungellensis</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium falcorostrum</i> Fitzg.	<i>Thelychiton</i>	
<i>Dendrobium fellowsii</i> F.Muell.	<i>Eleutheroglossum</i>	
<i>Dendrobium finniganense</i> D.L.Jones	<i>Thelychiton</i>	
<i>Dendrobium fleckeri</i> Rupp & C.T.White	<i>Thelychiton</i>	
<i>Dendrobium foelschei</i> F.Muell.	<i>Cepobaculum</i>	usually treated as a variety of <i>Dendrobium canaliculatum</i>
<i>Dendrobium funiforme</i> Blume	<i>Cadetia</i> , <i>Sarcocadetia</i>	formerly called <i>Cadetia waryana</i> & <i>Sarcocadetia waryana</i>
<i>Dendrobium glabrum</i> J.J.Sm.	<i>Diplocaulobium</i>	
<i>Dendrobium gracilicaule</i> F.Muell.	<i>Thelychiton</i>	
<i>Dendrobium johannis</i> Rchb.f.	<i>Cepobaculum</i>	
<i>Dendrobium jonesii</i> Rendle subspecies <i>bancroftianum</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Thelychiton</i>	doubtfully distinct from subsp. <i>jonesii</i>
<i>Dendrobium jonesii</i> Rendle subspecies <i>blackburnii</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Thelychiton</i>	doubtfully distinct from subsp. <i>jonesii</i>
<i>Dendrobium jonesii</i> Rendle subspecies <i>jonesii</i>	<i>Thelychiton</i>	
<i>Dendrobium jonesii</i> Rendle subspecies <i>magnificum</i> (Dockrill) D.P.Banks & Clemesha	<i>Thelychiton</i>	usually called var. <i>magnificum</i>
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>kingianum</i>	<i>Thelychiton</i>	several varieties have been named although these are recognised mainly in cultivation and not taxonomically; one variety included here
<i>Dendrobium kingianum</i> Bidw. ex Lindl. subspecies <i>canarvonense</i> P.B.Adams	<i>Thelychiton</i>	considered a full species <i>D. canarvonense</i> by Jones (2006)
<i>Dendrobium lichenastrum</i> (F.Muell.) Rolfe	<i>Davejonesia</i>	
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>huntianum</i> Rupp	<i>Dockrillia</i>	
<i>Dendrobium linguiforme</i> (Sw.) Brieger variety <i>linguiforme</i>	<i>Dockrillia</i>	
<i>Dendrobium lithocola</i> D.L.Jones & M.A.Clem.	<i>Vappodes</i>	usually treated as a variety of <i>Dendrobium bigibbum</i>
<i>Dendrobium lobbii</i> Teijsm. & Binn.	<i>Conostalix</i>	also called <i>Dendrobium paludica</i> Schltr.

Queensland Species	Alternate Genus	Notes
<i>Dendrobium luteociliium</i> Rupp	<i>Grastidium</i>	sometimes included in <i>Dendrobium pruinosum</i> Teijsm. & Binn
<i>Dendrobium maidenianum</i> Schltr.	<i>Cadetia</i>	
<i>Dendrobium malbrownii</i> Dockrill	<i>Monanthos</i>	
<i>Dendrobium mirbelianum</i> Gaudich.	<i>Durabaculum</i>	
<i>Dendrobium monophyllum</i> F.Muell.	<i>Australorchis</i>	
<i>Dendrobium mortii</i> F.Muell.	<i>Dockrillia</i>	
<i>Dendrobium nindii</i> W.Hill	<i>Durabaculum</i>	
<i>Dendrobium nitidum</i> F.M.Bailey	<i>Thelychiton</i>	a taxon of uncertain status; variously included in <i>Dendrobium speciosum</i> subs. <i>pedunculatum</i> var. <i>boreale</i> ; also considered a hybrid, a form of <i>Dendrobium xgracillimum</i>
<i>Dendrobium nugentii</i> (F.M.Bailey) D.L.Jones & M.A.Clem.	<i>Dockrillia</i>	
<i>Dendrobium phalaenopsis</i> Fitzg.	<i>Vappodes</i>	usually treated as a variety of <i>Dendrobium bigibbum</i>
<i>Dendrobium prenticei</i> (F.Muell.) Nicholls	<i>Davejonesia</i>	sometimes included in <i>Dendrobium lichenastrum</i>
<i>Dendrobium pugioniforme</i> A.Cunn.	<i>Dockrillia</i>	
<i>Dendrobium racemosum</i> (Nicholls) Clemesha & Dockrill	<i>Dockrillia</i>	
( <i>Dendrobium</i> ) <i>radiata</i> (D.L.Jones & M.A.Clem.)	<i>Tropilis</i>	doubtfully distinct from <i>Dendrobium aemulum</i> ; described in <i>Tropilis</i> , new combination in <i>Dendrobium</i> not yet made
<i>Dendrobium rigidum</i> R.Br.	<i>Dockrillia</i>	
<i>Dendrobium schneiderae</i> F.M.Bailey variety <i>majus</i> Rupp	<i>Australorchis</i>	renamed as <i>Australorchis eungellensis</i> D.L. Jones & M.A. Clem.; that epithet combination in <i>Dendrobium</i> not yet made
<i>Dendrobium schneiderae</i> F.M.Bailey variety <i>schneiderae</i>	<i>Australorchis</i>	
<i>Dendrobium schoeninum</i> Lindl.	<i>Dockrillia</i>	
<i>Dendrobium semifusum</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Cepobaculum</i>	sometimes included in <i>Dendrobium trilamellatum</i>
<i>Dendrobium smillieae</i> (F.Muell.) Fitzg.	<i>Coelandria</i>	
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>blackdownense</i> P.B.Adams	<i>Thelychiton</i>	alternative name (in part) <i>Dendrobium (Thelychiton) coriaceum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>capricornicum</i> Clemesha	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) capricornicum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>carnavonense</i> P.B.Adams	<i>Thelychiton</i>	alternative name (in part) <i>Dendrobium (Thelychiton) coriaceum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>curvicaule</i> F.M.Bailey	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) spectabilis</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>grandiflorum</i> F.M.Bailey	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) rex</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>hillii</i> Masters	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) tarberi</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>pedunculatum</i> variety <i>pedunculatum</i> (Clemesha) D.P.Banks & Clemesha	<i>Thelychiton</i>	alternative name <i>Dendrobium (Thelychiton) pedunculatum</i>
<i>Dendrobium speciosum</i> Sm. subspecies <i>pedunculatum</i> variety <i>boreale</i> P.B.Adams, J.M.Burke and S.D.Lawson	<i>Thelychiton</i>	alternative names <i>Dendrobium (Thelychiton) biconvexum</i> , <i>D. (T.) curvicaule</i> , <i>D. (T.) rupicolum</i>
<i>Dendrobium stelliferum</i> J.J.Sm.	<i>Diplocaulobium</i>	Australian record doubtful
<i>Dendrobium stuartii</i> F.M.Bailey		sometimes included in <i>Dendrobium macrostachyum</i> Lindl.
<i>Dendrobium sulphureum</i> Schltr.	<i>Dockrillia</i>	
<i>Dendrobium tattonianum</i> Bateman ex Rchb.f.		sometimes included in <i>Dendrobium canaliculatum</i>
<i>Dendrobium taylori</i> (F.Muell.) F.M.Bailey	<i>Cadetia</i>	
<i>Dendrobium teretifolium</i> R.Br.	<i>Dockrillia</i>	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>cataractum</i> P.B.Adams, S.D.Lawson and G.A.Paterson	<i>Tetrabaculum</i>	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>giganteum</i> variety <i>giganteum</i> (Leaney) P.B.Adams	<i>Tetrabaculum</i>	alternative name <i>Dendrobium (Tetrabaculum) capitisyork</i>
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>giganteum</i> variety <i>cacatua</i> M.A.Clem. & D.L.Jones	<i>Tetrabaculum</i>	alternative name <i>Dendrobium (Tetrabaculum) cacatua</i>
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>tetragonum</i> variety <i>serpentis</i> P.B.Adams	<i>Tetrabaculum</i>	
<i>Dendrobium tetragonum</i> A.Cunn. subspecies <i>tetragonum</i> variety <i>tetragonum</i>	<i>Tetrabaculum</i>	
<i>Dendrobium toressae</i> (F.M.Bailey) Dockrill	<i>Stilbophyllum</i>	
<i>Dendrobium tozerense</i> Lavarack	<i>Grastidium</i>	
<i>Dendrobium trilamellatum</i> J.J.Sm.	<i>Cepobaculum</i>	
<i>Dendrobium uniflos</i> F.M.Bailey	<i>Cadetia</i>	possibly a synonym of <i>Dendrobium taylori</i>
<i>Dendrobium wassellii</i> S.T.Blake	<i>Dockrillia</i>	
<i>Didymoplexis pallens</i> Griffith		
<i>Dienia montana</i> (Blume) M.A.Clem. & D.L.Jones	<i>Malaxis</i>	

Queensland Species	Alternate Genus	Notes
<i>Dipodium atropurpureum</i> D.L.Jones		
<i>Dipodium elegantulum</i> D.L.Jones		
<i>Dipodium ensifolium</i> F.Muell.		
<i>Dipodium hamiltonianum</i> F.M.Bailey		
<i>Dipodium pandanum</i> F.M.Bailey		
<i>Dipodium pictum</i> (Lindl.) Rchb.f.		taxon of uncertain status
<i>Dipodium pulchellum</i> D.L.Jones & M.A.Clem		
<i>Dipodium punctatum</i> (Sm.) R.Br.		
<i>Dipodium stenochilum</i> O.Schwarz		
<i>Dipodium variegatum</i> M.A.Clem. & D.L.Jones		
<i>Diteilis collinsii</i> (B.Gray) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Diteilis petricola</i> D.L.Jones & B.Gray	<i>Liparis</i>	
<i>Diteilis simmondsii</i> (F.M.Bailey) M.A.Clem. & D.L. Jones	<i>Liparis</i>	
<i>Diuris abbreviata</i> Benth.		
<i>Diuris alba</i> R.Br.		
<i>Diuris aurea</i> Sm.		Qld records possibly erroneous, perhaps misidentifications of <i>Diuris chrysantha</i>
<i>Diuris chrysantha</i> D.L.Jones & M.A.Clem.		
<i>Diuris curta</i> D.L.Jones		
<i>Diuris exitela</i> D.L.Jones		
<i>Diuris goanoensis</i> Rupp		called <i>Diuris maculata</i> in Qld
<i>Diuris luteola</i> D.L.Jones & B.Gray		
<i>Diuris oporina</i> D.L.Jones		
<i>Diuris parvipetala</i> (Dockrill) D.L.Jones & M.A.Clem.		
<i>Diuris pedunculata</i> R.Br.		
<i>Diuris punctata</i> Sm. variety <i>punctata</i>		Qld records are most likely attributable to several undescribed species including 'granite ranges' and 'NSW North Coast'; possibly also misidentifications of <i>D. alba</i> , <i>D. oporina</i> and <i>D. parvipetala</i>
<i>Diuris sulphurea</i> R.Br.		
<i>Diuris tricolor</i> Fitzg.		
<i>Diuris unica</i> D.L.Jones		
<i>Diuris</i> species 'Byfield'		no information available (Queensland Herbarium record)
<i>Diuris</i> species 'Carnarvon'		aff. <i>punctata</i> ; sparse woodland on sandstone at Mt Moffat & Carnarvon Range; AVH records listed as <i>D. punctata</i> but almost certainly not this species; <i>Diuris parvipetala</i> is also recorded from the region
<i>Diuris</i> species 'Jollys Falls'		no information available (Qld Herbarium record)
<i>Diuris</i> species 'Tenterfield/Stanthorpe'		aff. <i>punctata</i> with pale mottled flowers with relatively short lateral sepals and white between the labellum callus ridges; open forest in the granite ranges around Tenterfield NSW and Stanthorpe Qld
<i>Drymoanthus minutus</i> Nicholls		
<i>Empusa habenarina</i> (F.Muell.) Benth.	<i>Liparis</i>	
<i>Epipogium roseum</i> (D.Don) Lindl.		
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus petricola</i> D.L.Jones & M.A.Clem.		
<i>Eriochilus</i> species 'smooth leaf'		aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<i>Erythrorchis cassythoides</i> (A.Cunn. ex Lindl.) Garay		
<i>Eulophia agrostophylla</i> Benth.		may be the same as and therefore a synonym of <i>Eulophia venosa</i>
<i>Eulophia bicallosa</i> (D.Don) P.F.Hunt & Summerh.		
<i>Eulophia pelorica</i> D.L.Jones & M.A.Clem.		
<i>Eulophia venosa</i> (F.Muell.) Benth.		
<i>Eulophia zollingeri</i> (Rchb.f.) J.J.Sm.		
<i>Gastrodia crebriflora</i> D.L.Jones		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Gastrodia urceolata</i> D.L.Jones		
<i>Geodorum densiflorum</i> (Lam.) Schltr.		also called <i>Geodorum terrestre</i> (L.) Garay
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Glossodia minor</i> R.Br.	<i>Caladenia</i>	alternative name <i>Caladenia minorata</i> M.A. Clem.

Queensland Species	Alternate Genus	Notes
<i>Goodyera grandis</i> (Blume) Blume		
<i>Goodyera viridiflora</i> (Blume) Blume		also called <i>Eucosia umbrosa</i> D.L.Jones & M.A.Clem.
<i>Habenaria chlorosepala</i> D.L.Jones		
<i>Habenaria elongata</i> R.Br.		
<i>Habenaria euryloba</i> D.L.Jones		
<i>Habenaria exilis</i> D.L.Jones		
<i>Habenaria fuscina</i> D.L.Jones		
<i>Habenaria harroldii</i> D.L.Jones		
<i>Habenaria hymenophylla</i> Schltr.		
<i>Habenaria macraithii</i> Lavarack		
<i>Habenaria praecox</i> Lavarack & Dockrill		
<i>Habenaria propinquior</i> Rchb.f.		
<i>Habenaria rumphii</i> (Brogn.) Lindl.		
<i>Habenaria triplonema</i> Schltr.		
<i>Habenaria xanthantha</i> F.Muell.		
<i>Hetaeria oblongifolia</i> Blume		
<i>Luisia atacta</i> D.L.Jones		called <i>Luisia teretifolia</i> in Qld
<i>Lyperanthus suaveolens</i> R.Br.		
<i>Micropera fasciculata</i> (Lindl.) Garay		
<i>Microtis parviflora</i> R.Br.		
<i>Microtis rara</i> R.Br.		identity and distribution uncertain, thought to be only NSW & Qld (Jones 2006), although also claimed to be in Sth Aust (Bates 2011)
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		
<i>Mobilabium hamatum</i> Rupp		
<i>Nervilia aragoana</i> Gaud.		
<i>Nervilia crocifformis</i> (Zoll. & Mor.) Seidenf.		
<i>Nervilia holochila</i> (F.Muell.) Schltr.		
<i>Nervilia peltata</i> B.Gray & D.L.Jones		
<i>Nervilia plicata</i> (Andrews) Schltr.		also called <i>Nervilia dallachyana</i> (F.Muell. ex. Benth.)
<i>Nervilia uniflora</i> (F.Muell.) Schltr.		
<i>Oberonia attenuata</i> Dockrill		
<i>Oberonia carnosa</i> Lavarack		
<i>Oberonia complanata</i> (A.Cunn.) M.A.Clem. & D.L.Jones		
<i>Oberonia crateriformis</i> D.L.Jones & M.A.Clem.		
<i>Oberonia flavescens</i> D.L.Jones & M.A.Clem.		
<i>Oberonia palmicola</i> F.Muell.		also included in <i>Oberonia titania</i>
<i>Oberonia rimachila</i> D.L.Jones & M.A.Clem.		
<i>Octarrhena pusilla</i> (F.M.Bailey) M.A.Clem. & D.L.Jones		
<i>Orthoceras strictum</i> R.Br.		
<i>Pachystoma pubescens</i> Blume		
<i>Papillilabium beckeri</i> (F.Muell. ex Benth.) Dockrill		
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Peristeranthus hillii</i> (F.Muell.) T.E.Hunt		
<i>Peristylus banfieldii</i> (F.M.Bailey) Lavarack		
<i>Peristylus candidus</i> J.J.Sm.		
<i>Peristylus chlorandrellus</i> D.L.Jones & M.A.Clem.		
<i>Peristylus papuanus</i> (Kraenzl.) J.J.Sm.		
<i>Phaius australis</i> F.Muell.		
<i>Phaius bernaysii</i> Rowland ex Rchb.f.		
<i>Phaius pictus</i> T.E.Hunt		
<i>Phalaenopsis rosenstromii</i> F.M.Bailey		also called <i>P. amabilis</i> subspecies <i>rosenstromii</i>
<i>Pholidota imbricata</i> Hook	<i>Colelogyne</i>	
<i>Pinalia fitzalanii</i> (F.Muell.) Kuntze	<i>Eria, Hymeneria</i>	
<i>Pinalia kingii</i> (F.Muell.) Kuntze	<i>Eria, Hymeneria</i>	
<i>Plectorrhiza brevilabris</i> (F.Muell.) Dockrill		
<i>Plectorrhiza tridentata</i> (Lindl.) Dockrill		
<i>Plexaure crassiuscula</i> (Nicholls) M.A.Clem. & D.L.Jones		
<i>Pomatocalpa macphersonii</i> (F.Muell.) T.E.Hunt		
<i>Pomatocalpa marsupiale</i> (Kraenzl.) J.J.Sm.		



Queensland Species	Alternate Genus	Notes
<i>Prasophyllum australe</i> R.Br.		
<i>Prasophyllum brevilabre</i> (Lindl.) Hook.f.		
<i>Prasophyllum campestre</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum exilis</i> D.L.Jones & R.J.Bates		
<i>Prasophyllum flavum</i> R.Br.		
<i>Prasophyllum incompositum</i> D.L.Jones		
<i>Prasophyllum wallum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum</i> species 'New England batholith'		late spring-summer flowering aff. <i>odoratum</i> with crowded flowers with large, deeply ruffled labellum; open forest in the granite ranges above 900 m on New England Tableland NSW (L. Copeland unpubl.) & near Stanthorpe Qld
<i>Pseudovanilla foliata</i> (F.Muell.) Garay		
<i>Pterostylis acuminata</i> R.Br.		
<i>Pterostylis anaclasta</i> (D.L.Jones) Janes & Duretto		
<i>Pterostylis anatona</i> D.L.Jones		
<i>Pterostylis antennifera</i> D.L.Jones	<i>Specularantha</i>	
<i>Pterostylis aquilonia</i> D.L.Jones & B.Gray	<i>Diplodium</i>	
<i>Pterostylis baptistii</i> Fitzg.		
<i>Pterostylis bicolor</i> M.A.Clem. & D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis bicornis</i> D.L.Jones & M.A.Clem.	<i>Petrochis</i>	
<i>Pterostylis caligna</i> M.T.Mathieson		
<i>Pterostylis cobarensis</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis crassichila</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis curta</i> R.Br.		
<i>Pterostylis cycnocephala</i> Fitzg.	<i>Hymenochilus</i>	
<i>Pterostylis daintreeana</i> Benth.	<i>Pharochilum</i>	also spelt <i>daintreana</i>
<i>Pterostylis depauperata</i> F.M.Bailey	<i>Crangonorchis</i>	
<i>Pterostylis erecta</i> T.E.Hunt		
<i>Pterostylis exquisita</i> D.L.Jones	<i>Specularantha</i>	
<i>Pterostylis extranea</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	
<i>Pterostylis fischii</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis grandiflora</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis hildae</i> Nicholls		
<i>Pterostylis hispidula</i> Fitzg.		
<i>Pterostylis longicurva</i> Rupp	<i>Diplodium</i>	
<i>Pterostylis longifolia</i> R.Br.	<i>Bunochilus</i>	
<i>Pterostylis mitchellii</i> Lindl.	<i>Oligochaetochilus</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis mystacina</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	
<i>Pterostylis nigricans</i> D.L.Jones & M.A.Clem	<i>Specularantha</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis obtusa</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis ophioglossa</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis parviflora</i> R.Br.	<i>Specularantha</i>	
<i>Pterostylis pearsonii</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	
<i>Pterostylis pedoglossa</i> Fitzg.	<i>Crangonorchis</i>	
<i>Pterostylis pedunculata</i> R.Br.		
<i>Pterostylis praetermissa</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis procera</i> D.L.Jones & M.A.Clem.		
<i>Pterostylis rufa</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis russellii</i> T.E.Hunt	<i>Diplodium</i>	
<i>Pterostylis scoliosa</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis setifera</i> M.A.Clem., Matthias & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis stricta</i> Clemesha & B.Gray		
<i>Pterostylis taurus</i> M.A.Clem. & D.L.Jones	<i>Taurantha</i>	
<i>Pterostylis thulia</i> (D.L.Jones) Janes & Duretto	<i>Oligochaetochilus</i>	
<i>Pterostylis torquata</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis vitrea</i> (D.L.Jones) Bostock	<i>Bunochilus</i>	

Queensland Species	Alternate Genus	Notes
<i>Pterostylis woollsii</i> Fitzg.	<i>Oligochaetochilus</i>	
<i>Pterostylis</i> species 'Border Ranges'	<i>Bunochilus</i>	aff. <i>melagramma</i> with an obovate labellum; open forest in Border Ranges region (Jones & Clements 2006)
<i>Pterostylis</i> species 'Gundiah'		no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'hooked sepals'	<i>Oligochaetochilus</i>	reddish aff. <i>hamata</i> with longer dorsal sepal point and sparser labellum hairs; Callitris woodland on rocky hills in the ranges, tablelands and western slopes of northern NSW (= 'Northern Tablelands' Bishop 2000) and southern Qld
<i>Pterostylis</i> species 'Joiners Ridge'		no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'large curved flowers'	<i>Diplodium</i>	aff. <i>revoluta</i> with larger, more curved flowers; widespread in dry woodland and open forest through inland south-eastern Australia (Bishop 2000; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Mount Maroon'		no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Mount Mee'		no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'Mount Moffat'		no information available (Qld Herbarium record)
<i>Pterostylis</i> species 'south-east Queensland'	<i>Oligochaetochilus</i>	aff. <i>chaetophora</i> from woodland and open forest on steep rocky hills; Gayndah to Tamborine Mountain; Qld records of <i>P. chaetophora</i> are this taxon
<i>Rhinerrhiza divitiflora</i> (F.Muell. ex Benth.) Rupp		
<i>Rhinerrhizopsis matutina</i> D.L.Jones & M.A.Clem.	<i>Sarcochilus</i>	formerly <i>Sarcochilus moorei</i>
<i>Rhipidorchis micrantha</i> (A.Rich.) D.L.Jones & M.A.Clem.		
<i>Rhizanthella omissa</i> D.L.Jones & M.A.Clem.		
<i>Rhomboda polygonoides</i> (F.Muell.) Ormerod	<i>Zeuxine</i>	
<i>Robiquetia gracilistipes</i> (Schltr.) J.J.Sm.		
<i>Robiquetia wassellii</i> Dockrill		
<i>Saccolabiopsis armitii</i> (F.Muell.) Dockrill		
<i>Saccolabiopsis rectifolia</i> (Dockrill) Garay		
<i>Salacistic ochroleuca</i> (F.M.Bailey) M.A.Clem. & D.L.Jones	<i>Goodyera</i>	
<i>Sarcochilus argochilus</i> D.L.Jones & M.A.Clem.		called <i>S. olivaceus</i> in Queensland
<i>Sarcochilus borealis</i> (Nicholls) M.A.Clem. & D.L.Jones		called <i>S. olivaceus</i> in Queensland
<i>Sarcochilus ceciliae</i> F.Muell.		
<i>Sarcochilus dilatatus</i> F.Muell.		
<i>Sarcochilus eriochilus</i> Fitzg.		
<i>Sarcochilus falcatus</i> R.Br.		
<i>Sarcochilus fitzgeraldii</i> F.Muell.		
<i>Sarcochilus hartmannii</i> F.Muell.		
<i>Sarcochilus hillii</i> (F.Muell.) F.Muell.		
<i>Sarcochilus hirticalcar</i> (Dockrill) M.A.Clem. & B.J.Wallace		
<i>Sarcochilus minutiflos</i> F.M.Bailey		
<i>Sarcochilus parviflorus</i> Lindl.		called <i>S. olivaceus</i> in Queensland
<i>Sarcochilus roseus</i> (Clemesha) Clemesha		
<i>Sarcochilus serrulatus</i> D.L.Jones		
<i>Sarcochilus spathulatus</i> R.S.Rogers		
<i>Sarcochilus weinthalii</i> F.M.Bailey		
<i>Sarcochilus</i> species 'brittle leaves'		late flowering aff. <i>falcatus</i> with larger, pale, cupped flowers and brittle leaves; mountain moist forest and rainforest from Dorrigo to the Border Ranges
<i>Schistotylus purpuratus</i> (Rupp) Dockrill		
<i>Schoenorchis micrantha</i> Reinwardt ex Blume		
<i>Schoenorchis sarcophylla</i> Schltr.		
<i>Spathoglottis paulinae</i> F.Muell.		
<i>Spathoglottis plicata</i> Blume		
<i>Spiranthes australis</i> (R.Br.) Lindl.		
<i>Spiranthes sinensis</i> (Pers.) Ames		uncertain whether this is 'true' <i>sinensis</i> or <i>Spiranthes australis</i>
<i>Taeniophyllum clementsii</i> (D.L.Jones & B.Gray) Kocyan & Schuit.	<i>Microtatorchis</i>	
<i>Taeniophyllum confertum</i> B.Gray & D.L.Jones		
<i>Taeniophyllum cymbiforme</i> T.E.Hunt		
<i>Taeniophyllum lobatum</i> Dockrill		
<i>Taeniophyllum malianum</i> Schltr.		
<i>Taeniophyllum muelleri</i> Benth.		
<i>Taeniophyllum wilkianum</i> T.E.Hunt		

Queensland Species	Alternate Genus	Notes
<i>Tainia trinervis</i> (Blume) Rchb.f.		
<i>Thelasis carinata</i> Blume		
<i>Thelymitra angustifolia</i> R.Br.		
<i>Thelymitra carnea</i> R.Br.		
<i>Thelymitra fragrans</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra ixiooides</i> Sw.		
<i>Thelymitra longiloba</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra malvina</i> M.A.Clem.		
<i>Thelymitra nuda</i> R.Br.		
<i>Thelymitra pauciflora</i> R.Br.		
<i>Thelymitra purpurata</i> Rupp		
<i>Thelymitra queenslandica</i> Jeanes		
<i>Thelymitra</i> species 'Toy Creek'		no information available (Qld Herbarium record)
<i>Thrixspermum congestum</i> (F.M.Bailey) Dockrill		
<i>Thrixspermum platystachys</i> (F.M.Bailey) Schltr.		
<i>Trachoma speciosum</i> D.L.Jones, B.Gray, M.A.Clem. & J.J.Wood		
<i>Trachoma stellatum</i> M.A.Clem., D.L.Jones, B.Gray & J.J.Wood		
<i>Trachoma subluteum</i> (Rupp) Garay		
<i>Trichoglottis australiensis</i> Dockrill		
<i>Vanda hindsii</i> Lindl.		
<i>Vrydagzynea grayi</i> D.L.Jones & M.A.Clem.		
<i>Zeuxine oblonga</i> R.S.Rogers & C.T.White		

**Total: 494 taxa (species, subspecies, varieties)**

Queensland Hybrids	Name and Authority	Notes
<i>Dendrobium antennatum</i> X <i>discolor</i>		Jones 2006
<i>Dendrobium antennatum</i> X <i>nindii</i>		Jones 2006
<i>Dendrobium biggibum</i> X <i>discolor</i>	× <i>superbiens</i> Rchb.f.	
<i>Dendrobium biggibum</i> X × <i>superbiens</i>	× <i>lavarackianum</i> M.A. Clem.	
<i>Dendrobium calamiformis</i> X <i>nugentii</i>	× <i>grimesii</i> C.T. White & Summerh.	
<i>Dendrobium calamiformis</i> X <i>rigidum</i>	× <i>foederata</i> St. Cloud	
<i>Dendrobium canaliculatum</i> X <i>johannis</i>		
<i>Dendrobium canaliculatum</i> X <i>semifusum</i>		
<i>Dendrobium canaliculatum</i> X <i>discolor</i>		
<i>Dendrobium discolor</i> X <i>johannis</i>		Jones 2006
<i>Dendrobium discolor</i> X <i>lithicola</i>	× <i>vinicolor</i> St. Cloud	only marginally distinct from and probably the same as <i>D. ×superbiens</i>
<i>Dendrobium discolor</i> X <i>nindii</i>		
<i>Dendrobium discolor</i> X <i>semifusum</i>		
<i>Dendrobium gracilicaule</i> X <i>jonesii</i>		
<i>Dendrobium gracilicaule</i> X <i>speciosum</i>	× <i>gracillimum</i> (Rupp) Leaney	
<i>Dendrobium jonesii</i> X <i>nitidum</i>		Jones 2006
<i>Dendrobium jonesii</i> X <i>speciosum</i>	× <i>rupiosum</i> Clemesha	
<i>Dendrobium kingianum</i> subsp. <i>kingianum</i> X <i>speciosum</i> subsp. <i>speciosum</i>	× <i>delicatum</i> (F.M. Bailey) F.M. Bailey	
<i>Dendrobium nugentii</i> X <i>racemosum</i>		Jones 2006
<i>Diuria alba</i> X <i>chrysantha</i>		
<i>Diuris behrii</i> X <i>pardina</i>	× <i>palachila</i> R.S. Rogers	Qld record from Mt Moffat; neither parent occurs within several hundred km of location so identification doubtful
<i>Pterostylis baptistii</i> X <i>nutans</i>		
<i>Sarcochilus fitzgeraldii</i> X <i>hartmanii</i>		Jones 2006

**Total: 23 hybrids**

South Australian Species	Alternate Genus	Notes
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	
<i>Acianthus caudatus</i> variety <i>pallidus</i> Rupp	<i>Nemacianthus</i>	doubtfully distinct variety described from Tasmania but recognised in South Australia (Bates 2011)
<i>Acianthus pusillus</i> D.L.Jones		
<i>Caladenia argocalla</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia aurulenta</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	
<i>Caladenia behrii</i> Schldl.	<i>Arachnorchis</i>	
<i>Caladenia bicallata</i> R.S.Rogers subspecies <i>bicallata</i>	<i>Jonesiopsis</i>	
<i>Caladenia bicallata</i> R.S.Rogers subspecies <i>cleistogama</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia brumalis</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia calcicola</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia capillata</i> D.L.Jones	<i>Jonesiopsis</i>	
<i>Caladenia cardiochila</i> Tate	<i>Arachnorchis</i>	
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia carnea</i> variety <i>subulata</i> Nicholls	<i>Petalochilus</i>	recognised in one recent publication (Kuitert 2010) but probably only an aberrant form of <i>Caladenia carnea</i>
<i>Caladenia clavigera</i> A.Cunn. ex Lindl.	<i>Arachnorchis</i>	
<i>Caladenia clavula</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia coactilis</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia colorata</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia conferta</i> D.L.Jones subspecies <i>conferta</i>	<i>Arachnorchis</i>	
( <i>Caladenia</i> ) <i>conferta</i> D.L.Jones subspecies <i>occidentalis</i> R.J.Bates	<i>Arachnorchis</i>	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia cucullata</i> Fitzg.	<i>Stegostyla</i>	
<i>Caladenia dilatata</i> R.Br.	<i>Arachnorchis</i>	
( <i>Caladenia</i> ) <i>diversiflora</i> R.J.Bates	<i>Arachnorchis</i>	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia flaccida</i> D.L.Jones	<i>Jonesiopsis</i>	South Australian records unlikely to be true <i>Caladenia flaccida</i> ; here listed as an undescribed taxon <i>Caladenia</i> species 'Flinders Ranges red'
<i>Caladenia flindersica</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	
<i>Caladenia formosa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia fragrantissima</i> D.L.Jones & G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia fuliginosa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia gladiolata</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia interanea</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	
<i>Caladenia intuta</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia iridescens</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia latifolia</i> R.Br.		
<i>Caladenia leptochila</i> Fitzg. subspecies <i>dentata</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	
<i>Caladenia leptochila</i> Fitzg. subspecies <i>leptochila</i>	<i>Arachnorchis</i>	
<i>Caladenia littoricola</i> R.J.Bates	<i>Petalochilus</i>	
<i>Caladenia macroclavia</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia mentiens</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia microchila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia minor</i> Hook.f.	<i>Petalochilus</i>	a taxon of highly uncertain identity; Australian records from SA and NSW but generally considered to be endemic to New Zealand
<i>Caladenia moschata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia necrophylla</i> D.L.Jones	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia dilatata</i>
<i>Caladenia ornata</i> (Nicholls) D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia ovata</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia parva</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia phaeoclavia</i> D.L.Jones	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia parva</i>
<i>Caladenia prolata</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia pusilla</i> W.M.Curtis	<i>Petalochilus</i>	
<i>Caladenia pygmaea</i> (R.S.Rogers) R.J.Bates	<i>Petalochilus</i>	

South Australian Species	Alternate Genus	Notes
<i>Caladenia reticulata</i> Fitzg.	<i>Arachnorchis</i>	almost certainly occurs elsewhere in south-eastern Australia and more widely distributed than <i>reticulata</i> as narrowly defined by Jones (2006)
<i>Caladenia richardsiorum</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia rigida</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia sanguinea</i> D.L.Jones	<i>Jonesiopsis</i>	
<i>Caladenia saxatilis</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia septuosa</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia stricta</i> (R.Bates) R.Bates	<i>Arachnorchis</i>	
<i>Caladenia strigosa</i> (D.L.Jones) R.J.Bates	<i>Arachnorchis</i>	
<i>Caladenia subglabripHYlla</i> (R.J.Bates) M.A.Clem.	<i>Arachnorchis</i>	
<i>Caladenia tensa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia tentaculata</i> Schldl.	<i>Arachnorchis</i>	
<i>Caladenia toxochila</i> Tate	<i>Arachnorchis</i>	
<i>Caladenia transitoria</i> D.L.Jones subspecies 'Kangaroo Island'	<i>Stegostyla</i>	small aff. <i>transitoria</i> ; heathy woodland on limestone on Kangaroo Island; (R. Bates unpublished) subsp. <i>isolata</i> ms
<i>Caladenia valida</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia venusta</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia verrucosa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia villosissima</i> (G.W.Carr) Hopper & A.P.Br.	<i>Arachnorchis</i>	
( <i>Caladenia</i> ) <i>viriosa</i> R.J.Bates	<i>Arachnorchis</i>	described in <i>Arachnorchis</i> , new combination in <i>Caladenia</i> not yet made
<i>Caladenia vulgaris</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia woolcockiorum</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia xantholeuca</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia zephyra</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia</i> species 'coastal dunes'	<i>Arachnorchis</i>	aff. <i>orientalis</i> with large pale yellow flowers; coastal heath and heathy woodland, often on coastal dunes; sporadic between Portland (Vic) and Millicent (SA) (Backhouse 2011; Bates 2011)
<i>Caladenia</i> species 'coastal Eyre Peninsula'	<i>Arachnorchis</i>	aff. <i>macroclavia</i> with smaller flowers with more distinctly clubbed petals; heath and heathy woodland on limestone, coastal and hinterland Eyre Peninsula (Bates 2011)
<i>Caladenia</i> species 'Coonawarra'	<i>Arachnorchis</i>	aff. <i>colorata</i> with larger, variably coloured flowers with paler brown tails; heathy woodland between Tallageira (Vic), Lucindale and Penola (SA) (Jeanes & Backhouse 2006; Backhouse 2011); usually included in <i>Caladenia colorata</i>
<i>Caladenia</i> species 'Coorong'	<i>Petalochilus</i>	aff. <i>carnea</i> with brighter pink flowers with broader sepals and petals; coastal scrubs on alkaline sands in the Coorong region (Bates 2011)
<i>Caladenia</i> species 'Finnis'	<i>Arachnorchis</i>	extinct: the photo of this taxon in Bates (2011) suggests it was probably a hybrid, perhaps involving <i>Caladenia cardiochila</i> and <i>Caladenia parva</i>
<i>Caladenia</i> species 'Fleurieu Peninsula'	<i>Arachnorchis</i>	aff. <i>patersonii</i> with large white to pink to red flowers; heathy woodland on Fleurieu Peninsula (Bates 2011; Backhouse 2011)
<i>Caladenia</i> species 'Flinders Ranges red'	<i>Jonesiopsis</i>	aff. <i>filamentosa</i> with larger flowers with white-margined lateral sepals; dry woodland on rocky hills in the central Flinders Ranges (Bates 2011; Backhouse 2011); usually included in <i>Caladenia flaccida</i>
<i>Caladenia</i> species 'glowing green'	<i>Petalochilus</i>	aff. <i>prolata</i> with larger, more translucent flowers; seasonally damp River Red Gum flats in lower south-east SA & western Vic (Bates 2011)
<i>Caladenia</i> species 'Kangaroo Island'	<i>Stegostyla</i>	aff. <i>transitoria</i> with self-pollinating flowers that do not open and have very short barely distinguishable teeth on labellum mid-lobe; heathy woodland on Kangaroo Island (Bates 2011)
<i>Caladenia</i> species 'limestone mallee'	<i>Jonesiopsis</i>	aff. <i>capillata</i> with smaller flowers with very small labellum and relatively stiff petals and sepals; limestone mallee woodland and broombush on Eyre Peninsula; mentioned in NOSSA Feb 2015 (Bates unpublished)
<i>Caladenia</i> species 'little clowns'	<i>Arachnorchis</i>	aff. <i>lowanensis</i> with slightly smaller flowers; heathy woodland near Bordertown SA & western Vic (Bates 2011); probably falls within variation of <i>Caladenia lowanensis</i> (Backhouse 2011)
<i>Caladenia</i> species 'lowan mallee'	<i>Arachnorchis</i>	aff. <i>toxochila</i> with smaller flowers with shorter, thinner, more spaced calli and a more sharply curved column; mallee heath in Little Desert, also dry woodland near Rostron (Backhouse 2011); usually included in <i>Caladenia toxochila</i>
<i>Caladenia</i> species 'mallee'	<i>Arachnorchis</i>	small aff. <i>tensa</i> with more sparsely hairy stem and leaf and colourful flowers with almost no white on the labellum; widespread in mallee on sandy soils in SA, extending to west Vic (Bates 2011)
<i>Caladenia</i> species 'mallee heath'	<i>Arachnorchis</i>	aff. <i>arenaria</i> with lge cream flower with long dark red tails; mallee heath & woodland, Little Desert & Miram Piram (Backhouse 2011)
<i>Caladenia</i> species 'Monarto South'	<i>Jonesiopsis</i>	aff. <i>capillata</i> with shorter pale yellow sepals and petals; open mallee woodland on sand and clay loams over limestone, Murray Bridge to Middleton (Bates 2011)
<i>Caladenia</i> species 'Murray mallee'	<i>Arachnorchis</i>	aff. <i>stellata</i> with larger flowers with the labellum apex usually rolled under; mallee and dry woodland between Mt Pleasant and Langhorne Creek (Backhouse 2011; Bates 2011); usually included in <i>C. stellata</i>



South Australian Species	Alternate Genus	Notes
<i>Caladenia</i> species 'sand dunes'	<i>Petalochilus</i>	tall aff. <i>carnea</i> with multiple self-pollinating flowers, coastal scrubs on calcareous sand dunes (Bates 2011); <i>brevicaulis</i> ms
<i>Caladenia</i> species 'short yellow clubs'	<i>Arachnorchis</i>	aff. <i>aurulenta</i> with shorter, thicker sepaline clubs; mallee and dry woodland on the Eyre Peninsula (Bates 2011)
<i>Caladenia</i> species 'small tensa'	<i>Arachnorchis</i>	small aff. <i>tensa</i> ; no further information (R. Bates unpublished)
<i>Caladenia</i> species 'south east'	<i>Arachnorchis</i>	large aff. <i>reticulata</i> with yellow flowers; sandplains, flowering after fire (R. Bates unpublished)
<i>Caladenia</i> species 'tiny white'	<i>Petalochilus</i>	aff. <i>pusilla</i> with tiny (segments 5 mm long), short-lived (1 day), self-pollinating white flowers; damp shaded sites in woodland in the southern Mt Lofty Ranges and lower south-east (Bates 2011)
<i>Caladenia</i> species 'western Murray Flats'	<i>Arachnorchis</i>	unknown affinity; mentioned in NOSSA Feb 2015; Hartley to Sedan (R. Bates unpublished)
<i>Caleana major</i> R.Br.		
<i>Calochilus cupreus</i> R.S.Rogers		
<i>Calochilus herbaceus</i> Lindl.		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus pruinosus</i> D.L.Jones		
<i>Calochilus robertsonii</i> Benth.		
<i>Calochilus</i> species 'Kangaroo Island'		small. aff. <i>campestris</i> ; heathy woodland on limestone on Kangaroo Island (R. Bates unpublished)
<i>Chiloglottis cornuta</i> Hook.f.	<i>Simpliglottis</i>	
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmecchila</i>	
<i>Chiloglottis</i> species 'lowlands'	<i>Simpliglottis</i>	aff. <i>valida</i> indistinguishable from <i>valida</i> but with a different wasp pollinator (Bower 2007); widespread in damp shady sites in south-eastern Aust. below 1,000 m, mostly at low altitudes (Jeanes & Backhouse 2006)
<i>Corunastylis ciliata</i> (Ewart & B.Rees) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis despectans</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morrisii</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nigricans</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis tepperi</i> (F.Muell. ex. Tepper) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis</i> species 'Adelaide Hills'	<i>Genoplesium</i>	aff. <i>tasmanica</i> with dark purplish brown and green flowers with a broad labellum with a shallowly toothed, upturned margin; widespread in woodland and heath, often on rocky or gravelly sites (Bates 2011)
<i>Corunastylis</i> species 'dark flowers'	<i>Genoplesium</i>	aff. <i>rufa</i> with pointed dark reddish black labellum; widespread in dry woodland across inland south-eastern Australia (=C. sp. 'inland' in Backhouse <i>et al.</i> 2016; C. sp. 'dark midge' in Bates 2011)
<i>Corunastylis</i> species 'intermediate'	<i>Genoplesium</i>	aff. <i>rufa</i> with numerous crowded dark purplish brown flowers with contrasting pale (sometimes white-edged) petals; woodland and heath on calcareous soils (Bates 2011)
<i>Corunastylis</i> species 'large bracts'	<i>Genoplesium</i>	robust aff. <i>rufa</i> with numerous crowded maroon and green flowers with long sepals and long pointed floral bracts (most obvious in bud); <i>Triodia</i> hummocks on rocky ridges in dry woodland (Bates 2011)
<i>Corunastylis</i> species 'Mount Billy'	<i>Genoplesium</i>	aff. <i>rufa</i> with uniformly maroon flowers with a longish pointed labellum lacking any raised plates; seasonally wet heaths on clay soils on the Fleurieu Peninsula (Bates 2011)
<i>Corybas dentatus</i> D.L.Jones	<i>Corysanthes</i>	possibly of hybrid origin: <i>C. despectans</i> X <i>C. diemenicus</i>
<i>Corybas despectans</i> D.L.Jones & R.C.Nash	<i>Corysanthes</i>	
<i>Corybas diemenicus</i> (Lindl.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas dilatatus</i> (Rupp & Nicholls) Rupp	<i>Corysanthes</i>	
<i>Corybas expansus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	
( <i>Corybas</i> ) <i>minimus</i> (R.J.Bates)	<i>Corysanthes</i>	described in <i>Corysanthes</i> , new combination in <i>Corybas</i> not yet made
<i>Corybas unguiculatus</i> (R.Br.) Rchb.f.	<i>Anzybas</i>	
<i>Corybas</i> species 'red-eyed ghost'	<i>Corysanthes</i>	aff. <i>despectans</i> with greenish white, more upright flowers with a narrower dorsal sepal and a red column; calcareous sands under mallee and cypress-pines <i>Callitris</i> spp., Marion Bay, Fleurieu Peninsula and Kangaroo Island (Bates 2011)
<i>Corybas</i> species 'Woolly Tea-tree'	<i>Corysanthes</i>	late spring flowering aff. <i>diemenicus</i> with dark labellum; Woolly Tea-Tree swamps (Jeanes & Backhouse 2006 as sp. aff. <i>diemenicus</i> 1; Bates 2011 as sp. Silky Tea-tree swamps); <i>arcanus</i> ms

<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		
<i>Cyrtostylis reniformi</i> R.Br.		
<i>Cyrtostylis robusta</i> D.L.Jones & M.A.Clem.		
<i>Dipodium campanulatum</i> D.L.Jones		
<i>Dipodium pardalinum</i> D.L.Jones		
<i>Dipodium punctatum</i> (Sm.) R.Br.		
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		
<i>Diuris behrii</i> Schldl.		
<i>Diuris brevifolia</i> R.S.Rogers		
<i>Diuris calcicola</i> R.J.Bates		
<i>Diuris chryseopsis</i> D.L.Jones		
<i>Diuris daltonii</i> (C.Walter) D.L.Jones & M.A.Clem.		probably the same as <i>Diuris punctata</i> , from which it is poorly differentiated
<i>Diuris orientis</i> D.L.Jones		
<i>Diuris palustris</i> Lindl.		
<i>Diuris pardina</i> Lindl.		
<i>Diuris sulphurea</i> R.Br.		
<i>Diuris</i> species 'Red Gum waterholes'		short aff. <i>behrii</i> with small pale sometimes brown-streaked flowers with a depressed dorsal sepal and short labellum; damp, sometimes swampy sites in grassy woodland in the lower south-east (Bates 2011)
<i>Diuris</i> species 'sandhills'		aff. <i>chryseopsis</i> with narrower petals, labellum & narrow black-striped dorsal sepal; heathy woodland on grey sandy rises in the lower south-east of SA & far western Vic (Bates 2011)
<i>Eriochilus collinus</i> D.L. Jones & R.J.Bates		
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus paludosus</i> D.L. Jones & R.J.Bates		
<i>Eriochilus</i> species 'dry plains'		aff. <i>cucullatus</i> with soft deciduous hairs and minor ribbing on leaf, which is mostly green above, paler below; limestone plains of SA from west coast to Vic border (R. Bates unpublished)
<i>Eriochilus</i> species 'smooth leaf'		aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<i>Gastrodia procera</i> G.W.Carr		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Gastrodia vescula</i> D.L.Jones		
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Leporella fimbriata</i> (Lindl.) A.S.George		
<i>Leptoceras menziesii</i> (R.Br.) Lindl.	<i>Caladenia</i>	
<i>Microtis arenaria</i> Lindl.		
<i>Microtis atrata</i> Lindl.	<i>Microtidium</i>	
<i>Microtis eremaea</i> R.J.Bates		
<i>Microtis frutetorum</i> Schldl.		
<i>Microtis gracilentia</i> (R.J.Bates) R.J.Bates		
<i>Microtis oblonga</i> R.S.Rogers		
<i>Microtis orbicularis</i> R.S.Rogers	<i>Hydrorchis</i>	
<i>Microtis parviflora</i> R.Br.		
<i>Microtis rara</i> R.Br.		identity and distribution uncertain, thought to be only NSW & Qld (Jones 2006), although also claimed to be in Sth Aust (Bates 2011)
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		also on Lord Howe Island, Norfolk Island
<i>Microtis</i> species 'curved lip'		aff. <i>unifolia</i> with a blunt dorsal sepal and curved labellum; swampy habitats in central SA, flowering after fire; <i>curvichila</i> ms (Bates 2011)
<i>Microtis</i> species 'dense spikes'		aff. <i>parviflora</i> with a dense spike of numerous flowers with a tiny labellum; swampy habitats in sth Mt Lofty Range; <i>brevichila</i> ms (Bates 2011); probably only a subspecies of <i>parviflora</i>
<i>Microtis</i> species 'Flinders Ranges'		aff. <i>eremaea</i> with short slender spike of tiny congested blue-green flowers with fat rounded ovary; dry ridges of Flinders Ranges; <i>flindersica</i> ms (Bates 2011)
<i>Microtis</i> species 'fragrant'		aff. <i>arenaria</i> with sparser scented flowers with a longer labellum with a raised central ridge; widespread in south-eastern Australia; <i>fragrans</i> ms (Bates 2011)
<i>Microtis</i> species 'hanging bogs'		aff. <i>oblonga</i> with tall spike of well-spaced flowers with a broad dorsal sepal; peat bogs in the Adelaide Hills SA (Bates 2011); probably only a form of <i>oblonga</i>
<i>Microtis</i> species 'late flowering'		late flowering aff. <i>unifolia</i> with glossy flowers with a quadrate, unnotched labellum; woodland in higher parts of the Adelaide Hills; <i>postremus</i> ms (Bates 2011)
<i>Microtis</i> species 'narrow lip'		aff. <i>oblonga</i> with a long, narrow labellum constricted in the middle; tea-tree swamps on black alkaline mud in sth-east SA, also Vic?; <i>leptochila</i> ms (Bates 2011)

South Australian Species	Alternate Genus	Notes
<i>Microtis</i> species 'short leaf'		small slender aff. <i>unifolia</i> with short thick leaf & well spaced flowers with a long slender labellum; widespread; <i>brevis</i> ms (Bates 2011)
<i>Microtis</i> species 'Topperwein'		aff. <i>oblonga</i> with tall spike of larger, well-spaced fleshy flowers; River Red Gum swamps in sth-east SA & west Vic, flowering without fire (Bates 2011)
<i>Orthoceras strictum</i> R.Br.		
<i>Paracaleana disjuncta</i> D.L.Jones	<i>Caleana</i>	
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Pheladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	
<i>Prasophyllum album</i> R.S.Rogers		
<i>Prasophyllum asinantum</i> R.J.Bates		
<i>Prasophyllum australe</i> R.Br.		
<i>Prasophyllum calcicola</i> R.J.Bates		
<i>Prasophyllum catenemum</i> D.L.Jones		
<i>Prasophyllum collinum</i> D.L.Jones		
<i>Prasophyllum constrictum</i> R.S.Rogers		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum fecundum</i> R.J.Bates		
<i>Prasophyllum fitzgeraldii</i> R.S.Rogers & Maiden		
<i>Prasophyllum frenchii</i> F.Muell.		
<i>Prasophyllum goldsackii</i> J.Z. Weber & R.J.Bates		
<i>Prasophyllum laxum</i> R.J.Bates		
<i>Prasophyllum litorale</i> R.J.Bates		
<i>Prasophyllum murfetii</i> D.L.Jones		
<i>Prasophyllum occidentale</i> R.S.Rogers		
<i>Prasophyllum occultans</i> R.J.Bates		
<i>Prasophyllum odoratum</i> R.S.Rogers		
<i>Prasophyllum pallidum</i> Nicholls		
<i>Prasophyllum praecox</i> D.L.Jones		
<i>Prasophyllum pruinatum</i> R.S.Rogers		
<i>Prasophyllum rotundiflorum</i> R.S.Rogers		
<i>Prasophyllum spicatum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum validum</i> R.S.Rogers		
<i>Prasophyllum vesuvium</i> R.J.Bates		
<i>Prasophyllum viretrum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum</i> species 'Alligator Gorge'		aff. <i>fitzgeraldii</i> with slender flowers with a pale pink labellum; woodland in southern Flinders Ranges; probably extinct (Bates 2011)
<i>Prasophyllum</i> species 'bright shiny maroon'		aff. <i>fitzgeraldii</i> with bright shiny maroon flowers; woodland and mallee scrub on sandy soils between Keith and Loxton; <i>leiochilum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'brown lip'		aff. <i>fitzgeraldii</i> with brown lip with narrow white margin; heathy woodland in lower south-east of SA and far western Vic (Bates 2011; Jeanes & Backhouse 2006); <i>kafechilum</i> ms
<i>Prasophyllum</i> species 'bushfires'		slender aff. <i>constrictum</i> with variably coloured, strongly scented flowers with a long, slender, constricted labellum; woodland on rocky ridges in the Mount Lofty Ranges, flowering only after fire (Bates 2011); <i>pyrophilum</i> ms
<i>Prasophyllum</i> species 'coast sandhills'		coastal clumping aff. <i>odoratum</i> with upright labellum and thick-textured flowers; widely distributed in coastal heaths and grasslands (Bates 2011); <i>carinosum</i> ms
<i>Prasophyllum</i> species 'desert'		robust aff. <i>odoratum</i> with long narrow petals; semi-arid sandy and rocky grassy shrublands, often with <i>Triodia</i> ; widespread across central SA and far western NSW (Bates 2011); <i>eremaemum</i> ms
<i>Prasophyllum</i> species 'enigma'		aff. <i>fitzgeraldii</i> with partially opening, often contorted flowers with a green labellum callus and purple lamina; hilly woodland from central Eyre Peninsula to the southern Mt Lofty Ranges (Bates 2011); <i>aenigmum</i> ms
<i>Prasophyllum</i> species 'Gilbert River'		aff. <i>frenchii</i> with larger green and maroon flowers; seasonally moist grasslands on heavy clay soils in the Gilbert River valley in the northern Mt Lofty Ranges; last seen in 1990 and almost certainly extinct (Bates 2011)
<i>Prasophyllum</i> species 'Innes'		aff. <i>occidentale</i> with larger, more numerous flowers with a broad pinkish labellum lamina; coastal sedges and short open scrub on the tip of the Yorke Peninsula; <i>ventosum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'Jip Jip'		aff. <i>fitzgeraldii</i> with greenish white labellum; dry woodland on sandy soils between Keith and Padthaway, almost certainly extinct (Bates 2011)
<i>Prasophyllum</i> species 'late coastal dunes'		robust, late flowering aff. <i>odoratum</i> with large fleshy flowers and senescent leaf; coastal scrubs on calcareous sand dunes and flats (Bates 2011)

South Australian Species	Alternate Genus	Notes
<i>Prasophyllum</i> species 'lower south-east'		aff. <i>fitzgeraldii</i> with fleshy bright green flowers with a green labellum callus and off white lamina; winter wet heath on silty clays in the lower south-east, flowering after fire; <i>crassum</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'mallee sands'		aff. <i>odoratum</i> with narrow overlapping green and white flowers; mallee heath on sand dunes in the Big Desert (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> B Rouse)
<i>Prasophyllum</i> species 'Mokota'		short, slender aff. <i>constrictum</i> with small green flowers; montane grassland on bare hills above 500 m in the Mt Lofy Ranges; <i>arvense</i> ms (R. Bates unpublished)
<i>Prasophyllum</i> species 'needle points'		tall, robust aff. <i>occidentale</i> with pinkish toned flowers with long fine points to the sepals and reddish column wings; dry woodland under she-oaks <i>Allocasuarina</i> spp. on the eastern Eyre Peninsula; <i>aciculare</i> ms (Bates 2011)
<i>Prasophyllum</i> species 'Piccaninnie Ponds'		aff. <i>parviflorum</i> with short broad petals and sepals and a broad deeply grooved labellum; swampy sites in far south-east (Bates 2011)
<i>Prasophyllum</i> species 'pink lip'		aff. <i>fitzgeraldii</i> with pink labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> A Rouse); <i>rosaechilum</i> ms
<i>Prasophyllum</i> species 'purple lip'		aff. <i>fitzgeraldii</i> with purple labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> E Rouse); <i>violescens</i> ms
<i>Prasophyllum</i> species 'sedge swamps'		short aff. <i>odoratum</i> with few widely spaced flowers; heath and sedge swamps, often flowering in water (= sp. 'Drajurk' Backhouse <i>et al.</i> 2016; sp. 'Topperwein' Bates 2011)
<i>Prasophyllum</i> species 'Verran Tanks'		aff. <i>validum</i> with fleshy pink to brown-toned flowers; dry scrubs on sand plains and rocky hills on the eastern Eyre Peninsula between Cleve and Cowell (Bates 2011); <i>eyreanum</i> ms
<i>Prasophyllum</i> species 'waterholes'		very late flowering aff. <i>odoratum</i> with very widely spreading lateral sepals and incurved petals; heath and sedge swamps in south-western Vic & south-eastern SA (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> C/D Rouse 2002)
<i>Prasophyllum</i> species 'Wimmera woodlands'		short aff. <i>occidentale</i> with smaller, more widely spaced flowers with a narrower labellum; dry woodlands in narrow band between Murtoa (Vic) and Keith (SA)
<i>Pterostylis arenicola</i> M.A.Clem. & J.Stewart	<i>Oligochaetochilus</i>	
<i>Pterostylis biseta</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	
<i>Pterostylis boormanii</i> Rupp	<i>Oligochaetochilus</i>	
<i>Pterostylis bryophila</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis chlorogramma</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis cobarensis</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis concinna</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis cucullata</i> R.Br. subspecies <i>cucullata</i>		extinct in South Australia (Bates 2011)
<i>Pterostylis cucullata</i> R.Br. subspecies <i>sylvicola</i> D.L.Jones		
<i>Pterostylis curta</i> R.Br.		
<i>Pterostylis cycnocephala</i> Fitzg.	<i>Hymenochilus</i>	
<i>Pterostylis despectans</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis dolichochila</i> D.L.Jones & M.A.Clem.	<i>Diplodium</i>	
<i>Pterostylis erythroconcha</i> M.A.Clem. & D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis exalla</i> D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis excelsa</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis falcata</i> R.S.Rogers		
<i>Pterostylis ferruginea</i> (D.L.Jones) G.N.Backh.	<i>Oligochaetochilus</i>	
<i>Pterostylis flavovirens</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	
<i>Pterostylis foliata</i> Hook.f.		
( <i>Pterostylis</i> ) <i>gracilens</i> (D.L.Jones)	<i>Plumatichilos</i>	described in <i>Plumatichilos</i> , new combination in <i>Pterostylis</i> not yet made
<i>Pterostylis lepida</i> D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis lingua</i> M.A.Clem.	<i>Oligochaetochilus</i>	there is doubt as to the correct identity of South Australian records of this species
<i>Pterostylis littoralis</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	
<i>Pterostylis lustra</i> D.L.Jones		
<i>Pterostylis maxima</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis melagramma</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis mirabilis</i> D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis nana</i> R.Br.	<i>Linguella</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis ovata</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis parva</i> D.L.Jones & C.J.French	<i>Linguella</i>	

South Australian Species	Alternate Genus	Notes
<i>Pterostylis parviflora</i> R.Br.	<i>Speculantha</i>	
<i>Pterostylis pedunculata</i> R.Br.		
<i>Pterostylis planulata</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis prasina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis psammophila</i> D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis pusilla</i> R.S.Rogers	<i>Oligochaetochilus</i>	
<i>Pterostylis robusta</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis robusta</i> R.S.Rogers subspecies 'gorges'	<i>Diplodium</i>	distinctive tall form of <i>robusta</i> with dark red and white-striped flowers with longer dorsal sepal; woodland in gorges in Flinders & Mt Lofty Ranges (Bates 2011); subspecies <i>calvicola</i> ms
<i>Pterostylis sanguinea</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	
<i>Pterostylis setifera</i> M.A.Clem., Matthias & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis simulans</i> R.J.Bates	<i>Oligochaetochilus</i>	
<i>Pterostylis smaragdina</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis striata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis tasmanica</i> D.L.Jones	<i>Plumatichilos</i>	
<i>Pterostylis tenuissima</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis uliginosa</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis viriosa</i> (D.L.Jones) R.J.Bates	<i>Bunochilus</i>	
<i>Pterostylis xerophila</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis</i> species 'Adelaide hills'	<i>Diplodium</i>	aff. <i>alata</i> with smaller, narrower flowers; heathy and shrubby woodland in the Mt Lofty Range and possibly Kangaroo Island (Bates 2011)
<i>Pterostylis</i> species 'Arkaroola'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with smaller flowers with more rounded lateral sepals and a larger, broadly triangular labellum; rocky slopes and ledges in dry woodland in the northern Flinders Ranges (Bates 2011)
<i>Pterostylis</i> species 'Bimbowrie'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with dark brown flowers and a broadly triangular labellum; under low shrubs in sheltered sites on rocky hills north of Olary (Bates 2011)
<i>Pterostylis</i> species 'bluebush plains'	<i>Oligochaetochilus</i>	tall aff. <i>biseta</i> with narrower petals and lateral sepals; widespread across inland bluebush & saltbush plains; <i>thamnophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Broughton River'	<i>Oligochaetochilus</i>	small red & green aff. <i>biseta</i> with short lateral sepals and fleshy labellum with a grooved base; woodland on rocky hills in Broughton River gorge and southern Flinders Ranges; probably extinct (Bates 2011)
<i>Pterostylis</i> species 'canegrass'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with paler flowers with a more sparsely bristled labellum; grows amongst Triodia on red sand dunes on sandy plains between Renmark and Olary (Bates 2011)
<i>Pterostylis</i> species 'chunky flowers'	<i>Diplodium</i>	aff. <i>dolichocheila</i> with a shorter labellum; coastal scrubs from Kingston (SA) to Portland (Vic) (Bates 2011; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'coastal SA'	<i>Linguella</i>	aff. <i>nana</i> with large raised crinkled basal leaves and large yellow-green flowers aging to brownish; coastal scrubs over limestone SA; <i>paracites</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Everard Range'	<i>Oligochaetochilus</i>	winter flowering aff. <i>excelsa</i> with extremely long free points to the lateral sepals; dry woodland on rocky hills, almost certainly extinct (Bates 2011)
<i>Pterostylis</i> species 'Eyre Peninsula'	<i>Hymenochilus</i>	aff. <i>cynnocephala</i> with short habit, small striped flowers with prominent labellum appendage & crenulate bracts; mallee and dry woodland on the Eyre Peninsula; <i>cymbella</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Flinders'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> with smaller drooping flowers on long slender pedicels; mallee & <i>Callitris</i> woodland on rocky hills in Flinders Ranges between Hawker & Narrina (Bates 2011)
<i>Pterostylis</i> species 'Flinders Ranges north 1'	<i>Oligochaetochilus</i>	unknown affiliation, mentioned in NOSSA Journal 39(11) Dec 2015
<i>Pterostylis</i> species 'Flinders Ranges north 2'	<i>Oligochaetochilus</i>	unknown affiliation, mentioned in NOSSA Journal 39(11) Dec 2015
<i>Pterostylis</i> species 'Gammon Range'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> with larger flowers with a narrower labellum and very long sepal free points; <i>Callitris</i> woodland and dry shrubland in the northern Flinders Ranges from Mt Fitton to Hawker (Bates 2011)
<i>Pterostylis</i> species 'Joanna'	<i>Diplodium</i>	aff. <i>alata</i> with smaller, narrower flowers with a small opening; heathy woodland near Narracoorte (Bates 2011)
<i>Pterostylis</i> species 'large striped flowers'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with large brownish striped flowers, mallee, <i>Callitris</i> woodland and dry woodland from Charlton (Vic) to Renmark (SA) (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'limestone'	<i>Hymenochilus</i>	aff. <i>cynnocephala</i> with prominent labellum appendage and distinctly striped flowers; woodland and scrubs on clay and sandy soils over limestone on the Yorke Peninsula and Murray mallee region; <i>calvicola</i> ms (Bates 2011)
<i>Pterostylis</i> species 'lower Murray'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with darker flowers with forward curved lateral sepals free points and a more rounded labellum; <i>Callitris</i> woodland on rocky areas along the lower Murray River, probably extinct (Bates 2011)
<i>Pterostylis</i> species 'mallee heath'	<i>Plumatichilos</i>	smallish glossy aff. <i>plumosa</i> with long pedicel and long dense labellum hairs; mallee heath on sandy to clay loam soils in western Vic & sth-eastern SA (Bates 2011; Jeanes & Backhouse 2006)

South Australian Species	Alternate Genus	Notes
<i>Pterostylis</i> species 'mallee snail'	<i>Linguella</i>	short aff. <i>nana</i> with small squat flowers with short lateral sepals; mallee on sandy to clay loam soils (Backhouse et al. 2016; Bates 2011) <i>tenuis</i> ms
<i>Pterostylis</i> species 'Mambray Creek'	<i>Bunochilus</i>	aff. <i>chlorogramma</i> with a broadly rectangular labellum; dry woodland at Mambray Creek (Jones & Clements 2006)
<i>Pterostylis</i> species 'Mokota'	<i>Hymenochilus</i>	aff. <i>cycnocephala</i> with short, stocky habit, thick fleshy leaves and barely striped flowers; montane grassland near Mt Bryan; <i>anemophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Mount Geharty'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> with slender flowers with long free points to the sepals; dry shrubland on rocky outcrops on the Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'Mount Olinthus'	<i>Oligochaetochilus</i>	short aff. <i>excelsa</i> with rigid flowers with a small basal rosette and an anvil-shaped labellum; dry shrubland on rocky ridges on the Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'Mount Victoria Uranium Mine'	<i>Oligochaetochilus</i>	dwarf aff. <i>boormanii</i> with smaller green flowers with flatter, hooked lateral sepals; sparse shrublands on granite hills north of Olary (Bates 2011)
<i>Pterostylis</i> species 'Newland Head'	<i>Diplodium</i>	aff. <i>dolichochila</i> with a more widely opening flowers, a protruding sinus and shorter labellum; coastal scrubs on Kangaroo Island and the lower south-east (Bates 2011)
<i>Pterostylis</i> species 'Nora Creina'	<i>Diplodium</i>	aff. <i>alata</i> with tricolored flowers, a long fine point on dorsal sepal and a distinct gap between lateral sepals and galea; coastal scrubs in the lower south-east SA and sth-west Vic (Bates 2011)
<i>Pterostylis</i> species 'Old Boolcoolmata'	<i>Oligochaetochilus</i>	aff. <i>lingua</i> with larger, dark maroon flowers with a larger labellum; Callitris woodland along drainage lines in rocky hills north of Olary (Bates 2011); <i>lingua</i> subspecies <i>ampla</i> ms
<i>Pterostylis</i> species 'Oratan Rock'	<i>Oligochaetochilus</i>	short aff. <i>excelsa</i> with very small flowers with hooked lateral sepals; shrublands on rocky hills near Lillydale (Bates 2011)
<i>Pterostylis</i> species 'Outback'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with broader, rounded lateral sepals; sparse shrublands on rocky hills between Renmark and Olary (SA) and near Broken Hill (NSW) (Bates 2011)
<i>Pterostylis</i> species 'quartz'	<i>Oligochaetochilus</i>	aff. <i>lingua</i> with thick, fleshy labellum; sparse shrublands with <i>Triodia</i> on rocky quartz ridges near Olary (Bates 2011)
<i>Pterostylis</i> species 'rock ledges'	<i>Oligochaetochilus</i>	aff. <i>aciculiformis</i> with greenish flowers with longer free points to the sepals; open woodland on rocky hills in the upper south-east and Adelaide Hills (Bates 2011)
<i>Pterostylis</i> species 'rocky ranges'	<i>Hymenochilus</i>	aff. <i>mutica</i> with short stocky habit and crowded, barely striped flowers; shrubland and dry woodland on rocky ranges in central SA; <i>pagophila</i> ms (Bates 2011)
<i>Pterostylis</i> species 'sandheath'	<i>Speculantha</i>	aff. <i>rubescens</i> from heathy woodland near Glencoe (Bates 2011); possibly the same as <i>Pterostylis rubescens</i> as the differences are unclear
<i>Pterostylis</i> species 'slender desert'	<i>Oligochaetochilus</i>	small, slender aff. <i>excelsa</i> with pale flowers with short free points to the sepals; sparse shrubland on red soils around granite boulders near Olary (Bates 2011)
<i>Pterostylis</i> species 'southern Eyre Peninsula'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> with a raised rosette of fleshy, crinkled strongly veined leaves and darker flowers with a broader labellum; sparse shrublands on rocky outcrops in the southern Eyre Peninsula (Bates 2011)
<i>Pterostylis</i> species 'tall mallee snail'	<i>Linguella</i>	aff. <i>nana</i> with tall hairless stem and narrow green flowers with long light brownish sepal tips; inland mallee and dry woodland (Bates 2011); <i>naniflora</i> ms
<i>Pterostylis</i> species 'tall purple'	<i>Diplodium</i>	aff. <i>dolichochila</i> with darker flowers with a narrower labellum; coastal scrubs and heaths in the south-east and possibly Kangaroo Island (Bates 2011)
<i>Pterostylis</i> species 'tall woodland'	<i>Oligochaetochilus</i>	tall aff. <i>biseta</i> with long pedicels, narrow galea, distinctive lateral sepals with a double bend and a much broader labellum; woodland on fertile soils in the Mount Lofty and Flinders Ranges (Bates 2011); <i>sigmoidea</i> ms
<i>Pterostylis</i> species 'tiny flowers'	<i>Hymenochilus</i>	aff. <i>mutica</i> with tall slender habit and small, spaced barely striped flowers; widely distributed in shrublands and woodland; <i>pisinna</i> ms (Bates 2011)
<i>Pterostylis</i> species 'upper Spencer Gulf'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> ; no other information (NOSSA Journal 37(9): 94 (2013))
<i>Pterostylis</i> species 'Warren snail'	<i>Linguella</i>	aff. <i>nana</i> with pale, fleshy, smooth, ovate leaves, a hairless stem, swollen stem bracts and inflated base to the flower; open woodland in the Adelaide Hills (Bates 2011)
<i>Pterostylis</i> species 'woodland'	<i>Plumatichilos</i>	aff. <i>plumosa</i> with ovate, bluntly pointed rosette leaves; widespread in woodland, the common form of <i>plumosa</i> in South Australia (Bates 2011)
<i>Pterostylis</i> species 'woodlands'	<i>Hymenochilus</i>	tall aff. <i>cycnocephala</i> with striped, spaced flowers; widespread in woodlands across SA and in west Vic; <i>nemoralis</i> ms (Bates 2011)
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		
<i>Spiranthes alticola</i> D.L.Jones		very difficult to distinguish from (and doubtfully distinct from) <i>Spiranthes australis</i>
<i>Spiranthes elytra</i> D.L.Jones & R.J.Bates		
<i>Thelymitra albiflora</i> Jeanes		
<i>Thelymitra alcockii</i> D.L.Jones ex Jeanes		
<i>Thelymitra antennifera</i> (Lindl.) Hook.f.		
<i>Thelymitra arenaria</i> Lindl.		
<i>Thelymitra aristata</i> Lindl.		
<i>Thelymitra azurea</i> R.S.Rogers		
<i>Thelymitra batesii</i> Jeanes		
<i>Thelymitra benthamiana</i> Rchb.f.		
<i>Thelymitra bracteata</i> J.Z.Weber ex Jeanes		



<i>Thelymitra brevifolia</i> Jeanes	
<i>Thelymitra brevifolia</i> Jeanes subspecies 'Porters'	has a paler, non-cleft column apex than <i>brevifolia</i> ; possibly a stabilised <i>T. albiflora</i> x <i>T. brevifolia</i> hybrid? (Bates 2011)
<i>Thelymitra carnea</i> R.Br.	
<i>Thelymitra circumsepta</i> Fitzg.	
<i>Thelymitra corrugata</i> R.J.Bates	
<i>Thelymitra crenulata</i> R.J.Bates	
<i>Thelymitra cyanapicata</i> Jeanes	
<i>Thelymitra cyanea</i> (Lindl.)	
<i>Thelymitra epipactoides</i> F. Muell.	
<i>Thelymitra exigua</i> Jeanes	
<i>Thelymitra flexuosa</i> Endl.	
<i>Thelymitra glaucophylla</i> Jeanes	
<i>Thelymitra grandiflora</i> Fitzg. subspecies <i>exposa</i> R.J.Bates	
<i>Thelymitra grandiflora</i> Fitzg. subspecies <i>grandiflora</i>	
<i>Thelymitra holmesii</i> Nicholls	
<i>Thelymitra hygrophila</i> R.J.Bates	
<i>Thelymitra inflata</i> Jeanes	
<i>Thelymitra ixioides</i> Sw.	
<i>Thelymitra juncifolia</i> Lindl.	
<i>Thelymitra latifolia</i> R.J.Bates	
<i>Thelymitra lucida</i> Jeanes	
<i>Thelymitra luteocilium</i> Fitzg.	
<i>Thelymitra malvina</i> M.A.Clem.	
<i>Thelymitra matthewsii</i> Cheesem.	
<i>Thelymitra mucida</i> Fitzg.	
<i>Thelymitra nuda</i> R.Br.	
<i>Thelymitra occidentalis</i> Jeanes	
<i>Thelymitra odora</i> R.J.Bates	
<i>Thelymitra odora</i> R.J.Bates subspecies 'horse manure'	unnamed subsp. of <i>odora</i> with strong horse manure-like odour (Bates 2011)
<i>Thelymitra orientalis</i> R.J.Bates	
<i>Thelymitra pallidifructus</i> R.J.Bates	
<i>Thelymitra pauciflora</i> R.Br.	
<i>Thelymitra peniculata</i> Jeanes	
<i>Thelymitra polychroma</i> D.L.Jones & M.A.Clem.	
<i>Thelymitra rubra</i> Fitzg.	
<i>Thelymitra rubricaulis</i> R.J.Bates	
<i>Thelymitra</i> species 'Claire'	short aff. <i>pauciflora</i> with two stem bracts and dark brown column apex with a black collar; woodlands in the Adelaide hills (Bates 2011)
<i>Thelymitra</i> species 'Glen Shera'	tiny aff. <i>polychroma</i> with a purplish truncate post-anther lobe, sparse spots and tiny hair tufts; damp sites in heathland on acid sands on the Fleurieu Peninsula and lower South East (Bates 2011)
<i>Thelymitra</i> species 'insipid'	very slender, late flowered aff. <i>pauciflora</i> with long narrow sheathing leaf, rarely opening dull blue flowers and flat red and yellow post-anther lobe; widely distributed in many habitats in cooler, higher rainfall areas (Bates 2011)
<i>Thelymitra</i> species 'Meadows'	aff. <i>pauciflora</i> with scabrid leaves and very rounded flowers with a truncate post-anther lobe; grassy woodland on steep rocky slopes in the southern Mount Lofty Ranges (Bates 2011)
<i>Thelymitra</i> species 'papillose'	small aff. <i>ixioides</i> with papillose post-anther lobe and mop-like hair tufts; heath on damp peaty sands in the lower south-east (Bates 2011)
<i>Thelymitra</i> species 'red gum swamps'	aff. <i>inflata</i> with more circular and narrower column apex; seasonal red gum swamps (SA, Vic) or wet forest (Tas) (Bates 2011)
<i>Thelymitra</i> species 'Topperwein'	aff. <i>holmesii</i> with horseshoe-shaped, rounded column apex and yellow hair tufts; sedge swamps in River Red Gum woodland in the lower south-east of SA and western Vic (Bates 2011)

**Total: 378 taxa (species, subspecies, varieties)**

South Australian Hybrids	Name and Authority	Notes
<i>Caladenia argocalla</i> X <i>tentaculata</i>	Bates 2011	
<i>Caladenia aurulenta</i> X <i>clavula</i>	Backhouse 2011	
<i>Caladenia behrii</i> X <i>reticulata</i>	Bates 2011	
<i>Caladenia behrii</i> X <i>rigida</i>	Bates 2011	
<i>Caladenia bicallata</i> X <i>capillata</i>	G. Backhouse personal observation	
<i>Caladenia brumalis</i> X <i>capillata</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>cardiochila</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>conferta</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>fuliginosa</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>latifolia</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>macroclavia</i>	Bates 2011	
<i>Caladenia brumalis</i> X <i>stricta</i>	Bates 2011	
<i>Caladenia calcicola</i> X <i>cardiochila</i>	Bates 2011	
<i>Caladenia calcicola</i> X <i>colorata</i>	Bates 2011	
<i>Caladenia capillata</i> X <i>toxochila</i>	Bates 2011	
<i>Caladenia capillata</i> X <i>zephyra</i>	G. Backhouse personal observation	
<i>Caladenia cardiochila</i> X <i>colorata</i>	Backhouse 2011	
<i>Caladenia cardiochila</i> X <i>reticulata</i>	plants on Kangaroo Island recognised in Bates (2011) as a species	
<i>Caladenia cardiochila</i> X <i>septuosa</i>	G. Backhouse personal observation	
<i>Caladenia cardiochila</i> X <i>stricta</i>	Backhouse <i>et al.</i> 2016	
<i>Caladenia cardiochila</i> X <i>strigosa</i>	Bates 2011	
<i>Caladenia cardiochila</i> X <i>xanthochila</i>	Backhouse 2011	
<i>Caladenia cardiochila</i> X <i>zephyra</i>	Bates 2011	
<i>Caladenia cardiochila</i> X species 'mallee heath'	Bates 2011	
<i>Caladenia carnea</i> X <i>fuscata</i>	Jones <i>et al.</i> 1999	
<i>Caladenia carnea</i> X <i>latifolia</i>	Bates 2011	
<i>Caladenia carnea</i> X <i>vulgaris</i>	Bates 2011	
<i>Caladenia clavigera</i> X <i>reticulata</i>	G. Backhouse personal observation	
<i>Caladenia clavula</i> X <i>stricta</i>	G. Backhouse personal observation	
<i>Caladenia clavula</i> X <i>zephyra</i>	Backhouse 2011	
<i>Caladenia clavula</i> X species 'short yellow clubs'	Backhouse 2011	
<i>Caladenia colorata</i> X <i>strigosa</i>	Bates 2011	
<i>Caladenia colorata</i> X <i>tensa</i>	Bates 2011	
<i>Caladenia colorata</i> X <i>verrucosa</i>	Bates 2011	
<i>Caladenia colorata</i> X species 'mallee heath'	Backhouse <i>et al.</i> 2016	
<i>Caladenia colorata</i> X species 'Murray mallee'	Bates 2011	
<i>Caladenia conferta</i> X <i>macroclavia</i>	Bates 2011	
<i>Caladenia flindersica</i> X <i>gladiolata</i>	<i>xensigera</i> D.L. Jones	
<i>Caladenia flindersica</i> X <i>saxatilis</i>	Backhouse 2011	
<i>Caladenia flindersica</i> X <i>tentaculata</i>	Backhouse 2011	
<i>Caladenia flindersica</i> X <i>toxochila</i>	Bates 2011	
<i>Caladenia fuliginosa</i> X <i>stricta</i>	Bates 2011	
<i>Caladenia fulva</i> X species 'little clowns'	Bates 2011	
<i>Caladenia gladiolata</i> X <i>saxatilis</i>	Bates 2011	
<i>Caladenia gladiolata</i> X <i>viriosa</i>	Bates 2011	
<i>Caladenia interanea</i> X <i>zephyra</i>	G. Backhouse personal observation	
<i>Caladenia leptochila</i> X <i>reticulata</i>	Bates 2011	
<i>Caladenia macroclavia</i> X <i>zephyra</i>	G. Backhouse personal observation	
<i>Caladenia necrophylla</i> X <i>venusta</i>	Bates 2011	
<i>Caladenia parva</i> X <i>stricta</i>	Backhouse 2011	
<i>Caladenia reticulata</i> X <i>rigida</i>	Bates 2011	
<i>Caladenia saxatilis</i> X <i>woolcockiorum</i>	Bates 2011	
<i>Caladenia septuosa</i> X <i>zephyra</i>	Bates 2012	
<i>Caladenia stricta</i> X <i>zephyra</i>	Bates 2011	
<i>Caladenia stricta</i> X species 'little clowns'	Bates 2011	
<i>Caladenia tentaculata</i> X <i>venusta</i>	Jeanes & backhouse 2006	
<i>Caladenia toxochila</i> X <i>zephyra</i>	Bates 2011	
<i>Caladenia toxochila</i> X <i>viriosa</i>	Bates 2011	
<i>Caladenia verrucosa</i> X species 'little clowns'	Bates 2011	

South Australian Hybrids	Name and Authority	Notes
<i>Caladenia verrucosa</i> X <i>viriosa</i>		Bates 2011
<i>Caladenia woolcockiorum</i> X <i>viriosa</i>		Bates 2011
<i>Caladenia zephyra</i> X <i>viriosa</i>		Backhouse 2011
<i>Caladenia zephyra</i> X species 'short yellow clubs'		Bates 2011
<i>Caladenia capillata</i> X <i>Pheladenia deformis</i>		Bates 2011
<i>Caladenia carnea</i> X <i>Pheladenia deformis</i>		Bates 2011
<i>Caladenia coactilis</i> X <i>Pheladenia deformis</i>		Bates 2011
<i>Caladenia latifolia</i> X <i>Pheladenia deformis</i>		Bates 2011
<i>Corunastylis ciliata</i> X <i>despectans</i>		Bates 2011
<i>Corybas despectans</i> X <i>diemenicus</i>		Bates 2011
<i>Corybas despectans</i> X <i>expansus</i>		Bates 2011
<i>Corybas despectans</i> X <i>incurvus</i>		Bates 2011
<i>Corybas dilatatus</i> X <i>incurvus</i>	× <i>miscellus</i> D.L. Jones	
<i>Dipodium pardalinum</i> X <i>roseum</i>		Bates 2011; Backhouse <i>et al.</i> 2016
<i>Diuris behrii</i> X <i>orientis</i>		Bates 2011
<i>Diuris behrii</i> X <i>palustris</i>		Bates 2011
<i>Diuris behrii</i> X <i>pardina</i>	× <i>palachila</i> R.S. Rogers	Bates 2011; Jeanes & Backhouse 2006
<i>Diuris behrii</i> X <i>sulphurea</i>		Bates 2011
<i>Diuris brevifolia</i> X <i>orientis</i>		Bates 2011
<i>Diuris brevifolia</i> X <i>pardina</i>		Bates 2011
<i>Diuris calcicola</i> X <i>orientis</i>		Bates 2016
<i>Diuris calcicola</i> X <i>palustris</i>		Bates 2016
<i>Diuris chrysiopsis</i> X <i>pardina</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Diuris orientis</i> X <i>palustris</i>		Bates 2011
<i>Diuris orientis</i> X <i>pardina</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Diuris palustris</i> X <i>pardina</i>		Bates 2011
<i>Glossodia major</i> X <i>Pheladenia deformis</i>	× <i>Glossadenia tutelata</i> Kavulak	
<i>Pterostylis cucullata</i> X <i>nutans</i>		Bates 2011
<i>Pterostylis curta</i> X <i>lustra</i>		Bates 2011
<i>Pterostylis curta</i> X <i>pedunculata</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Pterostylis dolichochila</i> X <i>erythroconcha</i>		Bates 2011
<i>Pterostylis falcata</i> X <i>nutans</i>	× <i>ingens</i> (Rupp) D.L. Jones	
<i>Pterostylis lustra</i> X <i>nutans</i>		Bates 2011
<i>Pterostylis nutans</i> X <i>pedunculata</i>		Bates 2011
<i>Pterostylis robusta</i> X species 'Adelaide hills'		Bates 2011
<i>Pterostylis</i> species 'limestone' X species 'woodlands'		Bates 2011 ( <i>calcicola</i> ms X <i>nemoralis</i> ms)
<i>Thelymitra albiflora</i> X <i>antennifera</i>		Jeanes 2004; Bates 2011
<i>Thelymitra albiflora</i> X <i>bracteata</i>		Bates 2011
<i>Thelymitra albiflora</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra albiflora</i> X <i>pallidifructus</i>		Bates 2011
<i>Thelymitra albiflora</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra alcockiae</i> X <i>antennifera</i>		Backhouse <i>et al.</i> 2016
<i>Thelymitra alcockiae</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra antennifera</i> X <i>bracteata</i>		Bates 2011
<i>Thelymitra antennifera</i> X <i>inflata</i>		Bates 2011
<i>Thelymitra antennifera</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra antennifera</i> X <i>megcalyptera</i>		Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>nuda</i>	× <i>macmillanii</i> F. Muell.	Bates 2011
<i>Thelymitra antennifera</i> X <i>pauciflora</i>		Bates 2011
<i>Thelymitra antennifera</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra arenaria</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra azurea</i> X <i>nuda</i>		Jones 2006
<i>Thelymitra bracteata</i> X <i>ixioides</i>		Bates 2011
<i>Thelymitra bracteata</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra bracteata</i> X <i>nuda</i>		Bates 2011
<i>Thelymitra brevifolia</i> X <i>juncifolia</i>		Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra brevifolia</i> X <i>ixioides</i>		Bates 2011
<i>Thelymitra brevifolia</i> X <i>nuda</i>		Bates 2011
<i>Thelymitra brevifolia</i> X <i>pauciflora</i>		Bates 2011

South Australian Hybrids	Name and Authority	Notes
<i>Thelymitra carnea</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra corrugata</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra crenulata</i> X <i>ixioides</i>		Bates 2011
<i>Thelymitra crenulata</i> X <i>malvina</i>		Bates 2011
<i>Thelymitra crenulata</i> X <i>peniculata</i>		Bates 2011
<i>Thelymitra cyanapicata</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra cyanapicata</i> X <i>nuda</i>		Bates 2011
<i>Thelymitra cyanapicata</i> X <i>pauciflora</i>		Bates 2011
<i>Thelymitra cyanapicata</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra glaucopylla</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra holmesii</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra holmesii</i> X <i>pauciflora</i>		Bates 2011
<i>Thelymitra hygrophila</i> X <i>odora</i>		Bates 2011
<i>Thelymitra inflata</i> X <i>juncifolia</i>		Bates 2011
<i>Thelymitra inflata</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra ixioides</i> X <i>mucida</i>		Jones 2006
<i>Thelymitra ixioides</i> X <i>pallidifructus</i>		Bates 2010
<i>Thelymitra ixioides</i> X <i>pauciflora</i>	x <i>truncata</i> R.S. Rogers	
<i>Thelymitra ixioides</i> X <i>peniculata</i>	x <i>merraniae</i> Nicholls	
<i>Thelymitra juncifolia</i> X <i>mucida</i>		Bates 2011
<i>Thelymitra juncifolia</i> X <i>nuda</i>		Bates 2011
<i>Thelymitra juncifolia</i> X <i>pauciflora</i>		Bates 2011; Jones <i>et al.</i> 2008
<i>Thelymitra juncifolia</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra juncifolia</i> X species 'pale buds'		Bates 2011
<i>Thelymitra luteocilium</i> X <i>megcalyptera</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Thelymitra luteocilium</i> X <i>nuda</i>	x <i>chasmogama</i> R.S. Rogers	Bates 2011
<i>Thelymitra luteocilium</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra nuda</i> X <i>pallidifructus</i>		Bates 2010
<i>Thelymitra nuda</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra odora</i> subspecies 'horse manure' X <i>rubra</i>		Bates 2011
<i>Thelymitra pauciflora</i> X <i>rubra</i>		Bates 2011

**Total: 149 hybrids**

Tasmanian Species	Alternate Genus	Notes
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	
<i>Acianthus caudatus</i> variety <i>pallidus</i> Rupp	<i>Nemacianthus</i>	doubtfully distinct variety described from Tasmania but recognised in South Australia (Bates 2011)
<i>Acianthus pusillus</i> D.L.Jones		
<i>Burnettia cuneata</i> Lindl.		
<i>Caladenia alata</i> R.Br.	<i>Petalochilus</i>	
<i>Caladenia alpina</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia angustata</i> Lindl.	<i>Stegostyla</i>	
<i>Caladenia anthracina</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia atrata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia atrochila</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia aurantiaca</i> (R.S.Rogers) Rupp	<i>Petalochilus</i>	in Tasmania only on Deal Island Bass Strait
<i>Caladenia australis</i> G.W.Carr	<i>Arachnorchis</i>	in Tasmania only on Flinders Island Bass Strait; no recent records
<i>Caladenia brachyscapa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia campbellii</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia cardiochila</i> Tate	<i>Arachnorchis</i>	in Tasmania only on Flinders Island Bass Strait; no recent records
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia caudata</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia clavigera</i> A.Cunn. ex Lindl.	<i>Arachnorchis</i>	
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia cracens</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia dienema</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia dilatata</i> R.Br.	<i>Arachnorchis</i>	
<i>Caladenia echidnachila</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia gracilis</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia helvina</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia latifolia</i> R.Br.		
<i>Caladenia lindleyana</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia mentiens</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia pallida</i> Lindl.	<i>Arachnorchis</i>	possibly extinct
<i>Caladenia patersonii</i> R.Br.	<i>Arachnorchis</i>	
<i>Caladenia prolata</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia pusilla</i> W.M.Curtis	<i>Petalochilus</i>	
<i>Caladenia saggicola</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia sylvicola</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia tonellii</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia transitoria</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia vulgaris</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caleana major</i> R.Br.		
<i>Calochilus campestris</i> R.Br.		
<i>Calochilus herbaceus</i> Lindl.		
<i>Calochilus imberbis</i> R.S.Rogers		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Chiloglottis cornuta</i> Hook.f.	<i>Simpliglottis</i>	
<i>Chiloglottis grammata</i> G.W.Carr	<i>Simpliglottis</i>	
<i>Chiloglottis gunnii</i> Lindl.	<i>Simpliglottis</i>	
<i>Chiloglottis reflexa</i> (Labill.) Druce		
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmechila</i>	
<i>Chiloglottis triceratops</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis</i> species 'lowlands'	<i>Simpliglottis</i>	aff. <i>valida</i> indistinguishable from <i>valida</i> but with a different wasp pollinator (Bower 2007); widespread in damp shady sites in south-eastern Aust. below 1,000 m, mostly at low altitudes (Jeanes & Backhouse 2006)
<i>Chiloglottis</i> species 'Weilangta'	<i>Simpliglottis</i>	no information available (Tas. Herbarium record)
<i>Corunastylis archeri</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis brachystachya</i> (Lindl.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	

Tasmanian Species	Alternate Genus	Notes
<i>Corunastylis despectans</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis firthii</i> (L.Cady) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morrisii</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nuda</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nudiscapa</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis pumila</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis tasmanica</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corybas aconitiflorus</i> Salisb.		
<i>Corybas diemenicus</i> (Lindl.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas dienemus</i> D.L.Jones	<i>Nematoceras</i>	Macquarie Island only
<i>Corybas fimbriatus</i> (R.Br.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	
<i>Corybas sulcatus</i> (M.A.Clem. & D.L.Jones) G.N.Backh.	<i>Nematoceras</i>	Macquarie Island only
<i>Corybas unguiculatus</i> (R.Br.) Rchb.f.	<i>Anzybas</i>	
<i>Cryptostylis leptochila</i> Benth.		
<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		
<i>Cyrtostylis reniformi</i> R.Br.		
<i>Cyrtostylis robusta</i> D.L.Jones & M.A.Clem.		
<i>Dendrobium striolatum</i> Rchb.f. subspecies <i>chrysanthum</i> D.L.Jones	<i>Dockrillia</i>	
<i>Dendrobium striolatum</i> Rchb.f. Rauschert subspecies <i>striolatum</i>	<i>Dockrillia</i>	
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		
<i>Diuris chryseopsis</i> D.L.Jones		
<i>Diuris lanceolata</i> Lindl.		
<i>Diuris monticola</i> D.L.Jones		
<i>Diuris orientis</i> D.L.Jones		
<i>Diuris palustris</i> Lindl.		
<i>Diuris pardina</i> Lindl.		
<i>Diuris sulphurea</i> R.Br.		
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Gastrodia procera</i> G.W.Carr		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Leptoceras menziesii</i> (R.Br.) Lindl.	<i>Caladenia</i>	
<i>Lyperanthus suaveolens</i> R.Br.		
<i>Microtis arenaria</i> Lindl.		
<i>Microtis atrata</i> Lindl.	<i>Microtidium</i>	
<i>Microtis oblonga</i> R.S.Rogers		
<i>Microtis orbicularis</i> R.S.Rogers	<i>Hydrorchis</i>	
<i>Microtis parviflora</i> R.Br.		
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		
<i>Orthoceras strictum</i> R.Br.		
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Pheladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	
<i>Prasophyllum alpinum</i> R.Br.		
<i>Prasophyllum amoenum</i> D.L.Jones		
<i>Prasophyllum apoxychilum</i> D.L.Jones		
<i>Prasophyllum atratum</i> D.L.Jones		
<i>Prasophyllum australe</i> R.Br.		
<i>Prasophyllum brevilabre</i> (Lindl.) Hook.f.		
<i>Prasophyllum castaneum</i> D.L.Jones		
<i>Prasophyllum concinnum</i> Nicholls		
<i>Prasophyllum crebriflorum</i> D.L.Jones		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum favonium</i> D.L.Jones		
<i>Prasophyllum flavum</i> R.Br.		



Tasmanian Species	Alternate Genus	Notes
<i>Prasophyllum incorrectum</i> D.L.Jones		
<i>Prasophyllum incurvum</i> D.L.Jones		
<i>Prasophyllum limnetes</i> D.L.Jones		
<i>Prasophyllum lindleyanum</i> Rchb.f.		
<i>Prasophyllum milfordense</i> D.L.Jones		
<i>Prasophyllum mimulum</i> D.L.Jones		
<i>Prasophyllum olidum</i> D.L.Jones		
<i>Prasophyllum perangustum</i> D.L.Jones		
<i>Prasophyllum pulchellum</i> D.L.Jones		
<i>Prasophyllum robustum</i> (Nicholls) M.A.Clem. & D.L.Jones		
<i>Prasophyllum rostratum</i> Lindl.		
<i>Prasophyllum secutum</i> D.L.Jones		
<i>Prasophyllum sphacelatum</i> D.L.Jones		
<i>Prasophyllum stellatum</i> D.L.Jones		
<i>Prasophyllum tadgellianum</i> R.S.Rogers		
<i>Prasophyllum taphanyx</i> D.L.Jones		
<i>Prasophyllum truncatum</i> Lindl.		
<i>Prasophyllum tunbridgense</i> D.L.Jones		
<i>Prasophyllum</i> species 'Arthurs Lake'		aff. <i>montanum</i> ; grows in the ecotone between wet grassland and woodland on rocky ground above 700 m information available; previously called <i>P. montanum</i> in Tasmania (Jones <i>et al.</i> 1999)
<i>Pterostylis alata</i> (Labill.) Rchb.f.	<i>Diplodium</i>	
<i>Pterostylis aphylla</i> Lindl.	<i>Specularantha</i>	
<i>Pterostylis atrans</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis atriola</i> D.L.Jones	<i>Specularantha</i>	
<i>Pterostylis commutata</i> D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis concinna</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis cucullata</i> R.Br. subspecies <i>cucullata</i>		
<i>Pterostylis curta</i> R.Br.		
<i>Pterostylis decurva</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis dubia</i> R.Br.		
<i>Pterostylis falcata</i> R.S.Rogers		
<i>Pterostylis foliata</i> Hook.f.		
<i>Pterostylis furcata</i> Lindl.		
<i>Pterostylis grandiflora</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis lustra</i> D.L.Jones		
<i>Pterostylis melagramma</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis nana</i> R.Br.	<i>Linguella</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis parviflora</i> R.Br.	<i>Specularantha</i>	
<i>Pterostylis pedoglossa</i> Fitzg.	<i>Crangonorchis</i>	
<i>Pterostylis pedunculata</i> R.Br.		
<i>Pterostylis plumosa</i> Cady	<i>Plumatichilos</i>	
<i>Pterostylis pratensis</i> D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis rubenachii</i> D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis sanguinea</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	
<i>Pterostylis scabrida</i> Lindl.		
<i>Pterostylis squamata</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis stenochila</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis tasmanica</i> D.L.Jones	<i>Plumatichilos</i>	
<i>Pterostylis tunstallii</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis uliginosa</i> D.L.Jones	<i>Specularantha</i>	
<i>Pterostylis wapstrarum</i> D.L.Jones	<i>Hymenochilus</i>	also spelt <i>wapstreorum</i>
<i>Pterostylis williamsonii</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis ziegeleri</i> D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis</i> species 'Flinders Island'	<i>Bunochilus</i>	aff. <i>chlorogramma</i> with a broadly rectangular labellum; heathy woodland on Flinders Island (Jones & Clements 2006)
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		

Tasmanian Species	Alternate Genus	Notes
<i>Sarcochilus australis</i> (Lindl.) Rchb.f.		
<i>Spiranthes alticola</i> D.L.Jones		very difficult to distinguish from (and doubtfully distinct from) <i>Spiranthes australis</i>
<i>Spiranthes australis</i> (R.Br.) Lindl.		
<i>Thelymitra aggericola</i> D.L.Jones		
<i>Thelymitra antennifera</i> (Lindl.) Hook.f.		
<i>Thelymitra arenaria</i> Lindl.		
<i>Thelymitra aristata</i> Lindl.		
<i>Thelymitra atronitida</i> Jeanes		
<i>Thelymitra benthamiana</i> Rchb.f.		
<i>Thelymitra bracteata</i> J.Z.Weber ex Jeanes		
<i>Thelymitra brevifolia</i> Jeanes		
<i>Thelymitra carnea</i> R.Br.		
<i>Thelymitra circumsepta</i> Fitzg.		
<i>Thelymitra cyanea</i> (Lindl.)		
<i>Thelymitra erosa</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra exigua</i> Jeanes		
<i>Thelymitra flexuosa</i> Endl.		
<i>Thelymitra holmesii</i> Nicholls		
<i>Thelymitra imbricata</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra improcera</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra inflata</i> Jeanes		
<i>Thelymitra ixioides</i> Sw.		
<i>Thelymitra jonesii</i> Jeanes		
<i>Thelymitra juncifolia</i> Lindl.		
<i>Thelymitra longiloba</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra lucida</i> Jeanes		
<i>Thelymitra malvina</i> M.A.Clem.		
<i>Thelymitra mucida</i> Fitzg.		
<i>Thelymitra nuda</i> R.Br.		
<i>Thelymitra pauciflora</i> R.Br.		
<i>Thelymitra peniculata</i> Jeanes		
<i>Thelymitra polychroma</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra rubra</i> Fitzg.		
<i>Thelymitra silena</i> D.L.Jones		
<i>Thelymitra simulata</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra spadicea</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra sparsa</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra versicolor</i> Lindl.		often treated as a synonym of <i>T. nuda</i> , although considered a distinct species by Jones & Clements (1998) and Jones <i>et al.</i> (1999)
<i>Thelymitra viridis</i> Jeanes		
<i>Thelymitra</i> species 'red gum swamps'		aff. <i>inflata</i> with more circular and narrower column apex; seasonal red gum swamps (SA, Vic) or wet forest (Tas) (Bates 2011)
<i>Thynniorchis huntianus</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	
<i>Thynniorchis nothofagicola</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	
<i>Townsonia viridis</i> (Hook.f.) Schltr.		

**Total: 210 taxa (species, subspecies)**

Tasmanian Hybrids	Name and Authority	Notes
<i>Caladenia caudata</i> X <i>saggicola</i>		Backhouse 2011
<i>Glossodia major</i> X <i>Pheladenia deformis</i>	× <i>Glossadenia tutelata</i> Kavulak	
<i>Pterostylis concinna</i> X <i>striata</i>	× <i>toveyana</i> Ewart & Sharman	
<i>Pterostylis dubia</i> X <i>scabrada</i>		Jones 2006
<i>Pterostylis falcata</i> X <i>nutans</i>	× <i>ingens</i> (Rupp) D.L. Jones	
<i>Thelymitra carnea</i> X <i>ixioides</i>	× <i>irregularis</i> Nicholls	
<i>Thelymitra ixioides</i> X <i>pauciflora</i>	× <i>truncata</i> R.S. Rogers	
<i>Thelymitra ixioides</i> X <i>peniculata</i>	× <i>merraniae</i> Nicholls	

**Total: 8 hybrids**

Victorian Species	Alternate Genus	Notes
<i>Acianthus caudatus</i> R.Br.	<i>Nemacianthus</i>	
<i>Acianthus collinus</i> D.L.Jones		
<i>Acianthus exsertus</i> R.Br.		
<i>Acianthus pusillus</i> D.L.Jones		
<i>Burnettia cuneata</i> Lindl.		
<i>Caladenia aestiva</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia alata</i> R.Br.	<i>Petalochilus</i>	
<i>Caladenia alpina</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia amoena</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia ampla</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia reticulata</i>
<i>Caladenia ancylosa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia atrovespa</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia audasii</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia aurantiaca</i> (R.S.Rogers) Rupp	<i>Petalochilus</i>	
<i>Caladenia australis</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia bicallata</i> R.S.Rogers subspecies <i>bicallata</i>	<i>Jonesiopsis</i>	
<i>Caladenia brachyscapa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia calcicola</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia capillata</i> D.L.Jones	<i>Jonesiopsis</i>	
<i>Caladenia cardiochila</i> Tate	<i>Arachnorchis</i>	
<i>Caladenia carnea</i> R.Br.	<i>Petalochilus</i>	a widespread, highly variable taxon incorporating several named varieties that are not generally recognised, but likely to consist of several distinct species
<i>Caladenia carnea</i> variety <i>subulata</i> Nicholls	<i>Petalochilus</i>	recognised in one recent publication (Kuitert 2010) but probably only an aberrant form of <i>Caladenia carnea</i>
<i>Caladenia catenata</i> (Sm.) Druce	<i>Petalochilus</i>	
<i>Caladenia clarkiae</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia clavescens</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia clavigera</i> A.Cunn. ex Lindl.	<i>Arachnorchis</i>	
<i>Caladenia cleistantha</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia colorata</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia concolor</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia congesta</i> R.Br.	<i>Stegostyla</i>	
<i>Caladenia cremna</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia cretacea</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia cruciformis</i> D.L.Jones	<i>Arachnorchis</i>	said to have once occurred in South Australia but now extinct there (Bates 2011)
<i>Caladenia cucullata</i> Fitzg.	<i>Stegostyla</i>	
<i>Caladenia dilatata</i> R.Br.	<i>Arachnorchis</i>	
<i>Caladenia douglasiorum</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia filamentosa</i> R.Br.	<i>Jonesiopsis</i>	
<i>Caladenia flavovirens</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia formosa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia fragrantissima</i> D.L.Jones & G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia fulva</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia fuscata</i> (Rchb.f.) M.A.Clem. & D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia grampiana</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia hastata</i> (Nicholls) Rupp	<i>Arachnorchis</i>	
<i>Caladenia hildae</i> E.E.Pescott & Nicholls	<i>Stegostyla</i>	
<i>Caladenia insularis</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia iridescens</i> R.S.Rogers	<i>Stegostyla</i>	
<i>Caladenia latifolia</i> R.Br.		
<i>Caladenia lowanensis</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia magnifica</i> (Nicholls) D.L.Jones & G.W.Carr	<i>Arachnorchis</i>	probably extinct
<i>Caladenia maritima</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia mentiens</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia montana</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia moschata</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia oenochila</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia oreophila</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia orientalis</i> (G.W.Carr) Hopper & A.P.Br.	<i>Arachnorchis</i>	

Victorian Species	Alternate Genus	Notes
<i>Caladenia ornata</i> (Nicholls) D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia osmera</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	distinguishable from <i>Caladenia montana</i> only by altitude
<i>Caladenia parva</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia peisleyi</i> (D.L.Jones) G.N.Backh.	<i>Arachnorchis</i>	
<i>Caladenia phaeoclavia</i> D.L.Jones	<i>Arachnorchis</i>	doubtfully distinct from and probably a synonym of <i>Caladenia parva</i>
<i>Caladenia pilotensis</i> D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia praecox</i> Nicholls	<i>Stegostyla</i>	
<i>Caladenia prolata</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia pumila</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia pusilla</i> W.M.Curtis	<i>Petalochilus</i>	
<i>Caladenia robinsonii</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia rosella</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia stricta</i> (R.Bates) R.Bates	<i>Arachnorchis</i>	
<i>Caladenia tensa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia tentaculata</i> Schldl.	<i>Arachnorchis</i>	
<i>Caladenia tessellata</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia thysanochila</i> G.W.Carr	<i>Arachnorchis</i>	probably extinct
<i>Caladenia transitoria</i> D.L.Jones	<i>Stegostyla</i>	
<i>Caladenia valida</i> (Nicholls) M.A.Clem. & D.L.Jones	<i>Arachnorchis</i>	
<i>Caladenia venusta</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia verrucosa</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia versicolor</i> G.W.Carr	<i>Arachnorchis</i>	
<i>Caladenia villosissima</i> (G.W.Carr) Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia vulgaris</i> D.L.Jones	<i>Petalochilus</i>	
<i>Caladenia xanthochila</i> D.Beardsell & C.Beardsell	<i>Arachnorchis</i>	
<i>Caladenia</i> species 'Ararat hills'	<i>Arachnorchis</i>	late winter flowering aff. <i>venusta</i> with smaller flowers, short marginal teeth and sparser calli; Ararat Hills (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'Buchan'	<i>Arachnorchis</i>	large aff. <i>montana</i> with longer slender sepals and petals with long indistinct clubs and more numerous marginal teeth; open forest near Buchan (Turner <i>et al.</i> 2014)
<i>Caladenia</i> species 'Carboor'	<i>Arachnorchis</i>	aff. <i>concolor</i> with dull red flowers; grassy open forest in nth-eastern Victoria (Backhouse 2011); possibly the same as <i>Caladenia clavescens</i>
<i>Caladenia</i> species 'central Victoria'	<i>Arachnorchis</i>	aff. <i>formosa</i> with smaller, dull maroon flowers with long filamentous tails; dry woodland in central Vic (Jeanes & Backhouse 2006; Backhouse 2011)
<i>Caladenia</i> species 'Chapple Vale'	<i>Stegostyla</i>	summer-flowering aff. <i>iridescens</i> with golden yellow flowers; Otway Ranges (Jeanes & Backhouse 2006); probably only a colour form of <i>Caladenia iridescens</i>
<i>Caladenia</i> species 'coastal dunes'	<i>Arachnorchis</i>	aff. <i>orientalis</i> with large pale yellow flowers; coastal heath and heathy woodland, often on coastal dunes; sporadic between Portland (Vic) and Millicent (SA) (Backhouse 2011; Bates 2011)
<i>Caladenia</i> species 'Coonawarra'	<i>Arachnorchis</i>	aff. <i>colorata</i> with larger, variably coloured flowers with paler brown tails; heathy woodland between Tallageira (Vic), Lucindale and Penola (SA) (Jeanes & Backhouse 2006; Backhouse 2011); usually included in <i>Caladenia colorata</i>
<i>Caladenia</i> species 'Earlston'	<i>Arachnorchis</i>	aff. <i>concolor</i> with glossy red flowers with dark red labellum & thickened tepal tails; dry woodland on low hills at Earlston (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'glowing green'	<i>Petalochilus</i>	aff. <i>prolata</i> with larger, more translucent flowers; seasonally damp River Red Gum flats in lower south-east SA & western Vic (Bates 2011)
<i>Caladenia</i> species 'Inverleigh'	<i>Arachnorchis</i>	aff. <i>patersonii</i> with large cream to pinkish flowers; open heathy woodland on sandy loam at Inverleigh (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'Kaniva'	<i>Petalochilus</i>	aff. <i>pusilla</i> with 1-5 pink flowers with cupped tepals; dry heathy woodland near Kaniva (Backhouse 2011)
<i>Caladenia</i> species 'Kaniva Giant'	<i>Jonesiopsis</i>	aff. <i>filamentosa</i> with large dark red flowers with long drooping segments; mallee heath from Kiata to Kaniva (Backhouse 2011)
<i>Caladenia</i> species 'Kilsyth South'	<i>Arachnorchis</i>	aff. <i>venusta</i> with smaller dull white flowers with narrower labellum; heathy open forest at Kilsyth South (Backhouse & Jeanes 1995) <i>consternans</i> ms
<i>Caladenia</i> species 'Kooyera'	<i>Arachnorchis</i>	aff. <i>venusta</i> with smaller flower with dark reddish black glandular tails; grassy woodland on sandy loam near Rheola (Backhouse 2011)
<i>Caladenia</i> species 'large scented'	<i>Petalochilus</i>	aff. <i>carnea</i> with large, pale, strongly scented flowers; hilly dry woodland between Ararat and Stuart Mill (Backhouse 2011)
<i>Caladenia</i> species 'little clowns'	<i>Arachnorchis</i>	aff. <i>lowanensis</i> with slightly smaller flowers; heathy woodland near Bordertown SA & western Vic (Bates 2011); probably falls within variation of <i>Caladenia lowanensis</i> (Backhouse 2011)
<i>Caladenia</i> species 'lowan mallee'	<i>Arachnorchis</i>	aff. <i>toxochila</i> with smaller flowers with shorter, thinner, more spaced calli and a more sharply curved column; mallee heath in Little Desert, also dry woodland near Rostron (Backhouse 2011); usually included in <i>Caladenia toxochila</i>

Victorian Species	Alternate Genus	Notes
<i>Caladenia</i> species 'lower Glenelg River'	<i>Arachnorchis</i>	aff. <i>fragrantissima</i> with variable cream to red flowers; heathy woodland on red clays over limestone on Glenelg River near Nelson (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'mallee'	<i>Arachnorchis</i>	small aff. <i>tensa</i> with more sparsely hairy stem and leaf and colourful flowers with almost no white on the labellum; widespread in mallee on sandy soils in SA, extending to west Vic (Bates 2011)
<i>Caladenia</i> species 'mallee heath'	<i>Arachnorchis</i>	aff. <i>arenaria</i> with lge cream flower with long dark red tails; mallee heath & woodland, Little Desert & Miram Piram (Backhouse 2011)
<i>Caladenia</i> species 'Mandurang'	<i>Arachnorchis</i>	aff. <i>fragrantissima</i> with large stiff white flowers with reddish calli and marginal teeth; dry sparsely shrubby woodland near Mandurang (Backhouse & Jeanes 1995)
<i>Caladenia</i> species 'Raymond Island'	<i>Arachnorchis</i>	aff. <i>ancylosa</i> with relatively large clubbed greenish flower with long marginal teeth; heathy woodland on Raymond Island (Jeanes & Backhouse 2006)
<i>Caladenia</i> species 'south Gippsland'	<i>Arachnorchis</i>	early flowering aff. <i>australis</i> with large flowers with long indistinct clubs; moist forest in south Gippsland (Backhouse 2011); usually included in <i>Caladenia australis</i>
<i>Caladenia</i> species 'southern slopes'	<i>Arachnorchis</i>	aff. <i>australis</i> with large flowers with long indistinct sepal clubs and shorter petal clubs and has a different wasp pollinator; moist foothill forest in Gippsland from Tonimbuk to Erica (Backhouse 2011)
<i>Caladenia</i> species 'St Andrews'	<i>Arachnorchis</i>	aff. <i>oenochila</i> with large pale flowers; shrubby woodland on gravelly clay soil at St Andrews Vic (Jeanes & Backhouse 2006); probably only a form of <i>Caladenia oenochila</i>
<i>Caladenia</i> species 'swamp heath'	<i>Arachnorchis</i>	aff. <i>venusta</i> with small creamy white flowers with dark tails; swampy heath west of Casterton, flowering mostly after fires (Backhouse 2011)
<i>Caladenia</i> species 'Yarram'	<i>Petalochilus</i>	aff. <i>catenata</i> with variable white to crimson flowers; heathy open forest on sandy loam near Yarram (Jeanes & Backhouse 2006); most likely a hybrid swarm involving <i>Caladenia carnea</i> and <i>Caladenia catenata</i>
<i>Caleana major</i> R.Br.		
<i>Calochilus campestris</i> R.Br.		
<i>Calochilus herbaceus</i> Lindl.		
<i>Calochilus imberbis</i> R.S.Rogers		
<i>Calochilus paludosus</i> R.Br.		
<i>Calochilus platyichilus</i> D.L.Jones		virtually indistinguishable from and probably a synonym of <i>Calochilus robertsonii</i>
<i>Calochilus pruinosus</i> D.L.Jones		
<i>Calochilus richiae</i> Nicholls		
<i>Calochilus robertsonii</i> Benth.		
<i>Calochilus saprophyticus</i> R.S.Rogers		
<i>Calochilus therophilus</i> D.L.Jones		
<i>Calochilus</i> species 'Moondarra'		small, colourful aff. <i>campestris</i> with short irregular basal plates and short very sparse bristles on labellum; wet heathy woodland near Erica (Backhouse <i>et al.</i> 2016)
<i>Chiloglottis cornuta</i> Hook.f.	<i>Simpliglottis</i>	
<i>Chiloglottis jeanesii</i> D.L.Jones	<i>Simpliglottis</i>	
<i>Chiloglottis reflexa</i> (Labill.) Druce		
<i>Chiloglottis seminuda</i> D.L.Jones		
<i>Chiloglottis trapeziformis</i> Fitzg.	<i>Myrmechila</i>	
<i>Chiloglottis trilabra</i> Fitzg.		
<i>Chiloglottis valida</i> D.L.Jones	<i>Simpliglottis</i>	'true' <i>valida</i> occurs in the mountains above 500 m but is morphologically indistinguishable from two related cryptic species (Bower 2007; see sp. 'Falls Creek' and sp. 'lowlands'), distribution overlaps with sp. 'Falls Creek'
<i>Chiloglottis</i> species 'highlands'	<i>Simpliglottis</i>	aff. <i>valida</i> with slightly smaller labellum with fewer calli and different wasp pollinator; montane damp forest and woodland above 800 m (Bower 2007; Jones <i>et al.</i> 2008); indistinguishable from <i>C. valida</i>
<i>Chiloglottis</i> species 'lowlands'	<i>Simpliglottis</i>	aff. <i>valida</i> indistinguishable from <i>valida</i> but with a different wasp pollinator (Bower 2007); widespread in damp shady sites in south-eastern Aust. below 1,000 m, mostly at low altitudes (Jeanes & Backhouse 2006)
<i>Corunastylis archeri</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	'true' <i>archeri</i> almost certainly does not occur in NSW & Qld, with these plants probably comprising a complex of several different, un-named species
<i>Corunastylis arrecta</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis ciliata</i> (Ewart & B.Rees) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis densa</i> (Fitzg.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis despectans</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis morrisii</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis nuda</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	



Victorian Species	Alternate Genus	Notes
<i>Corunastylis nudiscapa</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis ostrina</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis pumila</i> (Hook.f.) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis tepperi</i> (F.Muell. ex. Tepper) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis turfosa</i> (D.L.Jones) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corunastylis</i> species 'Beechworth'	<i>Genoplesium</i>	aff. <i>archeri</i> with narrower, bluntly pointed labellum with curved but not upturned apex; sparsely shrubby woodland on granite hills at Beechworth (Backhouse <i>et al.</i> 2016)
<i>Corunastylis</i> species 'dark flowers'	<i>Genoplesium</i>	aff. <i>rufa</i> with pointed dark reddish black labellum; widespread in dry woodland across inland south-eastern Australia (= <i>C.</i> sp. 'inland' in Backhouse <i>et al.</i> 2016; <i>C.</i> sp. 'dark midge' in Bates 2011)
<i>Corunastylis</i> species 'Gippsland Plains'	<i>Genoplesium</i>	aff. <i>rufa</i> with fewer, slightly smaller paler flowers with a broader labellum and more broadly triangular petals; heathy open forest Bairnsdale to Stratford (Jeanes & Backhouse 2006)
<i>Corunastylis</i> species 'Pyrete Range'	<i>Genoplesium</i>	tall, robust aff. <i>morrisii</i> with more flowers in a denser spike, a broader labellum and lateral sepals pouched at the base; heathy woodland in Otway & Pyrete Ranges (Jeanes & Backhouse 2006); probably a form of <i>C. morrisii</i>
<i>Corybas aconitiflorus</i> Salisb.		
<i>Corybas despectans</i> D.L.Jones & R.C.Nash	<i>Corysanthes</i>	
<i>Corybas diemenicus</i> (Lindl.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas fimbriatus</i> (R.Br.) Rchb.f.	<i>Corysanthes</i>	
<i>Corybas fordhamii</i> (Rupp) Rupp	<i>Anzybas</i>	
<i>Corybas grumulus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas hispidus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas incurvus</i> D.L.Jones & M.A.Clem.	<i>Corysanthes</i>	
<i>Corybas unguiculatus</i> (R.Br.) Rchb.f.	<i>Anzybas</i>	
<i>Corybas</i> species 'dark labellum'	<i>Corysanthes</i>	small aff. <i>diemenicus</i> with dark labellum; heathy woodland on Wilsons Promontory (Jeanes & Backhouse 2006); possibly <i>C. grumulus</i> but not seen for over a decade now
<i>Corybas</i> species 'Portland'	<i>Corysanthes</i>	aff. <i>diemenicus</i> with smaller, pale flower with fewer marginal teeth; coastal heath near Portland (= sp. aff. <i>diemenicus</i> 3 Jeanes & Backhouse 2006); possibly just a mutant form of <i>C. diemenicus</i>
<i>Corybas</i> species 'Wilson's Promontory'	<i>Corysanthes</i>	aff. <i>diemenicus</i> with smaller, darker flower with fewer marginal teeth; damp heathy woodland at Wilsons Promontory (= sp. aff. <i>diemenicus</i> 2 Jeanes & Backhouse 2006); possibly a misidentification of the named hybrid <i>C. xmiscellus</i>
<i>Corybas</i> species 'Woolly Tea-tree'	<i>Corysanthes</i>	late spring flowering aff. <i>diemenicus</i> with dark labellum; Woolly Tea-Tree swamps (Jeanes & Backhouse 2006 as sp. aff. <i>diemenicus</i> 1; Bates 2011 as sp. Silky Tea-tree swamps); <i>arcanus</i> ms
<i>Cryptostylis erecta</i> R.Br.		
<i>Cryptostylis hunteriana</i> Nicholls		
<i>Cryptostylis leptochila</i> Benth.		
<i>Cryptostylis subulata</i> (Labill.) Rchb.f.		
<i>Cyanicula caerulea</i> (R.Br.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyrtostylis reniformi</i> R.Br.		
<i>Cyrtostylis robusta</i> D.L.Jones & M.A.Clem.		
<i>Dendrobium speciosum</i> Sm. subspecies <i>speciosum</i> variety <i>speciosum</i>	<i>Thelychiton</i>	
<i>Dendrobium striolatum</i> Rchb.f. Rauschert subspecies <i>striolatum</i>	<i>Dockrillia</i>	
<i>Dipodium campanulatum</i> D.L.Jones		
<i>Dipodium hamiltonianum</i> F.M.Bailey		
<i>Dipodium pardalinum</i> D.L.Jones		
<i>Dipodium punctatum</i> (Sm.) R.Br.		
<i>Dipodium roseum</i> D.L.Jones & M.A.Clem.		
<i>Dipodium variegatum</i> M.A.Clem. & D.L.Jones		
<i>Diuris basaltica</i> D.L.Jones		
<i>Diuris behrii</i> Schldl.		
<i>Diuris chryseopsis</i> D.L.Jones		
<i>Diuris daltonii</i> (C.Walter) D.L.Jones & M.A.Clem.		probably the same as <i>Diuris punctata</i> , from which it is poorly differentiated
<i>Diuris dendrobioides</i> Fitzg.		
<i>Diuris fragrantissima</i> D.L.Jones & M.A.Clem.		

Victorian Species	Alternate Genus	Notes
<i>Diuris fucosa</i> D.L.Jones		
<i>Diuris gregaria</i> D.L.Jones		
<i>Diuris monticola</i> D.L.Jones		
<i>Diuris ochroma</i> D.L.Jones		
<i>Diuris orientis</i> D.L.Jones		
<i>Diuris palustris</i> Lindl.		
<i>Diuris pardina</i> Lindl.		
<i>Diuris protena</i> D.L.Jones		
<i>Diuris punctata</i> Sm. variety <i>punctata</i>		
<i>Diuris semilunulata</i> Messmer		
<i>Diuris subalpina</i> D.L.Jones		
<i>Diuris sulphurea</i> R.Br.		
<i>Diuris tricolor</i> Fitzg.		
<i>Diuris</i> species 'Gippsland plains'		small aff. <i>dendrobioides</i> with blotched flowers; plains grasslands in central Gippsland (Jeanes & Backhouse 2006)
<i>Diuris</i> species 'sandhills'		aff. <i>chryseopsis</i> with narrower petals, labellum & narrow black-striped dorsal sepal; heathy woodland on grey sandy rises in the lower south-east of SA & far western Vic (Bates 2011)
<i>Diuris</i> species 'thick sepals'		late flowering aff. <i>pardina</i> with shorter, thicker, less curved to straight lateral sepals; dry woodland and open forest in western Vic & probably SA (Jeanes & Backhouse 2006); may be same as <i>D. calcicola</i> R.J. Bates
<i>Eriochilus cucullatus</i> (Labill.) Rchb.f.		<i>E. cucullatus</i> sens.strict. is said to be restricted to Tasmania, with up to 5 cryptic species occurring in south-eastern Australia (Bates 2011)
<i>Eriochilus magenteus</i> D.L.Jones		
<i>Eriochilus</i> species 'smooth leaf'		aff. <i>cucullatus</i> with vertical petals and a smooth green leaf; said to be the common widespread taxon occurring on mainland south-eastern Australia (Bates 2011)
<i>Gastrodia procera</i> G.W.Carr		
<i>Gastrodia sesamoides</i> R.Br.		
<i>Glossodia major</i> R.Br.	<i>Caladenia</i>	
<i>Glossodia minor</i> R.Br.	<i>Caladenia</i>	alternative name <i>Caladenia minorata</i> M.A. Clem.
<i>Leporella fimbriata</i> (Lindl.) A.S.George		
<i>Leptoceras menziesii</i> (R.Br.) Lindl.	<i>Caladenia</i>	
<i>Lyperanthus suaveolens</i> R.Br.		
<i>Microtis arenaria</i> Lindl.		
<i>Microtis atrata</i> Lindl.	<i>Microtidium</i>	
<i>Microtis oblonga</i> R.S.Rogers		
<i>Microtis orbicularis</i> R.S.Rogers	<i>Hydrochis</i>	
<i>Microtis parviflora</i> R.Br.		
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		
<i>Microtis</i> species 'alpine'		aff. <i>unifolia</i> with fewer smaller spaced flowers with a smaller labellum; wet mountain meadows and woodland (Jones <i>et al.</i> 2008)
<i>Microtis</i> species 'fragrant'		aff. <i>arenaria</i> with sparser, scented flowers with a longer labellum with a raised central ridge; widespread in many habitats in south-eastern Australia; <i>fragrans</i> ms (Bates 2011)
<i>Microtis</i> species 'narrow lip'		aff. <i>oblonga</i> with a long, narrow labellum constricted in the middle; tea-tree swamps on black alkaline mud in sth-east SA, also Vic?; <i>leptochila</i> ms (Bates 2011)
<i>Microtis</i> species 'Topperwein'		aff. <i>oblonga</i> with tall spike of larger, well-spaced fleshy flowers; River Red Gum swamps in sth-east SA & west Vic, flowering without fire (Bates 2011)
<i>Orthoceras strictum</i> R.Br.		
<i>Orthoceras</i> species 'short bracts'		aff. <i>strictum</i> with short flower bracts, rounded labellum apex; heathy woodland Otway Ranges; resembles New Zealand species <i>O. novae-zeelandiae</i> (Jeanes & Backhouse 2006); grows with and probably only a variant of typical <i>O. strictum</i>
<i>Paracaleana disjuncta</i> D.L.Jones	<i>Caleana</i>	
<i>Paracaleana minor</i> R.Br.	<i>Caleana</i>	
<i>Phladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	
<i>Plectorrhiza tridentata</i> (Lindl.) Dockrill		
<i>Prasophyllum alpestre</i> D.L.Jones		
<i>Prasophyllum anticum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum appendiculatum</i> Nicholls		
<i>Prasophyllum australe</i> R.Br.		
<i>Prasophyllum barnettii</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum beatrix</i> D.L.Jones & D.T.Rouse		

Victorian Species	Alternate Genus	Notes
<i>Prasophyllum brevilabre</i> (Lindl.) Hook.f.		
<i>Prasophyllum chasmogamum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum colemaniae</i> R.S.Rogers		probably extinct
<i>Prasophyllum correctum</i> D.L.Jones		
<i>Prasophyllum diversiflorum</i> Nicholls		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum erythrocommum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum flavum</i> R.Br.		
<i>Prasophyllum fosteri</i> D.L.Jones		
<i>Prasophyllum frenchii</i> F.Muell.		
<i>Prasophyllum gilgai</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum hygrophilum</i> D.L.Jones & D.Rouse		
<i>Prasophyllum lindleyanum</i> Rchb.f.		
<i>Prasophyllum litorale</i> R.J.Bates		
<i>Prasophyllum maccannii</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum morganii</i> Nicholls		probably extinct
<i>Prasophyllum niphopedium</i> D.L.Jones		
<i>Prasophyllum occidentale</i> R.S.Rogers		
<i>Prasophyllum occultans</i> R.J.Bates		
<i>Prasophyllum odoratum</i> R.S.Rogers		
<i>Prasophyllum parviflorum</i> Nicholls		
<i>Prasophyllum patens</i> R.Br.		
<i>Prasophyllum pyriforme</i> E.Coleman		
<i>Prasophyllum readii</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum retroflexum</i> D.L.Jones		
<i>Prasophyllum sphacelatum</i> D.L.Jones		
<i>Prasophyllum spicatum</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum suaveolens</i> D.L.Jones & R.J.Bates		
<i>Prasophyllum subbisectum</i> Nicholls		
<i>Prasophyllum suttonii</i> R.S.Rogers & B.Rees		probably extinct
<i>Prasophyllum sylvestre</i> R.J.Bates & D.L.Jones		
<i>Prasophyllum tadgellianum</i> R.S.Rogers		
<i>Prasophyllum uvidulum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum viretrum</i> D.L.Jones & D.T.Rouse		
<i>Prasophyllum</i> species 'Anglesea'		late spring flowering aff. <i>odoratum</i> with fewer, smaller flowers with a shorter labellum; heathy woodland at Anglesea (Jeanes & Backhouse 2006; = aff. <i>odoratum</i> E Rouse 2002)
<i>Prasophyllum</i> species 'basalt plains'		spring-flowering, robust, clump-forming aff. <i>suaveolens</i> with larger flowers; western basalt plains grasslands (Jeanes & Backhouse 2006; = aff. <i>occidentale</i> E Rouse 2002)
<i>Prasophyllum</i> species 'Boorhaman'		spring-flowering aff. <i>petilum</i> with very narrow labellum and narrow petals; native grasslands near Boorhaman (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'brown lip'		aff. <i>fitzgeraldii</i> with brown lip with narrow white margin; heathy woodland in lower south-east of SA and far western Vic (Bates 2011; Jeanes & Backhouse 2006); <i>kafechilum</i> ms
<i>Prasophyllum</i> species 'Bunyip'		tall, fairly slender aff. <i>odoratum</i> with widely spaced flowers with narrowly spreading to incurved petals; open forest in west & central Gippsland, seen mostly after fire (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Dandenong Ranges'		aff. <i>odoratum</i> with tall spike of well-spaced recurved flowers; open forest in the Dandenong Ranges (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> G Rouse 2002)
<i>Prasophyllum</i> species 'Deep Lead'		aff. <i>fitzgeraldii</i> with short-lived flowers with brown labellum lamina; heathy woodland on clay soils at Deep Lead (Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> B Rouse 2002)
<i>Prasophyllum</i> species 'Digby'		aff. <i>fitzgeraldii</i> with green labellum lamina and pink callus; dense heathy woodland near Digby, flowering after fire (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'eastern highlands'		summer flowering aff. <i>odoratum</i> with tall, moderately crowded spike of large flowers; subalpine woodland of eastern Vic and southern NSW above 800 m (=aff. <i>odoratum</i> L Rouse 2002)
<i>Prasophyllum</i> species 'foothills'		small, early spring-flowering aff. <i>patens</i> with slender spike of widely spaced flowers with long, narrow pointed labellum; foothill open forest from Moondarra (Vic) to Yass (NSW) (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'grassy woodlands'		spring-flowering aff. <i>diversiflorum</i> with shorter spike of smaller, more crowded flowers; damp sites in grassy woodland in the north-east between Nagambie and Chesney (Jeanes & Backhouse 2006)

Victorian Species	Alternate Genus	Notes
<i>Prasophyllum</i> species 'Inglewood'		small aff. <i>pyriforme</i> with a raised, broadly and deeply grooved labellum callus; dry woodland at Inglewood & Arnold West (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'inland woodlands'		aff. <i>validum</i> with large, relatively broad dull brownish-green flowers; widespread across drier inland woodlands Chiltern to Stawell (Jeanes & Backhouse 2006; =aff. <i>pyriforme</i> 5 Rouse 2002)
<i>Prasophyllum</i> species 'Inverleigh'		aff. <i>validum</i> with dull green flowers with short, broad column arms; sparsely shrubby woodland on fertile sandy loam soils at Inverleigh (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Lara'		aff. <i>odoratum</i> with long narrow green callus plate; basalt plains grasslands near Lara; presumed extinct (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> M Rouse)
<i>Prasophyllum</i> species 'limestone heath'		small aff. <i>parviflorum</i> with smaller, paler flowers; coastal heath on limestone between Portland and Nelson (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Mallacoota'		robust aff. <i>spicatum</i> with smaller, less crowded, fragrant flowers with widely spreading petals; coastal heaths, heathy woodland in East Gippsland, mostly seen after fire (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> K Rouse 2002)
<i>Prasophyllum</i> species 'mallee sands'		aff. <i>odoratum</i> with narrow overlapping green and white flowers; mallee heath on sand dunes in the Big Desert (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> B Rouse)
<i>Prasophyllum</i> species 'Midlands'		tall aff. <i>odoratum</i> with widely spaced, somewhat narrow flowers; woodland on rocky hills from Chewton to St Arnaud (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'montane'		aff. <i>montanum</i> with narrower labellum callus plate, broader lamina and longer, narrower sepals; foothills and mountains of eastern Victoria from Alexandra to Wulgulmerang (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Mortlake'		tall, robust aff. <i>correctum</i> with larger, more numerous and more crowded flowers; basalt plains grassland at Mortlake (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'mountain bogs'		relatively tall, late summer-flowering aff. <i>alpestre</i> with narrow petals and sharply curved labellum; mountain swamps above 700 m, Tamboritha (Vic) to Adaminaby (NSW) (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Murchison'		aff. <i>petilum</i> with narrower, more brownish to greenish labellum; native grasslands near Murchison (Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Noorinbee'		slender early spring-flowering aff. <i>odoratum</i> with few widely spaced flowers; shrubby open forest in East Gippsland, seen mostly after fires (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> I Rouse 2002)
<i>Prasophyllum</i> species 'Nunniong'		larger, mid-late summer flowering aff. <i>sphacelatum</i> with broader lip and growing in damper sites; alpine plains and woodland in the Nunniong and Mt Cobberas regions (Backhouse <i>et al.</i> 2016)
<i>Prasophyllum</i> species 'Otway coast'		aff. <i>odoratum</i> with short, crowded spike; coastal heath from Anglesea to Port Campbell (=aff. <i>odoratum</i> E/F Rouse 2002)
<i>Prasophyllum</i> species 'Otway Ranges'		tall, slender aff. <i>odoratum</i> with small, widely spaced flowers; shrubby open forest in the Otway Ranges, seen mostly after fire (Jeanes & backhouse 2006; =aff. <i>odoratum</i> H Rouse 2002)
<i>Prasophyllum</i> species 'pink lip'		aff. <i>fitzgeraldii</i> with pink labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> A Rouse); <i>rosaechilum</i> ms
<i>Prasophyllum</i> species 'purple lip'		aff. <i>fitzgeraldii</i> with purple labellum lamina; heath and heathy woodland in sth-eastern SA & western Vic (Bates 2011; Jeanes & Backhouse 2006; =aff. <i>fitzgeraldii</i> E Rouse); <i>violescens</i> ms
<i>Prasophyllum</i> species 'Riverina grasslands'		tall, early flowering aff. <i>suaveolens</i> with larger flowers with a long narrowly triangular labellum; Riverina plains grasslands from Hunter (Vic) to Moama (NSW) (Jeanes & Backhouse 2006; =aff. <i>occidentale</i> D Rouse 2002)
<i>Prasophyllum</i> species 'sedge swamps'		short aff. <i>odoratum</i> with few widely spaced flowers; heath and sedge swamps, often flowering in water (= sp. 'Drajurk' Backhouse <i>et al.</i> 2016; sp. 'Topperwein' Bates 2011)
<i>Prasophyllum</i> species 'Smokers Gap'		large, late-summer flowering aff. <i>sphacelatum</i> ; alpine meadows in Namadgi National Park; mentioned in notes on <i>P. spacelatum</i> in Jones <i>et al.</i> (2008); possibly the same as <i>P. species</i> 'Nunniong'?
<i>Prasophyllum</i> species 'waterholes'		very late flowering aff. <i>odoratum</i> with very widely spreading lateral sepals and incurved petals; heath and sedge swamps in south-western Vic & south-eastern SA (Jeanes & Backhouse 2006; =aff. <i>odoratum</i> C/D Rouse 2002)
<i>Prasophyllum</i> species 'Wilson's Promontory'		aff. <i>frenchii</i> with smaller, duller flowers with more ruffled labellum and lower callus; moist coastal heath at Wilson's Promontory (= sp. aff. <i>frenchii</i> 2 Jeanes & Backhouse 2006)
<i>Prasophyllum</i> species 'Wimmera woodlands'		short aff. <i>occidentale</i> with smaller, more widely spaced flowers with a narrower labellum; dry woodlands in narrow band between Murtoa (Vic) and Keith (SA)
<i>Pterostylis aciculiformis</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis acuminata</i> R.Br.		
<i>Pterostylis aestiva</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis agrestis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis alpina</i> R.S.Rogers		
<i>Pterostylis alveata</i> Garnet	<i>Diplodium</i>	

Victorian Species	Alternate Genus	Notes
<i>Pterostylis aneba</i> D.L.Jones		Victorian records in doubt
<i>Pterostylis atrans</i> D.L.Jones	<i>Diplodium</i>	
<i>Pterostylis baptistii</i> Fitzg.		
<i>Pterostylis basaltica</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis bicolor</i> M.A.Clem. & D.L.Jones	<i>Hymenochilus</i>	
<i>Pterostylis boormanii</i> Rupp	<i>Oligochaetochilus</i>	
<i>Pterostylis cheraphila</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis chlorogramma</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis clivicola</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis clivosa</i> (D.L.Jones) D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis coccinea</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis concinna</i> R.Br.	<i>Taurantha</i>	
<i>Pterostylis conferta</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis crassa</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis crassicaulis</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis cucullata</i> R.Br. subspecies <i>cucullata</i>		
<i>Pterostylis cucullata</i> R.Br. subspecies <i>sylvicola</i> D.L.Jones		
<i>Pterostylis curta</i> R.Br.		
<i>Pterostylis cycnocephala</i> Fitzg.	<i>Hymenochilus</i>	
<i>Pterostylis decurva</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis despectans</i> (Nicholls) D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis diminuta</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis dolichochila</i> D.L.Jones & M.A.Clem.	<i>Diplodium</i>	
<i>Pterostylis falcata</i> R.S.Rogers		
<i>Pterostylis ferruginea</i> (D.L.Jones) G.N.Backh.	<i>Oligochaetochilus</i>	
<i>Pterostylis fischii</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis foliata</i> Hook.f.		
<i>Pterostylis grandiflora</i> R.Br.	<i>Diplodium</i>	
<i>Pterostylis hamata</i> Blackmore & Clemesha	<i>Oligochaetochilus</i>	
<i>Pterostylis incognita</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	probably extinct
<i>Pterostylis jonesii</i> G.N.Backh.	<i>Bunochilus</i>	originally described as <i>Bunochilus montanus</i> D.L. Jones
<i>Pterostylis laxa</i> Blackmore	<i>Diplodium</i>	
<i>Pterostylis lingua</i> M.A.Clem.	<i>Oligochaetochilus</i>	there is some doubt as to the correct identity of Victorian records of this species
<i>Pterostylis loganii</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis longipetala</i> Rupp	<i>Diplodium</i>	
<i>Pterostylis lustra</i> D.L.Jones		
<i>Pterostylis macilenta</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis maxima</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis melagramma</i> D.L.Jones	<i>Bunochilus</i>	
<i>Pterostylis monticola</i> D.L.Jones		
<i>Pterostylis multiflora</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis nana</i> R.Br.	<i>Linguella</i>	
<i>Pterostylis nutans</i> R.Br.		
<i>Pterostylis oreophila</i> Clemesha		
<i>Pterostylis parviflora</i> R.Br.	<i>Speculantha</i>	
<i>Pterostylis pedoglossa</i> Fitzg.	<i>Crangonorchis</i>	
<i>Pterostylis pedunculata</i> R.Br.		
<i>Pterostylis planulata</i> D.L.Jones & M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis plumosa</i> Cady	<i>Plumatichilos</i>	
<i>Pterostylis prasina</i> (D.L.Jones) G.N.Backh.	<i>Bunochilus</i>	
<i>Pterostylis pusilla</i> R.S.Rogers	<i>Oligochaetochilus</i>	
<i>Pterostylis robusta</i> R.S.Rogers	<i>Diplodium</i>	
<i>Pterostylis rubescens</i> (D.L.Jones) G.N.Backh.	<i>Speculantha</i>	
<i>Pterostylis sanguinea</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	
<i>Pterostylis setifera</i> M.A.Clem., Matthias & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis smaragdina</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	

Victorian Species	Alternate Genus	Notes
<i>Pterostylis spissa</i> (D.L.Jones) G.N.Backh.	<i>Hymenochilus</i>	
<i>Pterostylis squamata</i> R.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis striata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis tasmanica</i> D.L.Jones	<i>Plumatichilos</i>	
<i>Pterostylis tenuissima</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis truncata</i> Fitzg.	<i>Diplodium</i>	
<i>Pterostylis tunstallii</i> D.L.Jones & M.A.Clem.	<i>Bunochilus</i>	
<i>Pterostylis uliginosa</i> D.L.Jones	<i>Speculantha</i>	
<i>Pterostylis valida</i> (Nicholls) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis woollsii</i> Fitzg.	<i>Oligochaetochilus</i>	
<i>Pterostylis xerophila</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis</i> species 'Anglesea'	<i>Plumatichilos</i>	large aff. <i>plumosa</i> with long sepal points; coastal heathy woodland at Anglesea (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Beechworth'	<i>Oligochaetochilus</i>	aff. <i>boormanii</i> with less bristly, brown flowers; <i>Callitris</i> spp. woodland on granite hills around Beechworth (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Berrook'	<i>Oligochaetochilus</i>	small aff. <i>excelsa</i> ; mallee on sandy soils in the south-western Sunset Country near Berrook (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'bristly lip'		small non-colony forming aff. <i>nutans</i> with less nodding flower with a very bristly labellum; forest in south-western Victoria flowering after disturbance such as grading or fire (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'chunky flowers'	<i>Diplodium</i>	aff. <i>dolichochila</i> with a shorter labellum; coastal scrubs from Kingston (SA) to Portland (Vic) (Bates 2011; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Daylesford'	<i>Diplodium</i>	aff. <i>deflexa</i> with smaller flowers with only gently curved galea, tip just below horizontal, not pointing to ground; foothill forest between Woodend and Ballarat (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'Dundonnell'	<i>Hymenochilus</i>	aff. <i>bicolor</i> with very short spike of densely crowded flowers, basalt plains grasslands near Dundonnell (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'East Gippsland'	<i>Speculantha</i>	aff. <i>rubescens</i> with slightly larger, plumper flowers; heath and heathy open forest in East Gippsland (AVH record) <i>protera</i> ms
<i>Pterostylis</i> species 'foothill forest'	<i>Diplodium</i>	autumn flowering aff. <i>alveata</i> with more protruding sinus; foothill forests in East Gippsland east from Bruthen (Jeanes & Backhouse 2006; Turner <i>et al.</i> 2014)
<i>Pterostylis</i> species 'Gippsland'	<i>Plumatichilos</i>	aff. <i>plumosa</i> with smaller flowers with a green apical knob; heathy woodland and open forest in East Gippsland between Bruthen and Genoa (Turner <i>et al.</i> 2014)
<i>Pterostylis</i> species 'greater Grampians'	<i>Speculantha</i>	aff. <i>rubescens</i> ; woodland on gravelly clay soils in the Ararat-Stawell region and northern Grampians; no further information available (AVH record) <i>aurifera</i> ms
<i>Pterostylis</i> species 'Guildford'	<i>Bunochilus</i>	aff. <i>macrosepala</i> with a distinctly rectangular labellum with a very short midlobe; open forest at Guildford (Jones & Clements 2006)
<i>Pterostylis</i> species 'Lara'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with short, thick stem and smallish flowers; basalt plains grasslands near Lara, probably extinct (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'large curved flowers'	<i>Diplodium</i>	aff. <i>revoluta</i> with larger, more curved flowers; widespread in dry woodland and open forest through inland south-eastern Australia (Bishop 2000; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'large striped flowers'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with large brownish striped flowers, mallee, <i>Callitris</i> woodland and dry woodland from Charlton (Vic) to Renmark (SA) (Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'Maldon'	<i>Oligochaetochilus</i>	aff. <i>biseta</i> with a short stem bearing very large, striped flowers with a bristly lip; sparsely shrubby woodland on granite hills near Maldon (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'mallee heath'	<i>Plumatichilos</i>	smallish glossy aff. <i>plumosa</i> with long pedicel and long dense labellum hairs; mallee heath on sandy to clay loam soils in western Vic & sth-eastern SA (Bates 2011; Jeanes & Backhouse 2006)
<i>Pterostylis</i> species 'mallee snail'	<i>Linguella</i>	short aff. <i>nana</i> with small squat flowers with short lateral sepals; mallee on sandy to clay loam soils (Backhouse <i>et al.</i> 2016; Bates 2011) <i>tenuis</i> ms
<i>Pterostylis</i> species 'Nora Creina'	<i>Diplodium</i>	aff. <i>alata</i> with tricolored flowers, a long fine point on dorsal sepal and a distinct gap between lateral sepals and galea; coastal scrubs in the lower south-east SA and sth-west Vic (Bates 2011)
<i>Pterostylis</i> species 'north-east Vic hills'	<i>Oligochaetochilus</i>	aff. <i>aciculiformis</i> with long lateral sepals and broader more bristly labellum; dry woodland on rocky hills in the north-east (Jeanes & Backhouse 2006); falls within the variation of and doubtfully distinct from <i>Pterostylis aciculiformis</i>
<i>Pterostylis</i> species 'north-west mallee'	<i>Oligochaetochilus</i>	aff. <i>excelsa</i> with large green flowers; mallee on sandy to clay loam soils in the Big Desert and southern Sunset Country (Backhouse <i>et al.</i> 2016)
<i>Pterostylis</i> species 'Omeo'	<i>Bunochilus</i>	aff. <i>smaragdina</i> with a narrowly obovate labellum; montane open forest near Omeo (Jones & Clements 2006)
<i>Pterostylis</i> species 'red lip'	<i>Bunochilus</i>	aff. <i>smaragdina</i> with large, rectangular red labellum; foothills open forest between Tonimbuk and Mallacoota (Backhouse <i>et al.</i> 2016; Turner <i>et al.</i> 2015)
<i>Pterostylis</i> species 'western Victoria'	<i>Oligochaetochilus</i>	small aff. <i>aciculiformis</i> with strongly backswept lateral sepals; dry woodland and mallee, sometimes in rocky areas, in the west (Bishop 2000)



Victorian Species	Alternate Genus	Notes
<i>Pterostylis</i> species 'western woodlands'	<i>Oligochaetochilus</i>	early summer flowering aff. <i>biseta</i> with smaller, less prominently striped flowers; dry woodlands in western Vic (Melb to Stawell to Portland)
<i>Pterostylis</i> species 'Wilson's Promontory'		small aff. <i>alpina</i> ; coastal scrubs on sand dunes at Tidal River Wilson's Promontory (Jeanes & Backhouse 2006); probably only a form of <i>P. alpina</i>
<i>Pterostylis</i> species 'woodlands'	<i>Hymenochilus</i>	tall aff. <i>cynocephala</i> with striped, spaced flowers; widespread in woodlands across SA and in west Vic; <i>nemoralis</i> ms (Bates 2011)
<i>Pterostylis</i> species 'Woolhpooper'	<i>Bunochilus</i>	aff. <i>melagramma</i> with a thin labellum which has a large basal lobe; grassy woodlands near Woolhpooper (Jones & Clements 2006)
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		
<i>Sarcochilus australis</i> (Lindl.) Rchb.f.		
<i>Sarcochilus falcatus</i> R.Br.		
<i>Spiranthes alticola</i> D.L.Jones		very difficult to distinguish from (and doubtfully distinct from) <i>Spiranthes australis</i>
<i>Spiranthes australis</i> (R.Br.) Lindl.		
<i>Thelymitra alcockii</i> D.L.Jones ex Jeanes		
<i>Thelymitra alpicola</i> Jeanes		
<i>Thelymitra alpina</i> Jeanes		
<i>Thelymitra antennifera</i> (Lindl.) Hook.f.		
<i>Thelymitra arenaria</i> Lindl.		
<i>Thelymitra aristata</i> Lindl.		
<i>Thelymitra atronitida</i> Jeanes		
<i>Thelymitra azurea</i> R.S.Rogers		
<i>Thelymitra basaltica</i> Jeanes		
<i>Thelymitra benthamiana</i> Rchb.f.		
<i>Thelymitra bracteata</i> J.Z.Weber ex Jeanes		
<i>Thelymitra brevifolia</i> Jeanes		
<i>Thelymitra carnea</i> R.Br.		
<i>Thelymitra circumsepta</i> Fitzg.		
<i>Thelymitra cyanea</i> (Lindl.)		
<i>Thelymitra epipactoides</i> F.Muell.		
<i>Thelymitra exigua</i> Jeanes		
<i>Thelymitra flexuosa</i> Endl.		
<i>Thelymitra gregaria</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra hiemalis</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra holmesii</i> Nicholls		
<i>Thelymitra improcera</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra incurva</i> Jeanes		
<i>Thelymitra ixioides</i> Sw.		
<i>Thelymitra juncifolia</i> Lindl.		
<i>Thelymitra latifolia</i> R.J.Bates		
<i>Thelymitra longiloba</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra lucida</i> Jeanes		
<i>Thelymitra luteocilium</i> Fitzg.		
<i>Thelymitra mackibbinii</i> F.Muell.		
<i>Thelymitra malvina</i> M.A.Clem.		
<i>Thelymitra matthewsii</i> Cheesem.		
<i>Thelymitra media</i> R.Br.		
<i>Thelymitra megalyptra</i> Fitzg.		
<i>Thelymitra mucida</i> Fitzg.		
<i>Thelymitra nuda</i> R.Br.		
<i>Thelymitra orientalis</i> R.J.Bates		
<i>Thelymitra pallidiflora</i> Jeanes		
<i>Thelymitra pallidifructus</i> R.J.Bates		
<i>Thelymitra pauciflora</i> R.Br.		
<i>Thelymitra peniculata</i> Jeanes		
<i>Thelymitra planicola</i> Jeanes		
<i>Thelymitra reflexa</i> Jeanes		
<i>Thelymitra rubra</i> Fitzg.		
<i>Thelymitra simulata</i> D.L.Jones & M.A.Clem.		

Victorian Species	Alternate Genus	Notes
<i>Thelymitra</i> species 'insipid'		very slender, late flowered aff. <i>pauciflora</i> with long narrow sheathing leaf, rarely opening dull blue flowers and flat red and yellow post-anther lobe; widely distributed in many habitats in cooler, higher rainfall areas (Bates 2011)
<i>Thelymitra</i> species 'red gum swamps'		aff. <i>inflata</i> with more circular and narrower column apex; seasonal red gum swamps (SA, Vic) or wet forest (Tas) (Bates 2011)
<i>Thelymitra</i> species 'Topperwein'		aff. <i>holmesii</i> with horseshoe-shaped, rounded column apex and yellow hair tufts; sedge swamps in River Red Gum woodland in the lower south-east of SA and western Vic (Bates 2011)
<i>Thynninorchis huntianus</i> (F.Muell.) D.L.Jones & M.A.Clem.	<i>Arthrochilus</i>	

**Total: 454 taxa (species, varieties)**

Victorian Hybrids	Name and Authority	Notes
<i>Caladenia ampla (reticulata) X cardiochila</i>		Backhouse 2011
<i>Caladenia ampla (reticulata) X clavigera</i>		G. Backhouse personal observation
<i>Caladenia ampla (reticulata) X fulva</i>		Backhouse 2011
<i>Caladenia ampla (reticulata) X grampiana</i>		Backhouse 2011
<i>Caladenia ancylosa X peisleyi</i>		Backhouse 2011
<i>Caladenia australis X venusta</i>		Jeanes & Backhouse 2006
<i>Caladenia capillata X fuscata</i>		Jeanes & Backhouse 2006
<i>Caladenia cardiochila X clavigera</i>		Jeanes & Backhouse 2006
<i>Caladenia cardiochila X formosa</i>		Jeanes & Backhouse 2006
<i>Caladenia cardiochila X lowanensis</i>		Jeanes & Backhouse 2006
<i>Caladenia cardiochila X stricta</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia cardiochila X venusta</i>		Jeanes & Backhouse 2006
<i>Caladenia cardiochila X species 'mallee heath'</i>		Bates 2011
<i>Caladenia carnea X catenata</i>		Jeanes & Backhouse 2006
<i>Caladenia carnea X cucullata</i>		Kuiter 2010
<i>Caladenia carnea X fuscata</i>		Jones <i>et al.</i> 1999
<i>Caladenia carnea X maritima</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia carnea X vulgaris</i>		Bates 2011
<i>Caladenia catenata X clarkiae</i>		Backhouse 2011
<i>Caladenia clarkiae X hildae</i>		Jones 2006
<i>Caladenia clavigera X montana</i>		Jeanes & Backhouse 2006
<i>Caladenia clavigera X oenochila</i>		Jeanes & Backhouse 2006
<i>Caladenia clavigera X parva</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia colorata X formosa</i>		Backhouse 2011
<i>Caladenia colorata X species 'mallee heath'</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia colorata X species 'Murray mallee'</i>		Bates 2011
<i>Caladenia congesta X hildae</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia cretacea X cruciformis</i>		Backhouse 2011
<i>Caladenia cretacea X douglasiorum</i>		Backhouse 2011
<i>Caladenia cretacea X tentaculata</i>		Backhouse 2011
<i>Caladenia filamentosa X montana</i>		Jeanes & Backhouse 2006
<i>Caladenia flavovirens X parva</i>		Backhouse 2011
<i>Caladenia formosa X parva</i>		G. Backhouse personal observation
<i>Caladenia formosa X tentaculata</i>		Backhouse 2011
<i>Caladenia formosa X venusta</i>		Jeanes & Backhouse 2006
<i>Caladenia gracilis X hildae</i>		Jeanes & Backhouse 2006
<i>Caladenia grampiana X parva</i>		Backhouse 2011
<i>Caladenia hildae X moschata</i>		Jeanes & Backhouse 2006
<i>Caladenia lowanensis X stricta</i>		Jeanes & Backhouse 2006
<i>Caladenia lowanensis X tensa</i>		Jeanes & Backhouse 2006
<i>Caladenia oenochila X rosella</i>		Backhouse 2011
<i>Caladenia oenochila X tentaculata</i>		Jeanes & Backhouse 2006
<i>Caladenia orientalis X tessellata</i>	<i>xvariabilis</i> Nicholls	
<i>Caladenia osmera X peisleyi</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia parva X venusta</i>		Jeanes & Backhouse 22006
<i>Caladenia peisleyi X tessellata</i>		Backhouse <i>et al.</i> 2016
<i>Caladenia stricta X verrucosa</i>		Jeanes & Backhouse 2006
<i>Caladenia tentaculata X species 'Raymond Island'</i>		Backhouse 2011
<i>Caladenia tentaculata X venusta</i>		Jeanes & backhouse 2006
<i>Caladenia venusta X villosissima</i>		Backhouse 2011
<i>Caladenia fuscata X Pheladenia deformis</i>		Backhouse <i>et al.</i> 2016
<i>Calochilus paludosus X robertsonii</i>		Backhouse <i>et al.</i> 2016
<i>Chiloglottis cornuta X valida</i>		Jeanes & Backhouse 2006
<i>Chiloglottis trapeziformis X valida</i>	<i>xpescottiana</i> R.S. Rogers	
<i>Corunastylis archeri X despectans</i>		Jeanes & Backhouse 2006
<i>Corunastylis arrecta X despectans</i>		Jeanes & Backhouse 2006
<i>Corunastylis despectans X morrisii</i>		Bates 2011
<i>Cyanicula caerulea X Glossodia major</i>		
<i>Cyanicula caerulea X Glossodia minor</i>		Backhouse 2011

Victorian Hybrids	Name and Authority	Notes
<i>Cyanicula caerulea</i> X <i>Pheladenia deformis</i>		Brown <i>et al.</i> 2013
<i>Dipodium pardalinum</i> X <i>roseum</i>		Bates 2011; Backhouse <i>et al.</i> 2016
<i>Diuris basaltica</i> X <i>palustris</i>	× <i>fastidiosa</i> R.S. Rogers	
<i>Diuris behrii</i> X <i>orientis</i>		Bates 2011
<i>Diuris behrii</i> X <i>pardina</i>	× <i>palachila</i> R.S. Rogers	Bates 2011; Jeanes & Backhouse 2006
<i>Diuris behrii</i> X <i>punctata</i>		Backhouse, <i>et al.</i> 2016; Jones 2006
<i>Diuris behrii</i> X species 'thick sepals'		
<i>Diuris chrysiopsis</i> X <i>orientis</i>		Jeanes & Backhouse 2006
<i>Diuris chrysiopsis</i> X <i>pardina</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Diuris chrysiopsis</i> X <i>palustris</i>		Jeanes & Backhouse 2006
<i>Diuris chrysiopsis</i> X <i>punctata</i>		Jeanes & Backhouse 2006
<i>Diuris gregaria</i> X <i>palustris</i>		Jeanes & Backhouse 2006
<i>Diuris gregaria</i> X <i>pardina</i>		
<i>Diuris monticola</i> X <i>pardina</i>		Backhouse <i>et al.</i> 2016
<i>Diuris orientis</i> X <i>pardina</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Diuris orientis</i> X <i>sulphurea</i>		Jeanes & Backhouse 2006
<i>Diuris pardina</i> X <i>subalpina</i>		
<i>Diuris pardina</i> X <i>sulphurea</i>		Backhouse <i>et al.</i> 2016
<i>Diuris punctata</i> X species 'Gippsland plains'		Backhouse <i>et al.</i> 2016
<i>Glossodia major</i> X <i>minor</i>		Jeanes & Backhouse 2006
<i>Glossodia major</i> X <i>Pheladenia deformis</i>	× <i>Glossadenia tutelata</i> Kavulak	
<i>Prasophyllum alpestre</i> X <i>sphacelatum</i>		Backhouse <i>et al.</i> 2016
<i>Prasophyllum alpestre</i> X <i>tadgellianum</i>		Jeanes & Backhouse 2006
<i>Prasophyllum chasmogamum</i> X <i>frenchii</i>		Backhouse <i>et al.</i> 2016
<i>Prasophyllum niphopedium</i> X <i>sphacelatum</i>		Jeanes & Backhouse 2006
<i>Prasophyllum viretrum</i> X species 'Mortlake'		Backhouse <i>et al.</i> 2016
<i>Pterostylis aciculiformis</i> X species 'Beechworth'		Backhouse <i>et al.</i> 2016
<i>Pterostylis aestiva</i> X <i>atrans</i>		Jeanes & Backhouse 2006
<i>Pterostylis aestiva</i> X <i>decurva</i>		Jeanes & Backhouse 2006
<i>Pterostylis alpina</i> X <i>cucullata</i>		Jeanes & Backhouse 2006
<i>Pterostylis alpina</i> X <i>foliata</i>		Jeanes & Backhouse 2006
<i>Pterostylis atrans</i> X <i>decurva</i>		Jeanes & Backhouse 2006; Jones <i>et al.</i> 2008
<i>Pterostylis baptistii</i> X <i>falcata</i>		Jeanes & Backhouse 2006
<i>Pterostylis boormanii</i> X <i>setifera</i>		Jeanes & Backhouse 2006
<i>Pterostylis cheraphila</i> X <i>ferruginea</i>		Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>fischii</i>		Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>nutans</i>		Backhouse <i>et al.</i> 2016
<i>Pterostylis concinna</i> X <i>striata</i>	× <i>toveyana</i> Ewart & Sharman	
<i>Pterostylis cucullata</i> X <i>falcata</i>	× <i>aenigma</i> D.L. Jones & M.A. Clem.	
<i>Pterostylis cucullata</i> X <i>foliata</i>		Jeanes & Backhouse 2006
<i>Pterostylis cucullata</i> X <i>nutans</i>		Bates 2011
<i>Pterostylis curta</i> X <i>nutans</i>		Jeanes & Backhouse 2006
<i>Pterostylis curta</i> X <i>pedunculata</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Pterostylis falcata</i> X <i>nutans</i>	× <i>ingens</i> (Rupp) D.L. Jones	
<i>Pterostylis ferruginea</i> X <i>planulata</i>		Backhouse <i>et al.</i> 2016
<i>Pterostylis fischii</i> X <i>laxa</i>		Jeanes & Backhouse 2006
<i>Pterostylis lingua</i> X species 'large striped flowers'		Backhouse <i>et al.</i> 2016
<i>Pterostylis parviflora</i> X <i>rubescens</i>		
<i>Pterostylis robusta</i> X <i>striata</i>		Backhouse <i>et al.</i> 2016
<i>Pterostylis truncata</i> X species 'large curved flowers'		Backhouse <i>et al.</i> 2016
<i>Thelymitra alcockiae</i> X <i>antennifera</i>		Backhouse <i>et al.</i> 2016
<i>Thelymitra alpina</i> X <i>simulata</i>		Backhouse <i>et al.</i> 2016
<i>Thelymitra antennifera</i> X <i>gregaria</i>		Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>luteocilium</i>		Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>megcalyptera</i>		Jeanes & Backhouse 2006
<i>Thelymitra antennifera</i> X <i>nuda</i>	× <i>macmillanii</i> F. Muell.	Bates 2011
<i>Thelymitra arenaria</i> X <i>ixioides</i>		
<i>Thelymitra arenaria</i> X <i>nuda</i>		Jeanes 2004

Victorian Hybrids	Name and Authority	Notes
<i>Thelymitra aristata</i> X <i>ixioides</i>		Jeanes & Backhouse 2006
<i>Thelymitra aristata</i> X <i>media</i>		Jeanes & Backhouse 2006
<i>Thelymitra brevifolia</i> X <i>ixioides</i>		
<i>Thelymitra carnea</i> X <i>ixioides</i>	× <i>irregularis</i> Nicholls	
<i>Thelymitra carnea</i> X <i>peniculata</i>		Jeanes & Backhouse 2006
<i>Thelymitra exigua</i> X <i>gregaria</i>		Jeanes & Backhouse 2006
<i>Thelymitra exigua</i> X <i>ixioides</i>		Backhouse <i>et al.</i> 2016
<i>Thelymitra ixioides</i> X <i>pauciflora</i>	× <i>truncata</i> R.S. Rogers	
<i>Thelymitra ixioides</i> X <i>peniculata</i>	× <i>merraniae</i> Nicholls	
<i>Thelymitra juncifolia</i> X <i>megcalyptra</i>		Bates 2011
<i>Thelymitra juncifolia</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra luteocilium</i> X <i>megcalyptra</i>		Bates 2011; Jeanes & Backhouse 2006
<i>Thelymitra luteocilium</i> X <i>nuda</i>	× <i>chasmogama</i> R.S. Rogers	Bates 2011
<i>Thelymitra luteocilium</i> X <i>rubra</i>		Bates 2011
<i>Thelymitra luteocilium</i> X <i>pauciflora</i>		Backhouse <i>et al.</i> 2016

**Total: 132 hybrids**

Western Australian Species	Alternate Genus	Notes
<i>Arthrochilus byrnesii</i> Blaxell	<i>Phoringopsis</i>	
<i>Caladenia abbreviata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia ambusta</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia applanata</i> Hopper & A.P.Br. subspecies <i>applanata</i>	<i>Arachnorchis</i>	
<i>Caladenia applanata</i> Hopper & A.P.Br. subspecies <i>erubescens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia arenicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia arrecta</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>attingens</i>	<i>Arachnorchis</i>	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>effusa</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia attingens</i> Hopper & A.P.Br. subspecies <i>gracillima</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia barbarella</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	
<i>Caladenia barbarossa</i> Rchb.f.	<i>Drakonorchis</i>	
<i>Caladenia bicallata</i> R.S.Rogers subspecies <i>bicallata</i>	<i>Jonesiopsis</i>	
<i>Caladenia bicallata</i> R.S.Rogers subspecies <i>cleistogama</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia bigeminata</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia brevisura</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia brownii</i> Hopper	<i>Arachnorchis</i>	
<i>Caladenia bryceana</i> R.S.Rogers subspecies <i>bryceana</i>	<i>Arachnorchis</i>	
<i>Caladenia bryceana</i> R.S.Rogers subspecies <i>cracens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia busselliana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>caesarea</i>	<i>Jonesiopsis</i>	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>maritima</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia caesarea</i> (Domin) M.A.Clem. & Hopper subspecies <i>transiens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia cairnsiana</i> F.Muell.	<i>Jonesiopsis</i>	
<i>Caladenia chapmanii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia christineae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia citrina</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia corynephora</i> A.S.George	<i>Arachnorchis</i>	
<i>Caladenia crebra</i> A.S.George	<i>Arachnorchis</i>	
<i>Caladenia cristata</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia cruscula</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia decora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia denticulata</i> Lindl. subspecies <i>albicans</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia denticulata</i> Lindl. subspecies <i>denticulata</i>	<i>Jonesiopsis</i>	
<i>Caladenia denticulata</i> Lindl. subspecies <i>rubella</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia dimidia</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia discoidea</i> Lindl.	<i>Arachnorchis</i>	
<i>Caladenia dorrienii</i> Domin	<i>Jonesiopsis</i>	
<i>Caladenia douthchiai</i> O.Sarg.	<i>Jonesiopsis</i>	
<i>Caladenia drakeoides</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	
<i>Caladenia drummondii</i> Benth.	<i>Arachnorchis</i>	
<i>Caladenia dundasiae</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia elegans</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia ensata</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia erythrochila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia erythronema</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia evanescens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia excelsa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia exilis</i> Hopper & A.P.Br. subspecies <i>exilis</i>	<i>Jonesiopsis</i>	



Western Australian Species	Alternate Genus	Notes
<i>Caladenia exilis</i> Hopper & A.P.Br. subspecies <i>vanleeuwenii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia exstans</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia falcata</i> (Nicholls) M.A.Clem. & Hopper	<i>Arachnorchis</i>	
<i>Caladenia ferruginea</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia filifera</i> Lindl.	<i>Jonesiopsis</i>	
<i>Caladenia flava</i> R.Br. subspecies <i>flava</i>		
<i>Caladenia flava</i> R.Br. subspecies <i>maculata</i> Hopper & A.P.Br.		
<i>Caladenia flava</i> R.Br. subspecies <i>sylvestris</i> Hopper & A.P.Br.		
<i>Caladenia flava</i> R.Br. subspecies 'late red'		short, late-flowering <i>C. flava</i> with prominent red markings; wandoo woodland on granite outcrops between Beverley & Williams (Brown <i>et al.</i> 2013); probably only a colourful form of <i>C. flava</i> subsp. <i>flava</i>
<i>Caladenia fluvialis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia footeana</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia fuscolutescens</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia gardneri</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia georgei</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia graminifolia</i> A.S.George	<i>Arachnorchis</i>	
<i>Caladenia graniticola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia granitora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia harringtoniae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia heberleana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia hiemalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia hirta</i> Lindl. subspecies <i>hirta</i>	<i>Arachnorchis</i>	
<i>Caladenia hirta</i> Lindl. subspecies <i>rosea</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia hoffmanii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia hopperiana</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia horistes</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia huegelii</i> Rchb.f.	<i>Arachnorchis</i>	
<i>Caladenia incensa</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia incrassata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia infundibularis</i> A.S.George	<i>Arachnorchis</i>	
<i>Caladenia integra</i> E.Coleman	<i>Arachnorchis</i>	
<i>Caladenia interjacens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia latifolia</i> R.Br.		
<i>Caladenia leucochila</i> A.P.Br., R.D.Phillips & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia lobata</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia lodgeana</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>albella</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>australora</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>borealis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>calcigena</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>clivicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>crassa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>eminens</i> (Domin) Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>extrema</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>insularis</i> Hopper & A.P.Br. ex A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>longicauda</i>	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>merrittii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	

Western Australian Species	Alternate Genus	Notes
<i>Caladenia longicauda</i> Lindl. subspecies <i>minima</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>redacta</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies <i>rigidula</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia longicauda</i> Lindl. subspecies 'Bulbarnet'	<i>Arachnorchis</i>	<i>longicauda</i> subspecies with robust flowers with relatively short stiff petals and sepals with broad bases, sepals often clubbed, labellum lamina calli in numerous rows; woodland near Moora
<i>Caladenia longiclavata</i> E.Coleman	<i>Arachnorchis</i>	
<i>Caladenia longifimbriata</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia lorea</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia luteola</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia macrostylis</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia magniclavata</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia marginata</i> Lindl.		
<i>Caladenia melanema</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia meridionalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia mesocera</i> Hopper & A.P.Br.	<i>Drakonorchis</i>	
<i>Caladenia microchila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia multiclavia</i> Rchb.f.	<i>Jonesiopsis</i>	
<i>Caladenia nana</i> Endl. subspecies <i>nana</i>		
<i>Caladenia nana</i> Endl. subspecies <i>unita</i> (Fitzg.) Hopper & A.P.Br.		sometimes considered a full species <i>Caladenia unita</i>
<i>Caladenia nivalis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia nobilis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia occidentalis</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia pachychila</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia paludosa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia paradoxa</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia pectinata</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia pendens</i> Hopper & A.P.Br. subspecies <i>pendens</i>	<i>Jonesiopsis</i>	
<i>Caladenia pendens</i> Hopper & A.P.Br. subspecies <i>talbotii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia perangusta</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia petrensis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia pholcoidea</i> Hopper & A.P.Br. subspecies <i>augustensis</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia pholcoidea</i> Hopper & A.P.Br. subspecies <i>pholcoidea</i>	<i>Arachnorchis</i>	
<i>Caladenia plicata</i> Fitzg.	<i>Arachnorchis</i>	
<i>Caladenia pluvialis</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia polychroma</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia postea</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia procera</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia pulchra</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia radialis</i> R.S.Rogers	<i>Jonesiopsis</i>	
<i>Caladenia radiata</i> Nicholls	<i>Arachnorchis</i>	
<i>Caladenia remota</i> Hopper & A.P.Br. subspecies <i>parva</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia remota</i> Hopper & A.P.Br. subspecies <i>remota</i>	<i>Jonesiopsis</i>	
<i>Caladenia reptans</i> Lindl. subspecies <i>impensa</i> Hopper & A.P.Br.		
<i>Caladenia reptans</i> Lindl. subspecies <i>reptans</i>		
<i>Caladenia rhomboidiformis</i> (E.Coleman) M.A.Clem. & Hopper	<i>Arachnorchis</i>	
<i>Caladenia roei</i> Benth.	<i>Jonesiopsis</i>	
<i>Caladenia saxicola</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia serotina</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia sigmoidea</i> R.S.Rogers	<i>Arachnorchis</i>	
<i>Caladenia speciosa</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	

Western Australian Species	Alternate Genus	Notes
<i>Caladenia splendens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia starteorum</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia straminichila</i> A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia swartsiorum</i> A.P.Br. & G.Brockman	<i>Arachnorchis</i>	
<i>Caladenia thinicola</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia uliginosa</i> A.S.George subspecies <i>candicans</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia uliginosa</i> A.S.George subspecies <i>patulens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia uliginosa</i> A.S.George subspecies <i>uliginosa</i>	<i>Arachnorchis</i>	
<i>Caladenia ultima</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia validinerva</i> Hopper & A.P.Br. ex A.P.Br. & G.Brockman	<i>Jonesiopsis</i>	
<i>Caladenia viridescens</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia voigtii</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia vulgata</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia wanosa</i> A.S.George	<i>Jonesiopsis</i>	
<i>Caladenia williamsiae</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia winfieldii</i> Hopper & A.P.Br.	<i>Arachnorchis</i>	
<i>Caladenia xantha</i> Hopper & A.P.Br.	<i>Jonesiopsis</i>	
<i>Caladenia</i> species 'Bindoon'	<i>Jonesiopsis</i>	aff. <i>hiemalis</i> with larger often yellowish-white flowers; granite hills between Bindoon and Gillingarra
<i>Caladenia</i> species 'east of Esperance'	<i>Arachnorchis</i>	late flowering aff. <i>hiemalis</i> with larger flowers from east of Esperance; brief mention in Brown <i>et al.</i> 2013
<i>Caladenia</i> species 'Wandoo'	<i>Jonesiopsis</i>	aff. <i>fluvialis</i> with slightly smaller flowers and upland habitat; woodland west of York and Brookton (A. Brown unpublished)
<i>Caladenia</i> species 'Watheroo'	<i>Jonesiopsis</i>	aff. <i>lorea</i> with creamy white flowers with red apex to labellum; heathy woodland near Watheroo (Backhouse 2011)
<i>Caladenia</i> species 'western wheatbelt'	<i>Jonesiopsis</i>	aff. <i>dimidia</i> with larger creamy white to yellow flowers with longer drooping segments with brighter red tails; dry woodland from York to Quairrading (A. Brown unpublished)
<i>Calochilus caesius</i> D.L.Jones		
<i>Calochilus holtzei</i> F.Muell.		
<i>Calochilus pruinosis</i> D.L.Jones		
<i>Calochilus stramenicola</i> D.L.Jones		
<i>Calochilus uliginosus</i> D.L.Jones		
<i>Calochilus</i> species 'Booyup Brook'		aff. <i>stramenicola</i> with different habitat and well-spaced flowers; heathy woodland near Booyup Brook (Brown <i>et al.</i> 2013)
<i>Calochilus</i> species 'Theda Station'		aff. <i>caesius</i> with larger flowers with longer basal hairs and a naked, cupped labellum apex; on sandy soils amongst sedges on seasonally wet sites in the central Kimberley region (Brown <i>et al.</i> 2013); <i>barbarossa</i> ms
<i>Corunastylis tepperi</i> (F.Muell. ex. Tepper) D.L.Jones & M.A.Clem.	<i>Genoplesium</i>	
<i>Corybas abditus</i> D.L.Jones	<i>Anzybas</i>	
<i>Corybas despectans</i> D.L.Jones & R.C.Nash	<i>Corysanthes</i>	
<i>Corybas limpidus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas recurvus</i> D.L.Jones	<i>Corysanthes</i>	
<i>Corybas</i> species 'Busselton'	<i>Corysanthes</i>	aff. <i>despectans</i> with tiny dorsal sepal; swampy habitats near Busselton (Brown <i>et al.</i> 2013)
<i>Corybas</i> species 'Hammersley Inlet'	<i>Corysanthes</i>	aff. <i>limpidus</i> with smaller flowers and a proportionately smaller dorsal sepal; under tea-tree on sandy soils around Hammersley Inlet (Fitzgerald River National Park) (Brown <i>et al.</i> 2013)
<i>Corybas</i> species 'Naturaliste'	<i>Corysanthes</i>	aff. <i>despectans</i> with very small self-pollinating flowers with incurved labellum margins and peaked dorsal sepal; low coastal heath south between Busselton and Cowaramup (Brown <i>et al.</i> 2013)
<i>Cryptostylis ovata</i> R.Br.		
<i>Cyanicula amplexans</i> (A.S.George) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula aperta</i> Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula ashbyae</i> Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula fragrans</i> Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula gemmata</i> (Lindl.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula gertrudeae</i> (Ostenf.) Hopper & A.P.Br.	<i>Caladenia</i>	

Western Australian Species	Alternate Genus	Notes
<i>Cyanicula ixiooides</i> (Lindl.) Hopper & A.P.Br. subspecies <i>candida</i> Lindl. Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula ixiooides</i> (Lindl.) Hopper & A.P.Br. subspecies <i>ixiooides</i>	<i>Caladenia</i>	
<i>Cyanicula nikulinskyae</i> Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula sericea</i> (Lindl.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Cyanicula</i> species 'Dale'	<i>Caladenia</i>	aff. <i>gemmata</i> with an all-green leaf and pale blue flowers with a prominently striped labellum; shrubby woodland adjacent to seasonally wet drainage lines, near Dale, flowering mostly after fire (Brown <i>et al.</i> 2013)
<i>Cyanicula</i> species 'Esperance'	<i>Caladenia</i>	aff. <i>gemmata</i> with purplish underside to leaf and purple and white labellum with tall calli, scrubs and woodland, also on granite outcrops, east from Esperance (Brown <i>et al.</i> 2013)
<i>Cymbidium canaliculatum</i> R.Br.		many forms and varieties have been named (eg. var. <i>sparksii</i> ) although none have any formal taxonomic recognition
<i>Cyrtostylis huegelii</i> Endl.		
<i>Cyrtostylis robusta</i> D.L.Jones & M.A.Clem.		
<i>Cyrtostylis tenuissima</i> (Nicholls & Goadby) D.L.Jones & M.A.Clem.		
<i>Cyrtostylis</i> species 'Cape Naturaliste'		short aff. <i>robusta</i> with smaller flowers and narrower labellum; coastal scrubs and heathy woodland on the Leeuwin-Naturaliste Ridge near Dunsborough (Brown <i>et al.</i> 2013)
<i>Dendrobium dicuphum</i> F.Muell.	<i>Vappodes</i>	
<i>Didymoplexis pallens</i> Griffith		
<i>Dipodium stenochilum</i> O.Schwarz		it is likely that plants identified as <i>D. stenochilum</i> in WA will be redetermined as <i>Dipodium</i> species 'basalt woodland' and <i>Dipodium</i> species 'sandstone'
<i>Dipodium</i> species 'basalt woodland'		aff. <i>roseum</i> with spotted deep pink flowers with a rose pink labellum; woodland in the Kimberely region; <i>basalticum</i> ms (Brown <i>et al.</i> 2013)
<i>Dipodium</i> species 'sandstone'		aff. <i>roseum</i> with unspotted pale pink flowers with a rose pink labellum; woodland over sandstone in the Kimberely region; <i>ammolithum</i> ms (Brown <i>et al.</i> 2013)
<i>Diuris amplissima</i> D.L.Jones		
<i>Diuris brumalis</i> D.L.Jones		
<i>Diuris carinata</i> Lindl.		
<i>Diuris concinna</i> D.L.Jones		
<i>Diuris conspicillata</i> D.L.Jones		
<i>Diuris corymbosa</i> Lindl.		
<i>Diuris decrementa</i> D.L.Jones & C.J.French		
<i>Diuris drummondii</i> Lindl.		
<i>Diuris eburnea</i> D.L.Jones		
<i>Diuris emarginata</i> R.Br.		
<i>Diuris filifolia</i> Lindl.		
<i>Diuris hazeliae</i> D.L.Jones & C.J.French		
<i>Diuris heberlei</i> D.L.Jones		
<i>Diuris immaculata</i> D.L.Jones		
<i>Diuris insignis</i> D.L.Jones & C.J.French		
<i>Diuris jonesii</i> C.J.French & G.Brockman		
<i>Diuris laevis</i> Fitzg.		
<i>Diuris laxiflora</i> Lindl.		
<i>Diuris longifolia</i> R.Br.		
<i>Diuris magnifica</i> D.L.Jones		
<i>Diuris micrantha</i> D.L.Jones		
<i>Diuris pauciflora</i> R.Br.		
<i>Diuris perialla</i> D.L.Jones & C.J.French		
<i>Diuris picta</i> J. Drummond		
<i>Diuris porrifolia</i> Lindl.		
<i>Diuris pulchella</i> D.L.Jones		
<i>Diuris purdiei</i> Diels		
<i>Diuris recurva</i> D.L.Jones		
<i>Diuris refracta</i> D.L.Jones & C.J.French		
<i>Diuris segregata</i> D.L.Jones & C.J.French		
<i>Diuris septentrionalis</i> D.L.Jones & C.J.French		
<i>Diuris setacea</i> R.Br.		
<i>Diuris tinkeri</i> D.L.Jones & C.J.French		

Western Australian Species	Alternate Genus	Notes
<i>Diuris</i> species 'Augusta'		late-flowering aff. <i>laxiflora</i> with bright yellow flowers with a broader labellum mid-lobe; amongst sedges in seasonally damp habitats, flowering best after fire; only near Augusta (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Darling scarp'		late spring flowering, tall aff. <i>amplissima</i> with large brownish-yellow and mauve flowers (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Green Range'		winter flowering aff. <i>corymbosa</i> with colourful flowers; heaths and heathy woodlands on south coast between Denmark and Esperance (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Kemerton'		aff. <i>corymbosa</i> with yellow and red-brown flowers; coastal woodland on sandy soils between Lake Clifton and Capel (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'mid-north'		small, often clumping aff. <i>corymbosa</i> with pale flowers with recurved dorsal sepal and crossed, recurved lateral sepals; heath and woodland between Moora and Mingenew (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Murchison River'		tall aff. <i>corymbosa</i> with mainly yellow flowers with recurved petals and sepals; moist sedgeland near Murchison River (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Perth swamps'		small, short aff. <i>corymbosa</i> with widely spaced flowers with a short labellum midlobe; swamps just south of Perth (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'sandplain'		tall, colourful aff. <i>corymbosa</i> ; sandplain habitats between Mandurah and Bunbury (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Scaddan'		small, winter flowering aff. <i>corymbosa</i> ; woodland on rocky outcrops between Hyden and Scaddan; included in Brown <i>et al.</i> 2008 but not in later books
<i>Diuris</i> species 'south coast'		winter-flowering aff. <i>corymbosa</i> with small, mainly yellow flowers; coastal heaths and heathy woodland between Munglinup and Denmark (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'western wheatbelt'		short aff. <i>corymbosa</i> with pale yellow flowers; woodlands and rocky areas between York and Ravensthorpe (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Williams'		small aff. <i>corymbosa</i> ; <i>Allocasuarina</i> woodland thickets near Williams; included in Brown <i>et al.</i> 2008 but not in later books
<i>Diuris</i> species 'Wyalkatchem'		early spring-flowering aff. <i>corymbosa</i> with yellow flowers with small red patches; inland woodlands on sandy soils between Bencubbin and Wyalkatchmen (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Yalgorup'		tall colourful aff. <i>corymbosa</i> ; woodland and open forest between and Mandurah and Bunbury (Brown <i>et al.</i> 2013)
<i>Diuris</i> species 'Zuytdorp cliffs'		smallish, compact aff. <i>corymbosa</i> with mainly yellow flowers with short lateral sepals; coastal sedges on limestone sands around Kalbarri (Brown <i>et al.</i> 2013)
<i>Drakaea andrewsiae</i> Hopper & A.P.Br.		
<i>Drakaea concolor</i> Hopper & A.P.Br.		
<i>Drakaea confluens</i> Hopper & A.P.Br.		
<i>Drakaea elastica</i> Lindl.		
<i>Drakaea glyptodon</i> Fitzg.		
<i>Drakaea gracilis</i> Hopper & A.P.Br.		
<i>Drakaea isolata</i> Hopper & A.P.Br.		
<i>Drakaea livida</i> J.Drummond		
<i>Drakaea micrantha</i> Hopper & A.P.Br.		
<i>Drakaea thynniphila</i> A.S.George		
<i>Elythranthera brunonis</i> (Endl.) A.S.George	<i>Caladenia</i>	
<i>Elythranthera emarginata</i> (Lindl.) A.S.George	<i>Caladenia</i>	
<i>Empusa habenarina</i> (F.Muell.) Benth.	<i>Liparis</i>	
<i>Epiblema grandiflorum</i> R.Br.		
<i>Ericksonella saccharata</i> (Rchb.f.) Hopper & A.P.Br.	<i>Caladenia</i>	
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>brevifolius</i> (Benth.) Hopper & A.P.Brown		
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>dilatatus</i>		
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>magnus</i> Hopper & A.P.Br.		
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>multiflorus</i> Hopper & A.P.Br.		
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>orientalis</i> Lindl.		
<i>Eriochilus dilatatus</i> Lindl. subspecies <i>undulatus</i> Hopper & A.P.Br.		
<i>Eriochilus helonomos</i> Hopper & A.P.Br.		
<i>Eriochilus pulchellus</i> Hopper & A.P.Br.		
<i>Eriochilus scaber</i> Lindl. subspecies <i>orbifolia</i> Hopper & A.P.Br.		
<i>Eriochilus scaber</i> Lindl. subspecies <i>scaber</i>		
<i>Eriochilus tenuis</i> Lindl.		

Western Australian Species	Alternate Genus	Notes
<i>Eriochilus valens</i> Hopper & A.P.Br.		
<i>Eulophia bicallosa</i> (D.Don) P.F.Hunt & Summerh.		
<i>Gastrodia lacista</i> D.L.Jones		
<i>Geodorum densiflorum</i> (Lam.) Schltr.		also called <i>Geodorum terrestre</i> (L.) Garay
<i>Geodorum neocaledonicum</i> Kraenzl.		record most likely false, probably resulting from an incorrect determination of <i>Eulophia bicallosa</i>
<i>Habenaria elongata</i> R.Br.		
<i>Habenaria hymenophylla</i> Schltr.		
<i>Habenaria ochroleuca</i> R.Br.		
<i>Habenaria</i> species 'Beverley Springs Station'		aff. <i>praecox</i> with a bluntly pointed, tri-lobed labellum; damp sites in woodland in the Kimberley region; <i>joesmithii</i> ms (Brown <i>et al.</i> 2013)
<i>Leporella fimbriata</i> (Lindl.) A.S.George		
<i>Leptoceras menziesii</i> (R.Br.) Lindl.	<i>Caladenia</i>	
<i>Lyperanthus serratus</i> Lindl.		
<i>Microtis alba</i> R.Br.		
<i>Microtis alboviridis</i> R.J.Bates		
<i>Microtis atrata</i> Lindl.	<i>Microtidium</i>	
<i>Microtis brownii</i> Rchb.f.		
<i>Microtis cupularis</i> (D.L.Jones & G.Brockman) A.P.Br.	<i>Hydrorchis</i>	
<i>Microtis eremaea</i> R.J.Bates		
<i>Microtis eremicola</i> (R.J.Bates) D.L.Jones & M.A.Clem.		
<i>Microtis familiaris</i> R.J.Bates		
<i>Microtis globula</i> R.J.Bates		
<i>Microtis granitica</i> R.J.Bates		
<i>Microtis media</i> R.Br. subspecies <i>densiflora</i> R.J.Bates		
<i>Microtis media</i> R.Br. subspecies <i>media</i>		
<i>Microtis orbicularis</i> R.S.Rogers	<i>Hydrorchis</i>	
<i>Microtis pulchella</i> R.Br.		
<i>Microtis quadrata</i> (R.J.Bates) D.L.Jones & M.A.Clem.		
<i>Microtis</i> species 'Windy Harbour'	<i>Hydrorchis</i>	summer flowering aff. <i>orbicularis</i> with smaller, more crowded reddish flowers with cupped labellum; seasonal swamps between Northcliff and Windy Harbour (Brown <i>et al.</i> 2013)
<i>Nervilia holochila</i> (F.Muell.) Schltr.		
<i>Paracaleana alcockii</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana brockmanii</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana disjuncta</i> D.L.Jones	<i>Caleana</i>	
<i>Paracaleana dixonii</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana gracilicordata</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana granitica</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana hortiorum</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana lyonsii</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana nigrata</i> J. Drummond ex Lindl.	<i>Caleana</i>	
<i>Paracaleana parvula</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana terminalis</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana triens</i> Hopper & A.P.Br.	<i>Caleana</i>	
<i>Paracaleana</i> species 'Darling Range'	<i>Caleana</i>	aff. <i>nigrata</i> with leaf still green at flowering and flat labellum with calli covering apical half; grows with <i>P. brockmanii</i> but flowers earlier and has smaller flower; woodland on lateritic soils near Kelmscott (Brown <i>et al.</i> 2013)
<i>Pheladenia deformis</i> (R.Br.) D.L.Jones & M.A.Clem.	<i>Caladenia</i>	
<i>Praecoxanthus aphyllus</i> (Benth.) Hopper & A.P.Br.		
<i>Prasophyllum brownii</i> Rchb.f.		
<i>Prasophyllum calcicola</i> R.J.Bates		
<i>Prasophyllum cucullatum</i> Rchb.f.	<i>Chiloterus</i>	
<i>Prasophyllum cyphochilum</i> Benth.		
<i>Prasophyllum drummondii</i> Rchb.f.		
<i>Prasophyllum elatum</i> R.Br.		
<i>Prasophyllum fimbria</i> Rchb.f.		
<i>Prasophyllum gibbosum</i> R.Br.	<i>Chiloterus</i>	
<i>Prasophyllum giganteum</i> Lindl.		
<i>Prasophyllum gracile</i> Lindl.		



Western Australian Species	Alternate Genus	Notes
<i>Prasophyllum hians</i> Rchb.f.		
<i>Prasophyllum macrostachyum</i> R.Br.		
<i>Prasophyllum macrotys</i> Lindl.		
<i>Prasophyllum odoratissimum</i> D.L.Jones		
<i>Prasophyllum ovale</i> Lindl.		
<i>Prasophyllum parvifolium</i> Lindl.	<i>Mecopodum</i>	
<i>Prasophyllum paulineae</i> D.L.Jones & M.A.Clem.		
<i>Prasophyllum plumiforme</i> Fitzg.		
<i>Prasophyllum regium</i> R.S.Rogers		
<i>Prasophyllum sargentii</i> (Nicholls) A.S.George		
<i>Prasophyllum triangulare</i> Fitzg.		
<i>Prasophyllum</i> species 'Brookton Highway'	<i>Mecopodum</i>	late winter-early spring flowering aff. <i>parvifolium</i> with few small, poorly opening, self-pollinating flowers; swampy sites and moist margins of granite outcrops from Aramadale to Brookton (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'crowded'		aff. <i>elatum</i> with crowded spike of poorly opening, self-pollinating flowers; woodland on sandy rises above seasonally wet flats between Perth and Ravensthorpe, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'early'	<i>Mecopodum</i>	autumn flowering aff. <i>parvifolium</i> with scented green and white flowers lacking red stripes; heath, woodland, forest and granite outcrops between Bunbury and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'eastern wheatbelt'		aff. <i>elatum</i> with smaller, dark flowers with a less ruffled labellum margin; flowers after fire and on lightly disturbed ground (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'late swamps'		late flowering aff. <i>drummondii</i> with smaller, darker, more widely spaced flowers; on seasonally wet flats between Augusta and Israelite Bay, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'southern'		short, late spring-early summer flowering aff. <i>fimbria</i> with smaller darker flowers; seasonally wet flats between Augusta and Israelite Bay, flowering after summer fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'striped'	<i>Chiloterus</i>	aff. <i>gibbosum</i> with green and white flowers with humped base to the lateral sepals; seasonally wet flats between Perth and Busselton, flowering only after fire (Brown <i>et al.</i> 2013)
<i>Prasophyllum</i> species 'Swan region'		aff. <i>elatum</i> with smaller paler flowers with a less ruffled labellum margin; woodland in rocky hills of the Darling Scarp between Greenmount and Aramadale, flowering only after summer fire (Brown <i>et al.</i> 2013)
<i>Pterostylis allantoidea</i> R.S.Rogers	<i>Eremorchis</i>	
<i>Pterostylis angusta</i> A.S.George	<i>Diplodium</i>	
<i>Pterostylis aspera</i> D.L.Jones & M.A.Clem.	<i>Diplodium</i>	
<i>Pterostylis barbata</i> Lindl.	<i>Plumatichilos</i>	
<i>Pterostylis brevichila</i> D.L.Jones & C.J.French	<i>Diplodium</i>	
<i>Pterostylis brunneola</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis ciliata</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis concava</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	
( <i>Pterostylis</i> ) <i>crebriflorus</i> (D.L.Jones & C.J.French)	<i>Urochilus</i>	in describing <i>crebriflorus</i> , Jones & French have taken a narrower interpretation than that of Brown <i>et al.</i> 2013 for <i>P.</i> species 'crowded'; described in <i>Urochilus</i> , new combination in <i>Pterostylis</i> not yet made
<i>Pterostylis dilatata</i> A.S.George	<i>Linguella</i>	
<i>Pterostylis echinulata</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis elegantissima</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis erubescens</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis exserta</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis frenchii</i> (D.L.Jones) A.P.Br.	<i>Oligochaetochilus</i>	
<i>Pterostylis glebosa</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis hadra</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis hamiltonii</i> Nicholls	<i>Diplodium</i>	
<i>Pterostylis insectifera</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis jacksonii</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis karri</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis leptochila</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis lortensis</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis macrocalymma</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis macrosceles</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	

Western Australian Species	Alternate Genus	Notes
<i>Pterostylis microglossa</i> D.L.Jones & C.J.French	<i>Diplodium</i>	
<i>Pterostylis microphylla</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis mutica</i> R.Br.	<i>Hymenochilus</i>	
<i>Pterostylis parva</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis perculata</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis picta</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis platypetala</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis pyramidalis</i> Lindl.	<i>Linguella</i>	
<i>Pterostylis recurva</i> Benth.	<i>Stamnorchis</i>	
<i>Pterostylis roensis</i> M.A.Clem. & D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis rogersii</i> E.Coleman	<i>Diplodium</i>	
<i>Pterostylis sanguinea</i> D.L.Jones & M.A.Clem.	<i>Urochilus</i>	
<i>Pterostylis sargentii</i> C.R.P.Andrews	<i>Ranorchis</i>	
<i>Pterostylis scabra</i> Lindl.	<i>Diplodium</i>	
<i>Pterostylis scitula</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis sinuata</i> (D.L.Jones) Janes & Duretto	<i>Hymenochilus</i>	
<i>Pterostylis spathulata</i> M.A.Clem.	<i>Oligochaetochilus</i>	
<i>Pterostylis telmata</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis timothyi</i> (D.L.Jones) Janes & Duretto	<i>Linguella</i>	
<i>Pterostylis turfosa</i> Endl.	<i>Plumatichilos</i>	
<i>Pterostylis vittata</i> Lindl.	<i>Urochilus</i>	
<i>Pterostylis voigtii</i> D.L.Jones & C.J.French	<i>Linguella</i>	
<i>Pterostylis zebrina</i> (D.L.Jones & C.J.French) D.L.Jones	<i>Oligochaetochilus</i>	
<i>Pterostylis</i> species 'bloated'	<i>Plumatichilos</i>	late flowering aff. <i>barbata</i> with larger more bloated flowers; damp open forest and seasonally wet flats between Augusta and Mt Barker (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'bloated snail'	<i>Linguella</i>	aff. <i>nana</i> with large 'bloated flowers; open forest on sandy soils between Armadale and Margaret River; included in Brown <i>et al.</i> 2008 but not in later books; possibly just a broad flowered form of <i>P.</i> species 'crinkled leaf'
<i>Pterostylis</i> species 'Boyup'	<i>Plumatichilos</i>	early flowering aff. <i>plumosa</i> with light green flowers with a shorter, upturned dorsal sepal tip and a more densely bristled labellum; woodland between Augusta and Tenderden (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Cape Le Grande'	<i>Linguella</i>	aff. <i>nana</i> with broad fleshy leaves in rosette and uniformly thickened lateral sepals; coastal scrubs along moist drainage lines and on granite rock sheets between Esperance and Cape Arid (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'cauline leaves'	<i>Linguella</i>	aff. <i>nana</i> with fleshy rosette, robust habit with short thickened lateral sepals; woodland and open forest on sandy soils between Perth and Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'chocolate'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> type with dark brown flowers; in shallow soil pockets on granite outcrops near Karonia (Brown <i>et al.</i> 2008); not included in later books but still thought to be distinct
<i>Pterostylis</i> species 'Chudalup'	<i>Linguella</i>	aff. <i>nana</i> with sparsely leafy rosette, short stem with narrow flower with short clubbed lateral sepals; moss mats on granite rocks near Northcliff (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'coastal'	<i>Urochilus</i>	aff. <i>vitatta</i> with smaller, more widely spaced flowers; scrubs and woodland from Dongara to Bunbury (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'coastal clubbed sepals'	<i>Linguella</i>	aff. <i>nana</i> with short stem and bloated flowers with very short clubbed lateral sepals; widespread on south-west coast, in scrubs, heath and woodland on sandy soils (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'coastal snail'	<i>Linguella</i>	aff. <i>nana</i> with short stem and somewhat bloated flowers with short thickened lateral sepals; coastal scrubs on sand dunes between Bremer Bay and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'crinkled leaf'	<i>Linguella</i>	aff. <i>nana</i> with crinkled leaves and tall slender flower with pointed galea and pointed tips to the lateral sepals; widely distributed between Perth and Albany, inland to Hyden, in woodland and open forest (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'dainty brown'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with small reddish brown with an extensively translucent hood and short forward-curved tips to the lateral sepals; open woodland on rocky hills and around salt lakes between Kalbarri and Eyre (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'dwarf'	<i>Plumatichilos</i>	aff. <i>barbata</i> with small dark green flowers with short sepal tips; widespread in shrublands and mallee heath across inland sth-west WA (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Esperance'	<i>Plumatichilos</i>	short, small aff. <i>plumosa</i> with light green flowers with long galea point and long dense bristles on the labellum; mallee woodland, shrubland and on granite rocks between Munghlinup and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Eyre'	<i>Urochilus</i>	aff. <i>vitatta</i> with fewer smaller paler flowers; shrubland and woodland on sand dunes, limestone cliffs south of Cocklebidy (Brown <i>et al.</i> 2013)

Western Australian Species	Alternate Genus	Notes
<i>Pterostylis</i> species 'fine sepals'	<i>Linguella</i>	aff. <i>nana</i> with small flowers with very long slender lateral sepals; wandoo woodland between Perth and Bindoon (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'granite bird'	<i>Plumatichilos</i>	aff. <i>plumosa</i> with very long sepal free points and the dorsal sepal tip upcurved; shrubland on granite outcrops between Albany and Shannon (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'greasy'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with smaller, nodding, 'greasy' reddish green to greenish flowers with a short upcurved tip to the dorsal sepal; woodland and shrubland between Wubin and Norseman (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'green flowers'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with large, fairly crowded, nodding green flowers with a large spoon-shaped labellum; shrubland on granite outcrops between Salmon Gums and Menzies (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Greenbushes'	<i>Plumatichilos</i>	aff. <i>barbata</i> with large pale flowers constricted and flattened in middle, a broad lower opening & sigmoid-shaped dorsal sepal; woodland and open forest between Dardanup and Augusta (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Gull Rock'	<i>Plumatichilos</i>	aff. <i>barbata</i> with long narrow flowers with very long sepal free points; heathy woodland and coastal scrubs near Albany (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Helena River'	<i>Linguella</i>	aff. <i>nana</i> with large broad flowers with extended dorsal sepal and narrowly clubbed lateral sepals; winter-wet sites in woodland and coastal heaths, Perth to Dunsborough (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'inland'	<i>Linguella</i>	tall aff. <i>nana</i> with a hairy stem; widespread in inland areas in woodland and shrublands (Brown <i>et al.</i> 2013); uncertain how it differs from and is almost certainly the same as <i>P. nana</i>
<i>Pterostylis</i> species 'large fawn'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with large tawny nodding flowers with forward-pointing lateral sepal tips and a broad, ovate labellum; dry woodland and shrubland on rocky soils, also granite outcrops near Karonie (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'late flowering'	<i>Linguella</i>	late spring-flowering aff. <i>nana</i> with short stems and clubbed lateral sepals; damp open forest near Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'limestone sands'	<i>Linguella</i>	tall aff. <i>nana</i> with large flowers; near coastal woodland on well-drained calcareous sands from Yanchep to Busselton; included in Hoffman & Brown 2011 although not in later books; very similar to <i>P. sp.</i> 'Helena River'
<i>Pterostylis</i> species 'long sepals'	<i>Linguella</i>	short, small aff. <i>nana</i> with long narrow lateral sepals and small, compact rosette of pointed leaves; moss mats on granite outcrops between Denmark and Windy Harbour (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Mehinup'	<i>Plumatichilos</i>	aff. <i>barbata</i> with a very sparsely hairy labellum and widely spreading lateral sepals with upturned tips; on and around granite outcrops, Denmark to Walpole (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Muir Highway'	<i>Plumatichilos</i>	aff. <i>barbata</i> with very long sepal free points and labellum bristles in opposite pairs; woodland on sandy soils along the Muir Highway (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Murchison'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with nodding flowers with bright red lateral sepals with green inner margins and forward curved tips; dry woodland and shrubland on sandy soils between Murchison and Shark Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'northern'	<i>Urochilus</i>	short aff. <i>vitatta</i> with fewer smaller flowers; dense shrubland on sandy soils between Cataby and Binu (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'northern thick sepals'	<i>Linguella</i>	aff. <i>nana</i> with broad fleshy rosette and short uniformly thickened lateral sepals; wandoo woodland between Perth and Cataby (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Paynes Find'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with smallish nodding red-striped flowers with cupped lateral sepals with forward-pointing tips; sparse shrublands on granite outcrops and rocky breakaways between Wubin and Mount Magnet (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'plumed'	<i>Plumatichilos</i>	aff. <i>plumosa</i> with smallish, dark green flowers with very short free point on dorsal sepal; woodland and shrubland between Esperance and Israelite Bay (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'Ravensthorpe'	<i>Linguella</i>	aff. <i>nana</i> with moderately tall stem and large bloated flowers with moderately long lateral sepals; open woodland and shrubland between the Stirling Range and Esperance (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'robust'	<i>Linguella</i>	robust, spring flowering aff. <i>nana</i> with a blunt galea and thickened lateral sepals; damp sites in open forest on heavy clay soils (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'short sepals'	<i>Linguella</i>	aff. <i>nana</i> with short stem and narrow flowers with short lateral sepals; heath and heathy woodland on sandy soils between Perth and Yanchep (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'skinny'	<i>Linguella</i>	aff. <i>nana</i> with tall, slender habit and tall, narrow flowers with long slender lateral sepals; open forest and woodland between Perth and Mt Barker (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'small bands'	<i>Urochilus</i>	aff. <i>vitatta</i> with fewer, smaller flowers with rounded lateral sepals; shrublands and woodlands between Reagans Ford and Three Springs (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'southern granites'	<i>Linguella</i>	aff. <i>nana</i> with small rosette, short stem and long lateral sepals; moss mats on granite outcrops between Manjimup and Hopetoun (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'sparse'	<i>Plumatichilos</i>	aff. <i>barbata</i> with very sparse bristles on labellum; woodland and open forest between Dardanup and Walpole (Brown <i>et al.</i> 2013)

Western Australian Species	Alternate Genus	Notes
<i>Pterostylis</i> species 'straight tops'	<i>Oligochaetochilus</i>	aff. <i>rufa</i> with nodding mostly translucent flowers with almost solid red lateral sepals with straight upper margins and moderately long tips; widely distributed in inland woodland and shrublands often on rocky areas (Brown <i>et al.</i> 2013)
<i>Pterostylis</i> species 'The Knoll'	<i>Plumatichilos</i>	late flowering aff. <i>plumosa</i> with very long bristles on labellum; woodland and open forest between Albany and Dardanup (Brown <i>et al.</i> 2013)
<i>Pyrorchis forrestii</i> (F.Muell.) D.L.Jones & M.A.Clem.		
<i>Pyrorchis nigricans</i> (R.Br.) D.L.Jones & M.A.Clem.		
<i>Rhizanthella gardneri</i> R.S.Rogers		
<i>Rhizanthella</i> species 'Munglinup'		winter flowering aff. <i>gardneri</i> with fewer, larger, paler coloured flowers; under dense <i>Melaleuca</i> scrubs on sandy loam soils near Munglinup; <i>johnstonii</i> MS (Brown <i>et al.</i> 2013)
<i>Spiculaea ciliata</i> Lindl.		
<i>Thelymitra antennifera</i> (Lindl.) Hook.f.		
<i>Thelymitra apiculata</i> (A.S.George) M.A.Clem. & D.L.Jones		
<i>Thelymitra benthamiana</i> Rchb.f.		
<i>Thelymitra campanulata</i> Lindl.		
<i>Thelymitra canaliculata</i> R.Br.		
<i>Thelymitra cornicina</i> Rchb.f.		
<i>Thelymitra crinita</i> Lindl.		
<i>Thelymitra cucullata</i> Rupp		
<i>Thelymitra dedmaniarum</i> R.S.Rogers		
<i>Thelymitra flexuosa</i> Endl.		
<i>Thelymitra frenchii</i> Jeanes		virtually identical to and very difficult to separate from <i>T. granitora</i>
<i>Thelymitra fuscolutea</i> R.Br.		
<i>Thelymitra graminea</i> Lindl.		
<i>Thelymitra granitora</i> D.L.Jones & M.A.Clem.		
<i>Thelymitra jacksonii</i> Hopper & A.P.Brown ex Jeanes		
<i>Thelymitra latiloba</i> Jeanes		
<i>Thelymitra macrophylla</i> Lindl.		
<i>Thelymitra maculata</i> Jeanes		
<i>Thelymitra magnifica</i> Jeanes		
<i>Thelymitra mucida</i> Fitzg.		
<i>Thelymitra occidentalis</i> Jeanes		
<i>Thelymitra paludosa</i> Jeanes		
<i>Thelymitra petrophila</i> Jeanes		
<i>Thelymitra psammophila</i> C.R.P.Andrews		
<i>Thelymitra pulcherrima</i> Jeanes		
<i>Thelymitra sargentii</i> R.S.Rogers		
<i>Thelymitra speciosa</i> Jeanes		
<i>Thelymitra spiralis</i> (Lindl.) F.Muell.		
<i>Thelymitra stellata</i> Lindl.		
<i>Thelymitra tigrina</i> R.Br.		
<i>Thelymitra uliginosa</i> Jeanes		
<i>Thelymitra variegata</i> (Lindl.) F.Muell.		
<i>Thelymitra villosa</i> Lindl.		
<i>Thelymitra vulgaris</i> Jeanes		
<i>Thelymitra xanthotricha</i> Jeanes		
<i>Thelymitra yorkensis</i> Jeanes		
<i>Thelymitra</i> species 'Murchison'		aff. <i>macrophylla</i> with clumping habit and smaller, paler flowers; seasonally moist heathlands between Geraldton and Nerrin Nerrin (Brown <i>et al.</i> 2013)
<i>Thelymitra</i> species 'northern swamps'		short, early-flowering aff. <i>paludosa</i> ; swamps between Perth and Bunbury (Brown <i>et al.</i> 2008 but not in later books); may just be a northern form of <i>T. paludosa</i>
<i>Thelymitra</i> species 'Ongerup'		aff. <i>yorkensis</i> with solid bicoloured flowers and more ornate column wings; heathy woodland near Ongerup (Brown <i>et al.</i> 2013)
<i>Thelymitra</i> species 'reed sun orchid'		large aff. <i>petrophila</i> with a longer leaf; called aff. <i>macrophylla</i> in Hoffman & Brown 2011; grows amongst sedges not reeds; possibly just a robust form of <i>T. petrophila</i>
<i>Zeuxine oblonga</i> R.S.Rogers & C.T.White		

**Total: 488 taxa (species, subspecies)**

Western Australian Hybrids	Name and Authority	Notes
<i>Caladenia applanata</i> X <i>attingens</i>		A. Brown personal observation
<i>Caladenia applanata</i> X <i>infundibularis</i>		A. Brown personal observation
<i>Caladenia applanata</i> X <i>nivalis</i>		A. Brown personal observation
<i>Caladenia arenicola</i> X <i>huegelii</i>		A. Brown personal observation
<i>Caladenia arenicola</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia arenicola</i> X <i>lorea</i>		Jones 2006
<i>Caladenia arenicola</i> X <i>swartziorum</i>		Brown & Brockman 2015
<i>Caladenia attingens</i> X <i>barbarossa</i>		Jones 2006
<i>Caladenia attingens</i> X <i>exstans</i>		Jones 2006
<i>Caladenia attingens</i> X <i>georgei</i>		Backhouse 2011
<i>Caladenia attingens</i> X <i>horistes</i>		Brown & Brockman 2015
<i>Caladenia attingens</i> X <i>infundibularis</i>		Brown <i>et al.</i> 2013
<i>Caladenia attingens</i> X <i>longicauda</i>		Brown & Brockman 2015
<i>Caladenia attingens</i> X <i>procera</i>		Jones 2006
<i>Caladenia attingens</i> X <i>thinicola</i>		Jones 2006
<i>Caladenia barbarossa</i> X <i>caesarea</i>		Brown <i>et al.</i> 2013
<i>Caladenia barbarossa</i> X <i>falcata</i>		Backhouse 2011
<i>Caladenia barbarossa</i> X <i>lobata</i>		A. Brown personal observation
<i>Caladenia barbarossa</i> X <i>longicauda</i>	<i>×enigma</i> Hopper & A.P. Br.	
<i>Caladenia barbarossa</i> X <i>radiata</i>		Brown <i>et al.</i> 2013
<i>Caladenia brevisura</i> X <i>cruscula</i>		Brown <i>et al.</i> 2013
<i>Caladenia brevisura</i> X <i>doutchiae</i>		Jones 2006
<i>Caladenia brevisura</i> X <i>horistes</i>		Brown <i>et al.</i> 2008
<i>Caladenia brevisura</i> X <i>polychroma</i>		Brown <i>et al.</i> 2008
<i>Caladenia brevisura</i> X <i>voigtii</i>		Jones 2006
<i>Caladenia brownii</i> X <i>ferruginea</i>		A. Brown personal observation
<i>Caladenia brownii</i> X <i>longicauda</i>		Backhouse 2011
<i>Caladenia brownii</i> X <i>pectinata</i>		A. Brown personal observation
<i>Caladenia brownii</i> X <i>serotina</i>		G. Backhouse personal observation
<i>Caladenia busselliana</i> X <i>paludosa</i>		Backhouse 2011
<i>Caladenia busselliana</i> X <i>procera</i>		A. Brown personal observation
<i>Caladenia caesarea</i> X <i>polychroma</i>		A. Brown personal observation
<i>Caladenia cairnsiana</i> X <i>dimidia</i>	<i>×ericksoniae</i> Nicholls	
<i>Caladenia cairnsiana</i> X <i>filifera</i>		Jones 2006
<i>Caladenia cairnsiana</i> X <i>polychroma</i>		A. Brown personal observation
<i>Caladenia cairnsiana</i> X <i>pulchra</i>		Brown <i>et al.</i> 2008
<i>Caladenia cairnsiana</i> X <i>vulgata</i>		Jones 2006
<i>Caladenia chapmanii</i> X <i>longicauda</i>	<i>×eludens</i> Hopper & A.P. Br.	
<i>Caladenia chapmanii</i> X <i>polychroma</i>		Jones 2006
<i>Caladenia christineae</i> X <i>harringtoniae</i>		Brown <i>et al.</i> 2008
<i>Caladenia christineae</i> X <i>latifolia</i>		Brown <i>et al.</i> 2013
<i>Caladenia citrina</i> X <i>rhomboidiformis</i>		Brown <i>et al.</i> 2013
<i>Caladenia corynephora</i> X <i>serotina</i>	<i>×aestantha</i> Hopper & A.P. Br.	
<i>Caladenia crebra</i> X <i>longicauda</i>	<i>×coactescens</i> Hopper & A.P. Br.	
<i>Caladenia cristata</i> X <i>denticulata</i>		Brown <i>et al.</i> 2013
<i>Caladenia cristata</i> X <i>vulgata</i>		Jones 2006
<i>Caladenia decora</i> X <i>longicauda</i>		Backhouse 2011
<i>Caladenia denticulata</i> X <i>filifera</i>		A. Brown personal observation
<i>Caladenia denticulata</i> X <i>footeana</i>		Brown & Brockman 2015
<i>Caladenia dimidia</i> X <i>doutchiae</i>		Backhouse 2011
<i>Caladenia dimidia</i> X <i>erythronema</i>		A. Brown personal observation
<i>Caladenia dimidia</i> X <i>filifera</i>		A. Brown personal observation
<i>Caladenia dimidia</i> X <i>footeana</i>		A. Brown personal observation
<i>Caladenia dimidia</i> X <i>hirta</i>		A. Brown personal observation
<i>Caladenia dimidia</i> X <i>melanema</i>		A. Brown personal observation
<i>Caladenia dimidia</i> X <i>pachychila</i>		Brown <i>et al.</i> 2013
<i>Caladenia dimidia</i> X <i>paradoxa</i>		Jones 2006
<i>Caladenia dimidia</i> X <i>polychroma</i>		Jones 2006
<i>Caladenia dimidia</i> X <i>radialis</i>		Backhouse 2011

Western Australian Hybrids	Name and Authority	Notes
<i>Caladenia dimidia</i> X <i>roei</i>	× <i>exoleta</i> Hopper & A.P. Br.	
<i>Caladenia discoidea</i> X <i>hirta</i>		Brown <i>et al.</i> 2013
<i>Caladenia dorrienii</i> X <i>vulgata</i>		A. Brown personal observation
<i>Caladenia doutchiae</i> X <i>exilis</i>		Jones 2006
<i>Caladenia doutchiae</i> X <i>incrassata</i>		Jones 2006
<i>Caladenia doutchiae</i> X <i>polychroma</i>		Jones 2006
<i>Caladenia doutchiae</i> X <i>pulchra</i>		Jones 2006
<i>Caladenia doutchiae</i> X <i>radialis</i>		Backhouse 2011
<i>Caladenia doutchiae</i> X <i>vulgata</i>		A. Brown personal observation
<i>Caladenia drakeoides</i> X <i>exilis</i> subsp. <i>exilis</i>	× <i>hopperi</i> J.M.H.Shaw	the original name × <i>ornata</i> Hopper & A.P.Br. was invalid as <i>Caladenia ornata</i> was already named
<i>Caladenia drakeoides</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia drakeoides</i> X <i>vulgata</i>		Jones 2006
<i>Caladenia dundasiae</i> X <i>vulgata</i>		A. Brown personal observation
<i>Caladenia elegans</i> X <i>nobilis</i>		Jones 2006
<i>Caladenia elegans</i> X <i>vulgata</i>		Brown <i>et al.</i> 2013
<i>Caladenia erythrochila</i> X <i>straminichila</i>		A. Brown personal observation
<i>Caladenia erythronema</i> X <i>polychroma</i>		A. Brown personal observation
<i>Caladenia erythronema</i> X <i>vulgata</i>		A. Brown personal observation
<i>Caladenia excelsa</i> X <i>longicauda</i>		Jones 2006
<i>Caladenia exilis</i> X <i>roei</i>		Jones 2006
<i>Caladenia exstans</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia falcata</i> X <i>integra</i>		A. Brown personal observation
<i>Caladenia falcata</i> X <i>longicauda</i>	× <i>cala</i> Hopper & A.P. Br.	
<i>Caladenia ferruginea</i> X <i>leucochila</i>		Brown & Brockman 2015
<i>Caladenia ferruginea</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia ferruginea</i> X <i>pectinata</i>		Brown <i>et al.</i> 2008
<i>Caladenia ferruginea</i> X <i>rhomboidiformis</i>		Brown <i>et al.</i> 2013
<i>Caladenia filifera</i> X <i>hiemalis</i>		Brown <i>et al.</i> 2013
<i>Caladenia flava</i> X <i>hirta</i>		Brown <i>et al.</i> 2008
<i>Caladenia flava</i> X <i>latifolia</i>	× <i>spectabilis</i> Hopper & A.P. Br.	
<i>Caladenia flava</i> X <i>longicauda</i>	× <i>triangularis</i> R.S. Rogers	
<i>Caladenia flava</i> X <i>marginata</i>	× <i>erminae</i> Hopper & A.P. Br.	
<i>Caladenia flava</i> X <i>nana</i>		Brown <i>et al.</i> 2008
<i>Caladenia flava</i> subspecies <i>flava</i> X <i>reptans</i> subspecies <i>reptans</i>		Brown <i>et al.</i> 2008
<i>Caladenia flava</i> subspecies <i>maculata</i> X <i>reptans</i> subspecies <i>impensa</i>		Brown <i>et al.</i> 2008
<i>Caladenia fluvialis</i> X <i>footeana</i>		A. Brown personal observation
<i>Caladenia footeana</i> X <i>hiemalis</i>		Jones 2006
<i>Caladenia footeana</i> X <i>horistes</i>		A. Brown personal observation
<i>Caladenia footeana</i> X <i>incensa</i>		A. Brown personal observation
<i>Caladenia footeana</i> X <i>nobilis</i>		Jones 2006
<i>Caladenia footeana</i> X <i>remota</i>		A. Brown personal observation
<i>Caladenia gardneri</i> X <i>interjacens</i>		A. Brown personal observation
<i>Caladenia gardneri</i> X <i>latifolia</i>	× <i>idiastes</i> Hopper & A.P. Br.	
<i>Caladenia georgei</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia georgei</i> X <i>speciosa</i>		Jones 2006
<i>Caladenia georgei</i> X <i>swartziorum</i>		Backhouse 2011
<i>Caladenia harringtoniae</i> X <i>longicauda</i>		Backhouse 2011
<i>Caladenia heberleana</i> X <i>longicauda</i>		Backhouse 2011
<i>Caladenia heberleana</i> X <i>pectinata</i>		A. Brown personal observation
<i>Caladenia hiemalis</i> X <i>polychroma</i>		Jones 2006
<i>Caladenia hiemalis</i> X <i>xantha</i>		Jones 2006
<i>Caladenia hirta</i> X <i>longicauda</i>	× <i>suffusa</i> Hopper & A.P. Br.	
<i>Caladenia hirta</i> X <i>pendens</i>		A. Brown personal observation
<i>Caladenia hirta</i> X <i>remota</i>		Brown <i>et al.</i> 2013
<i>Caladenia hirta</i> X <i>saxicola</i>		A. Brown personal observation
<i>Caladenia hoffmanii</i> X <i>longicauda</i>		Brown <i>et al.</i> 2013
<i>Caladenia hopperiana</i> X <i>longicauda</i>		Brown & Brockman 2015



Western Australian Hybrids	Name and Authority	Notes
<i>Caladenia horistes</i> X <i>multiclavia</i>		incorrectly named as <i>xresupina</i> , the type of which is <i>polychroma</i> X <i>multiclavia</i>
<i>Caladenia horistes</i> X <i>pachychila</i>		Jones 2006
<i>Caladenia huegelii</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia huegelii</i> X <i>paludosa</i>		A. Brown personal observation
<i>Caladenia huegelii</i> X <i>speciosa</i>		Jones 2006
<i>Caladenia huegelii</i> X <i>swartsiorum</i>		Brown & Brockman 2015
<i>Caladenia incensa</i> X <i>incrassata</i>		A. Brown personal observation
<i>Caladenia incensa</i> X <i>pachychila</i>		Brown <i>et al.</i> 2013
<i>Caladenia incensa</i> X <i>radialis</i>		A. Brown personal observation
<i>Caladenia incensa</i> X <i>roei</i>		Brown <i>et al.</i> 2008
<i>Caladenia incensa</i> X <i>vulgata</i>		Jones 2006
<i>Caladenia incrassata</i> X <i>pluvialis</i>		A. Brown personal observation
<i>Caladenia infundibularis</i> X <i>lodgeana</i>		A. Brown personal observation
<i>Caladenia integra</i> X <i>longicauda</i>		A. Brown personal observation
<i>Caladenia latifolia</i> X <i>longicauda</i>		Brown <i>et al.</i> 2013
<i>Caladenia latifolia</i> X <i>nana</i>		Jones 2006
<i>Caladenia leucochila</i> X <i>longicauda</i>		Brown & Brockman 2015
<i>Caladenia lobata</i> X <i>longicauda</i>	<i>xhypata</i> Hopper & A.P. Br.	
<i>Caladenia lobata</i> X <i>macrostylis</i>		A. Brown personal observation
<i>Caladenia lodgeana</i> X <i>serotina</i>		A. Brown personal observation
<i>Caladenia longicauda</i> X <i>lorea</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>macrostylis</i>		Jones 2006
<i>Caladenia longicauda</i> X <i>magniclavata</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>nobilis</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>pectinata</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>splendens</i>		Hopper & Brown 2001
<i>Caladenia longicauda</i> X <i>stareorum</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>thinicola</i>		Backhouse 2011
<i>Caladenia longicauda</i> X <i>uliginosa</i>	<i>xexserta</i> Hopper & A.P. Br.	
<i>Caladenia longicauda</i> X <i>winfieldii</i>		Backhouse 2011
<i>Caladenia luteola</i> X <i>pendens</i>		A. Brown personal observation
<i>Caladenia luteola</i> X <i>polychroma</i>		A. Brown personal observation
<i>Caladenia macrotylis</i> X <i>pectinata</i>		Jones 2006
<i>Caladenia marginata</i> X <i>nana</i>		Jones 2006
<i>Caladenia marginata</i> X <i>reptans</i>		Jones 2006
<i>Caladenia melanema</i> X <i>vulgata</i>		A. Brown personal observation
<i>Caladenia microchila</i> X <i>sigmoidea</i>	<i>xtryphera</i> Hopper & A.P. Br.	
<i>Caladenia multiclavia</i> X <i>polychroma</i>	<i>xresupina</i> Hopper & A.P. Br.	
<i>Caladenia multiclavia</i> X <i>pulchra</i>		Brown <i>et al.</i> 2008
<i>Caladenia nana</i> X <i>reptans</i>		Jones 2006
<i>Caladenia nobilis</i> X <i>occidentalis</i>		Jones 2006
<i>Caladenia nobilis</i> X <i>vulgata</i>		Hopper & Brown 2001
<i>Caladenia occidentalis</i> X <i>pendens</i>		A. Brown personal observation
<i>Caladenia pachychila</i> X <i>paradoxa</i>		A. Brown personal observation
<i>Caladenia pachychila</i> X <i>vulgata</i>		Brown <i>et al.</i> 2008
<i>Caladenia paludosa</i> X <i>speciosa</i>		Jones 2006
<i>Caladenia paradoxa</i> X <i>pulchra</i>		Jones 2006
<i>Caladenia paradoxa</i> X <i>radialis</i>		A. Brown personal observation
<i>Caladenia paradoxa</i> X <i>roei</i>		A. Brown personal observation
<i>Caladenia pendens</i> X <i>postea</i>		A. Brown personal observation
<i>Caladenia pholcoidea</i> X <i>serotina</i>		Jones 2006
<i>Caladenia pluvialis</i> X <i>radialis</i>		Backhouse 2011
<i>Caladenia pluvialis</i> X <i>roei</i>		A. Brown personal observation
<i>Caladenia polychroma</i> X <i>pulchra</i>		Jones 2006
<i>Caladenia polychroma</i> X <i>roei</i>		G. Backhouse personal observation
<i>Caladenia polychroma</i> X <i>xantha</i>		Jones 2006
<i>Caladenia radialis</i> X <i>remota</i>		A. Brown personal observation
<i>Caladenia radialis</i> X <i>roei</i>		A. Brown personal observation

Western Australian Hybrids	Name and Authority	Notes
<i>Caladenia radiata</i> X <i>serotina</i>		Brown <i>et al.</i> 2013
<i>Caladenia remota</i> X <i>roei</i>		Brown <i>et al.</i> 2008
<i>Caladenia roei</i> X <i>voigtii</i>		A. Brown personal observation
<i>Caladenia roei</i> X <i>vulgata</i>		Brown <i>et al.</i> 2013
<i>Caladenia roei</i> X <i>wanosa</i>		A. Brown personal observation
<i>Caladenia vulgata</i> X <i>wanosa</i>		Backhouse 2011
<i>Caladenia dimidia</i> X <i>Pheladenia deformis</i>		Brown <i>et al.</i> 2013
<i>Caladenia hirta</i> X <i>Pheladenia deformis</i>		A. Brown personal observation
<i>Caladenia radialis</i> X <i>Pheladenia deformis</i>		Jones 2006
<i>Cyanicula gemmata</i> X <i>sericea</i>		A. Brown personal observation
<i>Cyanicula amplexans</i> X <i>Pheladenia deformis</i>		Brown <i>et al.</i> 2013
<i>Cyanicula sericea</i> X <i>Elythranthera brunonis</i>	× <i>Cyanthera glossodioides</i> Hopper & A.P. Br.	
<i>Diuris corymbosa</i> X <i>laxiflora</i>		Brown <i>et al.</i> 2013
<i>Diuris laxiflora</i> X <i>purdei</i>		Brown <i>et al.</i> 2013
<i>Diuris septentrionalis</i> X <i>tinkeri</i>		A. Brown personal observation
<i>Diuris septentrionalis</i> X species 'Zuytdorp cliffs'		
<i>Diuris</i> species 'sandplains' X species 'Yalgorup'		Brown <i>et al.</i> 2013
<i>Drakaea glyptodon</i> X <i>livida</i>		
<i>Drakaea gracilis</i> X <i>livida</i>		Jones 2006
<i>Drakaea livida</i> X <i>thynniphila</i>		A. Brown personal observation
<i>Elythranthera brunonis</i> X <i>emarginata</i>	× <i>intermedia</i> (Fitzg.) M.A. Clem.	
<i>Ericksonella saccharata</i> X <i>Pheladenia deformis</i>		Brown <i>et al.</i> 2013
<i>Microtis alba</i> X <i>media</i>		A. Brown personal observation
<i>Prasophyllum brownii</i> X <i>regium</i>		Brown <i>et al.</i> 2013
<i>Prasophyllum fimbria</i> X <i>giganteum</i>		Brown <i>et al.</i> 2013
<i>Pterostylis allantoidea</i> X species 'inland'		A. Brown personal observation
<i>Thelymitra antennifera</i> X <i>graminea</i>		Brown <i>et al.</i> 2013
<i>Thelymitra antennifera</i> X <i>macrophylla</i>		Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>maculata</i>		Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>petrophila</i>		Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>vulgaris</i>		Brown <i>et al.</i> 2008
<i>Thelymitra antennifera</i> X <i>vulgaris</i>		Brown <i>et al.</i> 2008
<i>Thelymitra apiculata</i> X <i>pulcherrima</i>		
<i>Thelymitra benthamiana</i> X <i>macrophylla</i>		Brown <i>et al.</i> 2008
<i>Thelymitra benthamiana</i> X <i>villosa</i>		Brown <i>et al.</i> 2013
<i>Thelymitra campanulata</i> X <i>spiralis</i>		
<i>Thelymitra cornicina</i> X <i>crinita</i>		Jones 2006
<i>Thelymitra crinita</i> X <i>macrophylla</i>		Brown <i>et al.</i> 2013
<i>Thelymitra flexuosa</i> X <i>macrophylla</i>		Jones 2006
<i>Thelymitra flexuosa</i> X <i>vulgaris</i>		Brown <i>et al.</i> 2013
<i>Thelymitra latiloba</i> X <i>graminea</i>		Brown <i>et al.</i> 2013 as <i>latiloba</i> x <i>petrophila</i>
<i>Thelymitra macrophylla</i> X <i>vulgaris</i>		Jones 2006
<i>Thelymitra sargentii</i> X species 'Murchison'		A. Brown personal observation
<i>Thelymitra villosa</i> X species 'Murchison'		Brown <i>et al.</i> 2013 as <i>macrophylla</i> x <i>villosa</i>

**Total: 218 hybrids**

Australia's Island Territories Species	Alternate Genus	Notes
<i>Brachypeza archytas</i> (Ridl.) Garay		Christmas Island
<i>Bryobium retusum</i> (Blume) Y.P.Ng & P.J.Cribb	<i>Eria</i>	Christmas Island
<i>Bulbophyllum argyropus</i> (Endl.) Rchb.f.	<i>Adelopetalum</i>	also on Lord Howe Island, Norfolk Island
<i>Corybas acuminatus</i> M.A.Clem. & Hatch		Norfolk Island
<i>Corymborkis veratrifolia</i> (Reinw.) Blume		Christmas Island
<i>Dendrobium brachypus</i> (Endl.) Rchb.f.	<i>Thelychiton</i>	Norfolk Island
<i>Dendrobium crumenatum</i> Sw.	<i>Ceraia</i>	Christmas Island; species name <i>saaronica</i> sometimes used
<i>Dendrobium insigne</i> Blume	<i>Dichopus, Grastidium</i>	Dauan Island, Torres Strait
<i>Dendrobium litorale</i> Schltr.	<i>Aporopsis</i>	Dauan Island, Torres Strait; also called <i>Dendrobium macfarlanei</i>
<i>Dendrobium macropus</i> (Endl.) Rchb.f. ex Lindl.	<i>Thelychiton</i>	Norfolk Island
<i>Dendrobium nativitatis</i> Ridl.	<i>Flickingeria</i>	Christmas Island
<i>Dendrobium reconditum</i> Schuit. & P.B.Adams	<i>Cadetia</i>	Moa Island, Torres Strait; also called <i>Cadetia clausa</i>
<i>Didymoplexis pallens</i> Griffith		Christmas Island
<i>Habenaria vatia</i> D.L.Jones		Moa Island, Torres Strait
<i>Microtis unifolia</i> (G.Forst.) Rchb.f.		also on Lord Howe Island, Norfolk Island
<i>Oberonia titania</i> Lindl.		Norfolk Island
<i>Phreatia paleata</i> (Rchb.f.) Rchb.f.		Norfolk Island
<i>Plexaure limenophylax</i> Endl.		Norfolk Island
<i>Plexaure listeri</i> (Rolfe) M.A.Clem. & D.L.Jones		Christmas Island
<i>Sarcanthopsis warocqueana</i> (Rolfe) Garay		Moa Island, Torres Strait
<i>Taeniophyllum hasseltii</i> Rchb.f.		Christmas Island
<i>Taeniophyllum norfolkianum</i> D.L.Jones, B.Gray & M.A.Clem.		Norfolk Island
<i>Thelasis capitata</i> Blume		Christmas Island
<i>Thelymitra longifolia</i> J.R.Forst. & G.Forst.		Norfolk Island
<i>Thrixspermum carinatifolium</i> (Ridl.) Schltr.		Christmas Island
<i>Tropidia viridifusca</i> Kraenzl.		Norfolk Island
<i>Zeuxine exilis</i> Ridl.		Christmas Island

**Total: 27 species**