

TUESDAY, 13TH MAY, 1884.

THE PRESIDENT HON. A. C. GREGORY, C.M.G., IN THE CHAIR.

The following Donations were announced:—

1. "Report of Progress" for 1880-81-82. Geological and Natural History Survey of Canada. Montreal, 1883. From Alfred R. C. Selwyn, L.L.D., F.R.S., Director, Geological and Natural History Survey of Canada.

2. "Maps to accompany Report of Progress for 1880-81-82. Montreal, 1883. 1d.

3. "Catalogue of Canadian Plants." Part I. Polypetalæ, by John Macoun, M.A., F.L.S., Montreal, 1883. 1d.

4. "Journals of Australian Explorations," by A. C. Gregory, C.M.G., F.R.G.S., &c, and F. T. Gregory, F.R.G.S., etc., Brisbane, 1884. From Hon. A. C. Gregory, C.M.G., &c.

5. "Eucalyptographia," by Baron Ferd. Von Mueller, K.C.M.G., M. and Ph. D., F.R.S., etc. From the Author.

6. "Systematic Census of Australian Plants. First Annual Supplement," by Baron Ferd. Von Mueller, K.C.M.G., etc., Melbourne, 1884. From the author.

The following Papers were read:—

A CONTRIBUTION TOWARDS A FLORA OF MOUNT PERRY

BY

F. M. BAILEY, F.L.S., GOVERNMENT BOTANIST.

Mr. James Keys, of Mt. Perry, has from time to time sent me down the plants of his neighbourhood for determination; and it is by reason of his efforts, in this direction, that the present attempt at a local Flora has been rendered possible.

ORDER RANUNCULACEÆ.

Clematis glycinoides, DC. (Virgin's Bower, or Traveller's Joy.)

ORDER DILLENIACEÆ.

Hibbertia linearis, var. *obtusifolia*,

ORDER ANONACEÆ.

Melodorum Leichhardtii, *Benth.*

ORDER MENISPERMACEÆ.

Stephania hernandiæfolia, *Walp.*

ORDER CRUCIFERÆ.

Cardamine hirsuta, *Linn.*

ORDER CAPPARIDEÆ.

Capparis sarmentosa, *A. Cunn.*

„ *nobilis*, *F.v.M.*

„ *canescens*, *Banks.*

ORDER VIOLARIEÆ.

Viola betonicaefolia, *Sm.* (Violet.)

„ *hederacea*, *Labill.* (Creeping Violet.)

Ionidium suffruticosum, *Ging.*

ORDER PITTOSPOREÆ.

Pittosporum rhombifolium, *A. Cunn.*

„ *revolutum*, *Ait.*

Citriobatus multiflorus, *A. Cunn.*

ORDER POLYGALEÆ.

Polygala japonica, *Houtt.* (Milk wort.)

ORDER PORTULACEÆ.

Portulaca oleracea, *Linn.* (Pigweed.)

ORDER HYPERICINEÆ.

Hypericum gramineum, *Forst.* (St. John's wort.)

ORDER MALVACEÆ.

Malvastrum spicatum, *A. Gray.*

„ *tricuspidatum*, *A. Gray.*

Sida corrugata, *Linn.*

„ *subspicata*, *F.v.M.*

„ *rhombifolia*, *Linn.* (Sida-weed.)

Abutilon oxycarpum, *F.v.M.*

„ *auritum*, *G. Don.*

Hibiscus rhodopetalus, *F.v.M.*

„ *splendens*, *Fraser.!*

ORDER STERCULIACEÆ.

Sterculia quadrifida, *R. Br.*

„ *Bidwilli*, *Hook.*

- Sterculia diversifolia*, *G. Don.* (Currijong.)
 „ *rupestris*, *Benth.* (Narrow-leaved Bottle-tree.)
Melhania incana, *Heyne.*
Waltheria americana, *Linn.*
Commersonia echinata, *Forst.*

ORDER TILIACEÆ.

- Grewia latifolia*, *F.v.M.*
Elæocarpus obovatus, *G. Don.*

ORDER LINEÆ.

- Erythroxylon australe*, *F.v.M.*

ORDER ZYGOPHYLLÆÆ.

- Tribulus terrestris*, *Linn.*

ORDER GERANIACEÆ.

- Geranium dissectum*, *Linn.*
Erodium cygnorum, *Nees.*
Oxalis corniculata, *Linn.* (Small horned Wood-Sorrel.)

ORDER RUTACEÆ.

- Zanthoxylum brachyacanthum*, *F.v.M.*
Geijera Muelleri, *Benth.*
 „ *salicifolia*, *Schott.*
Acronychia Baueri, *Schott.*
 „ *lævis*, *Forst.*
Micromelum pubescens, *Bl.*
Clausena brevistyla, *Oliver.*

A very rare shrub, not before met with so far south, said to bear an intense red fruit, which might probably be utilised as that of its near ally *Atalantia*, for preserves.

ORDER SIMARUBEÆ.

- Ailanthus imberbiflora*, *F.v.M.*

ORDER BURSERACEÆ.

- Canarium australasicum*, *F.v.M.*

ORDER MELIACEÆ.

- Turræa pubescens*, *Hellen.*
Melia composita, *Willd.* (White Cedar.)
Dysoxylon Muelleri, *Benth.*
Synoum glandulosum, *A. Juss.*
Owenia venosa, *F.v.M.*
Cedrela Toona, *Roxb.* (Red Cedar.)
Flindersia Australis, *R. Br.* (Yellow-wood.)

ORDER CELASTRINEÆ.

- Elæodendron australe, *Vent.*
 „ melanocarpum, *F.v.M.*
 Siphonodon australe, *Benth.*

ORDER STACKHOUSIÆ.

- Stackhousia monogyna, *Labill.*

ORDER RHAMNEÆ.

- Alphitonia excelsa, *Reissek.* (Red Ash.)

ORDER AMPELIDEÆ.

- Vitis antarctica, *Benth.*
 „ oblonga, *Benth.*
 „ nitens, *F.v.M.*
 „ acris, *F.v.M.*
 „ clematidea, *F.v.M.*
 „ hypoglauca, *F.v.M.*
 „ opaca, *F.v.M.*

ORDER SAPINDACEÆ.

- Cupania anacardioides, *A. Rich.*
 „ pseudorhus, *A. Rich.*
 „ xylocarpa, *A. Cunn.*
 „ nervosa, *F.v.M.*
 Ratonia pyriformis, *Benth.*
 Nephelium tomentosum, *F.v.M.*
 Harpullia pendula, *Planch.* (Tulip-wood.)
 Dodonæa triquetra, *Andr.* (Hop-bush.)
 „ viscosa, *Linn.*
 „ cuneata, *Rudge.*

ORDER LEGUMINOSÆ SUBOR. PAPILIONACEÆ.

- Jacksonia scoparia, *R. Br.* (Dogwood.)
 Crotalaria alata, *Hamilt.* (Rattle pod.)
 „ linifolia, *Linn. f.*
 „ Mitchelli, *Benth.*
 „ trifolium, *Willd.*
 „ incana, *Linn.*
 Lotus, australis, *Andr.*

This Australian Bird's-foot Trefoil is looked upon in some parts as a poison herb.

- Indigofera enneaphylla, *Linn.*
 „ hirsuta, *Linn.*
 „ pratensis, *F.v.M.*

Indigofera Baileyi, *F.v.M.*

„ *australis*, *Willd.*

Tephrosia filipes, *Benth.*

„ *purpurea*, *Pers.*

In some parts of Australia this has been found poisonous to stock.

Swainsona galegifolia, *R. Br.* (Indigo or Darling Pea.)

„ *brachycarpa*, *Benth.*

Æschynomene falcata, *DC.*

Zornia diphylla, *Pers.*

Desmodium gangeticum, *DC.*

„ *brachypodum*, *A. Gray.*

„ *varians*, *Endl.*

„ *parvifolium*, *DC.*

Uraria picta, *Desv.*

Lespedeza cuneata, *G. Don.*

Glycine clandestina, *Wendl.*

„ *tabacina*, *Benth.*

Hardenbergia monophylla, *Benth.* (Bushmen's Sarsaparilla.)

Erythrina vespertilio, *Benth.* (Coral or Cork-tree.)

Canavalia obtusifolia, *DC.*

Vigna vexillata, *Benth.*

Atylosia reticulata, *Benth.*

Rhynchosia Cunninghamii, *Benth.*

„ *minima*, *DC.*

Lonchocarpus Blackii, *Benth.*

Castanospermum australe, *A. Cunn.* (Moreton Bay Chestnut.)

SUBORDER CÆSALPINIÆ.

Cæsalpinia sepiaria, *Roxb.* (A strong hedge plant, naturalised.)

Cassia Sophera var. *schinifolia*

„ *concinna*, *Benth.*

„ *mimosoides*, *Linn.*

SUBORDER MIMOSÆ.

Neptunia gracilis, *Benth.* (Native Sensitive Plant.)

Acacia penninervis, *Sieb.*

„ *decora*, *Reichb.*

„ *podalyriæfolia*, *A. Cunn.*

„ *implexa*, *Benth.*

„ *aulacocarpa*, *A. Cunn.*

„ *Bidwilli*, *Benth.*

Pithecolobium pruinatum, *Benth.*

ORDER ROSACEÆ.

- Rubus moluccanus*, *Linn.*
 „ *parvifolius*, *Linn.*
 „ *rosæfolius*, *Sm.* (Raspberry.)

ORDER SAXIFRAGEÆ.

- Argophyllum Lejournanii*, *F.v.M.*
Abrophyllum ornans, *Hook, F.*

ORDER CRASSULACEÆ

- Tillæa verticillaris*, *DC.*

ORDER DROSERACEÆ.

- Drosera Burmanni*, *Vahl.* (Sundew.)
 „ *auriculata*, *Backh.* (Sundew.)

ORDER HALORAGEÆ.

- Haloragis heterophylla*, *Brongn.*
Myriophyllum verrucosum, *Labill.*
 „ *variæfolium*, *Hook.*

ORDER MYRTACEÆ.

- Callistemon lanceolatus*, *DC.* (Bottle-brush.)
Melaleuca linariifolia, *Sm.*
Angophora subvelutina, *F.v.M.* (Apple tree.)
Eucalyptus acmenioides, *Schau.* (Stringy-bark.)
 „ *melanophloia*, *F.v.M.* (Silver-leaved Ironbark.)
 „ *crebra*, *F.v.M.* (Ironbark.)
 „ *tereticornis*, *Sm.* (Blue Gum)
 „ *tesselaris*, *F.v.M.* (Moreton Bay Ash.)
 „ *corymbosa*, *Sm.* (Bloodwood)
 „ *maculata*, *Hook.* (Spotted Gum.)
 „ *var. citriodora*. (Citron scented Gum.)
Tristania suaveolens, *Sm.*
 „ *conferta*, *R. Br.* (Brisbane Box.)
Myrtus Hillii, *Benth.* (Scrub Iron-wood.)
 „ *racemulosa*, *Benth.*
Eugenia Smithii, *Poir.* (Lilly Pilly.)
 „ *Ventenatii*, *Benth.*
 „ *myrtifolia*, *Sims.* (Scrub Cherry.)

ORDER MELASTOMACEÆ.

- Melastoma malabathricum*, *Linn.*

ORDER LYTHRARIÆ.

- Ammannia pentandra*, *Roxb.*

ORDER ONAGRARIÆ.

Jussiaea suffruticosa, *Linn.*

ORDER CUCURBITACEÆ.

Melothria Cunninghamii, *F.v.M.*

Mukia scabrella, *Arn.*

ORDER CACTEÆ.

Opuntia vulgaris, *Mill.* (Prickly Pear, naturalized.)

ORDER UMBELLIFERÆ.

Hydrocotyle hirta, *R. Br.*

„ *asiatica*, *Linn.*

In the Mauritius, where this plant is called “Bevilaqua,” it is highly extolled as a remedy in diseases of the skin.

Trachymene procumbens, *Benth.*

Eryngium expansum, *F.v.M.*

Apium leptophyllum, *F.v.M.*

ORDER ARALIACEÆ.

Panax elegans, *F.v.M.*

ORDER CORNACEÆ.

Marlea vitiensis, var. *tomentosa*.

ORDER CAPRIFOLIACEÆ.

Sambucus Gaudichaudiana, *DC.*

ORDER RUBIACEÆ.

Dentella repens, *Forst.*

Knoxia corymbosa, *Willd.*

Hodgkinsonia ovatiflora, *F.v.M.*

Plectronia odorata, *F.v.M.*

Ixora Becklerii, *Benth.*

Pavetta indica, *Linn.*

Psychotria loniceroides, *Sieb.*

„ *daphnoides*, *A. Cunn.*

Pomax unbellata, *Sol.*

Spermacoce multicaulis, *Benth.*

ORDER COMPOSITÆ.

Centratherum muticum, *Less.*

Ageratum conyzoides, *Linn.*

Lagenophora Billardieri, *Cass.*

Calotis dentex, *R. Br.*

„ *lappulacea*, *Benth.*

- Olearia Nernstii*, *F.v.M.*
Erigeron linifolius, *Linn.* (Cobbler's-pegs.)
Vittadinia brachycomoides, *F.v.M.*
 ,, *australis*, *A. Rich.*
 ,, *scabra*, *DC.*
Blumea hieracifolia, *DC.*
Pterocaulon sphacelatum, *Benth et Hook.*
Helipterum polyphyllum, *F.v.M.*
Helichrysum bracteatum, *Willd.*
 ,, *apiculatum*, *DC.*
Cassinia lævis, *R. Br.*
Podolepis acuminata, *R. Br.*
Xanthium spinosum, *Linn.* (Bathurst Burr.)
Siegesbeckia orientalis, *Linn.*
Eclipta platyglossa, *F.v.M.*
Spilanthes grandiflora, *Turez.*
Tagetes glandulifera, *Schränk.* (Stinking Rodger.)
Erechthites Ackinsoniæ, *F.v.M.*
Senecio amygdalifolius, *F.v.M.*
Picris hieracioides, *Linn.*

ORDER STYLIDIEÆ.

- Stylidium graminifolium*, *Sw.*

ORDER GOODENOVIÆ.

- Goodenia heterophylla*, *Sm.*
 ,, *glabra*, *R. Br.*
 ,, *grandiflora*, *Sims.*
Velleia spathulata, *R. Br.*

ORDER CAMPANULACEÆ.

- Lobelia gibbosa*, *Labill.*
 ,, *purpurascens*, *R. Br.*
Wahlenbergia gracilis, *A. DC.* (Blue-bell.)

ORDER EPACRIDEÆ.

- Acrotriche aggregata*, *R. Br.*

ORDER PLUMBAGINEÆ.

- Plumbago zeylanica*, *Linn.* (Leadwort.)

ORDER PRIMULACEÆ.

- Anagallis arvensis*, *Linn.* (Pimpernel, naturalized.)

ORDER MYRSINEÆ.

- Myrsine variabilis*, *R. Br.*
Samara australiana, *F.v.M.*

ORDER SAPOTACEÆ.

Lucuma (Sersalisia) *sericea*, *R. Br.*

Sideroxylon (*Achras*) *myrsinoides*, *A. Cunn.*

ORDER EBENACEÆ.

Maba *geminata*, *R. Br.*

„ *humilis*, *R. Br.*

ORDER OLEACEÆ.

Jasminum *racemosum*, *F.v.M.*

„ *lineare*, *R. Br.*

„ *simplicifolium*, *Forst.*

Notelæa *microcarpa*, *R. Br.*

Olea *paniculata*, *R. Br.* (Australian Olive.)

ORDER APOCYNACEÆ.

Chilocarpus *australis*, *F.v.M.*

Carissa *ovata*, *R. Br.*

Alyxia *ruscifolia*, *R. Br.*

Tabernæmontana *orientalis*, var. *angustifolia*.

Lyonsia *lilacina*, *F.v.M.*

„ *reticulata*, *F.v.M.*

„ *latifolia*, *Benth.*

„ *eucalyptifolia*, *F.v.M.*

„ *straminea*, *R. Br.*

Parsonsia, *ventricosa*, *F.v.M.*

ORDER ASCLEPIADEÆ.

Secamone *elliptica*, *R. Br.*

Asclepias *curassavica*, *Linn.*

(Red-head or Wild Ipecacuanha, naturalized.)

Sarcostemma *australe*, *R. Br.* (Gaoloowurrah.)

Gongronema *micradenia*, *Benth.*

Tylophora *grandiflora*, *R. Br.*

Marsdenia *rostrata*, *R. Br.*

Hoya *australis*, *R. Br.* (Honey Flower.)

ORDER LOGANIACEÆ.

Mitrasacme *indica*, *Wight.*

„ *pygmæa*, *R. Br.*

ORDER GENTIANEÆ.

Erythræa *australis*, *R. Br.* (Centaury.)

Limnanthemum *crenatum*, *F.v.M.* (Yellow fringed Water Lily.)

ORDER BORAGINEÆ.

Trichodesma *zeylanicum*, *R. Br.*

ORDER CONVULVULACEÆ.

- Ipomæa hederacea*, *Jacq.*
„ *plebeia*, *R. Br.*
Evolvulus alsinoides, *Linn.*

ORDER SOLANACEÆ.

- Lycopersicum esculentum*, *Mill.* (Tomato naturalized.)
Solanum nigrum, *Linn.* (Common black Nightshade.)
„ *aviculare*, *Først.*
„ *verbascifolium*, *Ait.*
„ *discolor*, *R. Br.*
„ *stelligerum*, *San.*
„ *ellipticum*, *R. Br.*
Datura fastuosa, *Mill.* (Proud Thorn-apple, naturalized.)
Nicotiana suaveolens, *Lehm.* (Sweet Tobacco.)

ORDER SCROPHULARINEÆ.

- Morgania glabra*, *R. Br.*
Gratiola pedunculata, *R. Br.*
Artanema fimbriatum, *Don.*
Veronica calycina, *R. Br.*
Buchnera ramosissima, *R. Br.*
Centranthera hispida, *R. Br.*

ORDER LENTIBULARIEÆ.

- Utricularia dichotoma*, *Labill.* (Bladder-wort.)

ORDER BIGNONIACEÆ.

- Tecoma australis*, *R. Br.*

ORDER ACANTHACEÆ.

- Hygrophila salicifolia*, *Nees.*
Ruellia bracteata, *R. Br.*
Eranthemum variabile, *R. Br.*
„ *tenellum*, *Benth.*
Justicia procumbens, *Linn.*

ORDER MYOPORINEÆ.

- Myoporum debile*, *R. Br.*

ORDER VERBENACEÆ.

- Pityrodia salvifolia*, *R. Br.*
Spartothamnus junceus, *A. Cunn.*
Verbena officinalis, *Linn.*
Callicarpa pedunculata, *R. Br.*
Premna Dallachyana, *Benth.*

Vitex lignum-vitæ, *A. Cunn.*

The fruit of this tree is largely devoured by the fruit-eating pigeon (*Megaloprepia magnifica*).

Clerodendron tomentosum, *R. Br.*

„ *floribunda*, *R. Br.* (Supposed poisonous to stock.)

ORDER LABIATÆ.

Plectranthus parviflorus, *Willd.*

Mentha satureioides, *R. Br.* (Pennyroyal.)

Salvia plebeia, *R. Br.*

Anisomeles salvifolia, *R. Br.*

Stachys arvensis, *Linn.* (Hedge-nettle, naturalized.)

Teucrium argutum, *R. Br.*

Ajuga australis, *R. Br.* (Bugle.)

ORDER PLANTAGINEÆ.

Plantago debilis, *R. Br.* (Plantain.)

ORDER NYCTAGINEÆ.

Boerhaavia diffusa, *Linn.*

ORDER AMARANTACEÆ.

Deeringia celosioides, *R. Br.*

„ *altissima*, *F.v. M.*

Amarantus viridis, *Linn.*

Achyranthes aspera, *Linn.*

Nyssanthes diffusa, *R. Br.*

Alternanthera nana, *R. Br.*

ORDER CHENOPODIACEÆ.

Chenopodium ambrosioides, *Linn.*

„ *carinatum*, *R. Br.*

ORDER POLYGONACEÆ.

Polygonum minus, *Huds.*

Rumex Brownii, *Camp.* (Dock.)

ORDER ARISTOLOCHIACEÆ.

Aristolochia pubera, *R. Br.* (Birthwort.)

ORDER PIPERACEÆ.

Piper novæ-hollandiæ, *Miq.*

This Pepper bears a large quantity of fruit scarcely inferior to the black pepper of commerce.

Peperomia leptostachya, *Hook et Arn.*

ORDER MONIMIACEÆ.

Kibara macrophylla, *Benth.*

ORDER LAURINEÆ.

Cassytha filiformis, *Linn.* (Laurel Dodder.)

ORDER PROTEACEÆ.

Banksia integrifolia, *Linn. f.* (Honeysuckle tree.)

ORDER LORANTHACEÆ.

Loranthus longiflorus, *Desv.*

„ *exocarp*i, *Behr.*

„ *pendulus*, *Sieb.*

Viscum articulatum, *Burm.* (Mistletoe.)

ORDER SANTALACEÆ.

Exocarpus latifolia, *R. Br.*

„ *cupressiformis*, *Labill.*

ORDER EUPHORBIACEÆ.

Euphorbia Drummondii, *Boiss.* (Sheep poison-herb.)

„ *Macgillivrayi*, *Boiss.*

„ *pilulifera*, *Linn.* (Asthma herb.)

Cleistanthus Cunninghamii, *Muell. Arg.*

Phyllanthus Ferdinandi, *Muell. Arg.*

„ *lobocarpus*, *Benth.*

„ *thesioides*, *Benth.*

„ *albiflorus*, *F.v.M.*

„ *Gasstroemii*, *Muell. Arg.*

„ *minutiflorus*, *F.v.M.*

Breynia oblongifolia, *Muell. Arg.*

Petalostigma quadriloculare, *F.v.M.* (Emu apple or bitter-bark.)

Croton insularis, *Baill.*

„ *acronychioides*, *F.v.M.*

Baloghia lucida, *Endl.* (Scrub Bloodwood.)

Adriana acerifolia, *Hook.*

Acalypha capillipes, *F.v.M.*

Alchornea ilicifolia, *Muell. Arg.* (Queensland Holly-bush.)

Mallotus philippinensis, *Muell. Arg.* ("Kamala" Tree.)

Ricinus communis, *Linn.* (Castor Oil Plant, naturalized.)

Tragia Novæ Hollandiæ, *Muell. Arg.* (Climbing Nettle.)

Homalanthus stillingiæfolius, *F.v.M.*

Excæcaria Dallachyana, *Baill.*

ORDER URTICACEÆ.

Trema aspera, *Blume*.

(Called sometimes Peach-leaved Poison Bush.)

Aphananthe philippinensis, *Planch*.

Malaisia tortuosa, *Blanco*.

Ficus Cunninghamii, *Miq*.

„ *macrophylla*, *Desf*. (Moreton Bay Fig.)

„ *aspera*, *Forst*. (Black Fig.)

„ *opposita*, *Miq*.

„ *glomerata*, *Willd*. (Cluster Fig.)

Cudrania javanensis, *Trécul*.

Laportea photiniphylla, *Wedd*. (Shiny-leaved Stinging Tree.)

Pipturus argenteus, *Wedd*.

ORDER CASUARINEÆ.

Casuarina glauca, *Sieb*. (River She-oak.)

„ *torulosa*, *Ait*. (Corky-barked She-oak.)

ORDER CONIFERÆ.

Araucaria Cunninghamii, *Ait*. (Moreton Bay or Hoop Pine.)

ORDER HYDROCHARIDEÆ.

Ottelia ovalifolia, *L. C. Rich*.

ORDER ORCHIDEÆ.

Dendrobium tetragonum, *A. Cunn*.

„ *linguiforme*, *Sw*.

„ *Mortii*, *F.v.M*.

Sarcochilus divitiflorus, *F.v.M*.

„ *Hillii*, *F.v.M*.

Pterostylis ophioglossa, *R. Br*.

„ *acuminata*, *R. Br*.

Glossodia minor, *R. Br*.

ORDER SCITAMINEÆ.

Alpinia cærulea, *Benth*. (Native Ginger.)

ORDER AMARYLLIDEÆ.

Curculigo ensifolia, *R. Br*.

Hypoxis hygrometrica, *Labill*.

ORDER DIOSCORIDEÆ.

Dioscorea transversa, *R. Br*. (Yam.)

ORDER LILIACEÆ.

Rhipogonum album, *R. Br*.

Dianella lævis, *R. Br*.

Eustrephus latifolius var. *angustifolia*
Cordyline terminalis, *Kunth.* (Lily Palm.)
Iphigenia indica, *Kunth.*
Thysanotus tuberosus *R. Br.* (Fringe Violet.)
Cæsia vittata *R. Br.*
Tricoryne elatior, *h. Br.*
Laxmannia gracilis, *R. Br.*

ORDER PHILYDRACEÆ.

Philydrum lanuginosum, *Banks.*

ORDER XYRIDEÆ.

Xyris complanata, *R. Br.*

ORDER COMMELYNACEÆ.

Commelina cyanea, *R. Br.* (Blue Spiderwort)
Aneilema gramineum, *R. Br.*
Polia macrophylla, *Benth.*

ORDER FLAGELLARIEÆ.

Flagellaria indica, *Linn.*

ORDER JUNCACEÆ.

Xerotes longifolia, *R. Br.*
 „ *multiflora*, *R. Br.*
Xanthorrhæa arborea, *R. Br.* (Grass-tree.)
Juncus communis, *E. Mey.* (Common rush.)

ORDER PALMÆ.

Archontophoenix Cunninghamii, *Wendl. et Drude.*
Livistona australis, *Mart.*

ORDER AROIDEÆ.

Typhonium Brownii, *Schott.*
Gymnostachys anceps, *R. Br.*

ORDER NAIADEÆ.

Triglochin procera, *R. Br.*

ORDER ERIOCAULEÆ.

Eriocaulon Smithii, *R. Br.*

ORDER CYPERACEÆ.

Kyllinga cylindrica, *Nees.*
Cyperus polystachyus *Rottb.*
 „ *difformis*, *Linn.*

Cyperus fulvus, *R. Br.*
 „ *exaltatus*, *Retz.*
Heleocharis cylindrostachys, *Bæcket.*
Fuirena glomerata, *Lam.*
Gahnia aspera, *Spreug.*
Scleria Brownii, *Kunth.*

ORDER GRAMINEÆ.

Paspalum scrobiculatum, *Linn.*
Panicum sanguinale, *Linn.*
 „ *parviflorum*, *R. Br.*
 „ *leucophœum*, *H. B. et K.*
 „ *semialatum*, *R. Br.*

In tropical Queensland, Dr. Lumholtz found that the white cockatoos rooted up this grass to feed on its thickened root-stock.

Panicum effusum, *R. Br.*
 „ *crus-galli*, *Linn.*
Oplismenus compositus, *Beauv.*
Setaria macrostachya, *H. B. et K.*
Arundinella nepalensis, *Trin.*
Imperata arundinacea, *Cyrs.* (Blady grass.)
Heteropogon contortus, *Ræm. et Schult.* (Bunch Spear grass)
Andropogon pertusus, *Willd.*
 „ *intermedius*, *R. Br.*
 „ *refractus*, *R. Br.*
 „ *lachnatherus*, *Benth.*
Chrysopogon parviflorus, *Benth.*
Anthistiria ciliata, *Linn.* (Kangaroo Grass.)
Aristida ramosa, *R. Br.*
 „ *calycina*, *R. Br.*
Sporobolus indicus, *R. Br.*
 „ *diander*, *Beauv.*
Cynodon dactylon, *Pers.* (Common Couch-grass.)
Chloris divaricata, *R. Br.* (Star-grass.)
Eleusine indica, *Gærtn.* (Crab-grass.)
Eragrostis Brownii, *Nees.* (Common Love-grass.)

ORDER LYCOPODIACEÆ.

Azolla rubra, *R. Br.* (Red Water-moss.)

ORDER MARSILEACEÆ.

Marsilea hirsuta, *R. Br.* (Water Clover.)

ORDER FILICES.

- Ophioglossum vulgatum*, *Linn.* (Adder's-tongue Fern.)
Gleichenia flabellata, *R. Br.* (Fan Fern.)
Todea barbara, *T. Moore.*
Alsophila australis, *R. Br.* (Common Tree Fern.)
Davallia dubia, *R. Br.* (Mountain Bracken.)
Adiantum æthiopicum, *Linn.* (Small Maiden-hair Fern.)
 ,, *hispidulum*, *Sw.* (Rough-stalked Maiden-hair Fern.)
 ,, *formosum*, *R. Br.*
Cheilanthes tenuifolia var. *Sieberi.* (Curly Fern.)
Pteris falcata, var. *nana.* (Ear-fern.)
 ,, ,, var. *paradoxa.*
 ,, *longifolia*, *Linn.*
 ,, *incisa*, *Thunb.* (Bats-wing Fern.)
Blechnum serrulatum, *Rich.*
 ,, *cartilagineum*, *Sw.*
Doodia aspera *R. Br.* (Prickly Fern.)
 ,, ,, var. *caudata.*
Asplenium nidus, *Linn.* (Bird's-nest Fern.)
Aspidium molle, *Sw.*
 ,, *decompositum*, *Spreng.*
Polypodium tenellum, *Forst.*
 ,, *serpens*, *Forst.*
 ,, *confluens*, *R. Br.*
 ,, *irioides*, *Poir.*
Platycterium grande, *J. Sm.* (Stag's-horn Fern.)
-



Bailey, Frederick Manson. 1884. "A Contribution Towards a Flora of Mount Perry." *The Proceedings of the Royal Society of Queensland* 1(2), 61–76.

<https://doi.org/10.5962/p.351004>.

View This Item Online: <https://www.biodiversitylibrary.org/item/138530>

DOI: <https://doi.org/10.5962/p.351004>

Permalink: <https://www.biodiversitylibrary.org/partpdf/351004>

Holding Institution

American Museum of Natural History Library

Sponsored by

BHL-SIL-FEDLINK

Copyright & Reuse

Copyright Status: Public domain. The BHL considers that this work is no longer under copyright protection.

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.