

Ecological and Phytochemical Significance of *Croton bonplandianum* (Bail)

Abstract:

The aim of current research work is to explore the ecological role of *Croton bonplandianum* specie prevalent to north eastern Punjab region of Pakistan. Under extreme drought condition the plant species survives modestly. The plant on blooming attract economic pollinator insects like honey bees even in dry climate periods. *Croton bonplandianum* (Bail) was collected from wild and re-cultivated under normal atmospheric conditions. Its flowering body parts and stems were plucked and kept under shade for about two weeks for drying. The flowering inflorescence and herbaceous green portions were subjected to maceration, then grinded to fine powder. The powdered samples were analyzed using FTIR spectroscopic technique. FTIR spectral lines have shown different characteristic peaks pattern in case of flowers and non-flowering portion of plant inflorescence segment. Appropriate correlations of peaks to functional groups present were debated.

Key Words: Ecological study, Phytochemicals, *Croton bonplandianum* (bail), FTIR analysis.

Introduction:

Croton is a diverse and complex taxonomic group of plants ranging from herbs and shrubs to trees, though native to south America, but now have global distribution [1]. *Croton* species are known as traditional medicinal plants in Africa, Asia and South America [2-6]. The *Croton* species are rich in alkaloids, flavonoids and terpenoids [7-13]. So due to presence of bioactive compounds, the *croton* species are commonly used to treat cancer, hypertension, inflammation, rheumatism, bleeding gums, malaria, asthma, syphilitic ulcers, diabetes, pain and ulcers [14-21]. *Croton. gratissimus* Burchell, *C. sylvaticus* Hoechst and *C. pseudopulchellus* Pax are well known *croton* species [22]. Surprisingly, sufficient information of *Croton bonplandianum* (Bail) is not available in literature.

Croton bonplandianum (Bail) an exotic weed is one of the important member of genus *Croton*. The plant is usually 1-2 ft. in height, it grows in sandy clay but unpolluted soil along road side. As mentioned earlier that the specie has been is native to Southern Bolivia, Paraguay, South Western Brazil, North Argentina, Bangladesh, South America, India and Pakistan. In Indo-Pak it is widely distributed in the Sub-Himalayan region [23]. In sub-Himalayan region it is often called Ban Tulsi (wild tulsi). *Croton bonplandianum* (bail) is shown in fig 1.



Fig. 1. *Croton bonplandianum* (bail)

Ecological Importance:

Under hot climatic environment the specie *Croton bonplandianum* serve as the source of nectar and food. Economically useful insects like honey bees frequently sit the flowers (fig 2) of *Croton* species, the stalked inflorescence attract the pollinator insects.






Fig. 2. Honey Bees looming *C. bonplandianum* flowers.

The plant species grow-well on arid land because of woody scaly nature of stem and deep penetrating roots, even a very small plant would have a long root penetrated deeper into the soil. *C. bonplandianum* growing on arid land as shown in fig 3.



Fig.3. Species of genus Croton in hot dry environment.

Phytochemical Importance:

Croton bonplandianum (ban tulsi), commonly called wild tulsi, belong to genus Croton which is famous among plant science community as rich source of medicinal compounds [24] So Croton b. will also be a good source of bioactive compounds. It was verified by successful correlation of FTIR spectral lines with expected bioactive compounds.

Biological Activities:

Croton bonplandianum possess immense medicinal value and its stem latex is used by different region as a medicinal plant for the treatment of fresh skin cuts and minor bleeding wounds [25-26]. The plant acquires important biological activities like antioxidant, anti-inflammatory and anticancer [27-28] These activities are attributed to the presence of total ester terpenoids-flavonoids contents [29].

Moreover the extracts of plant have known bio-medical applications in diseases like respiratory, gastro intestinal and skin etc.[30].

Instrumental Investigation:

For investigating bioactive compounds in Croton bonplandianum (bail), FTIR spectroscopic technique was used. FTIR is simple, cost effective and user friendly tool that provides basic information of sample. It is becoming popular technique, especially in testing of biological specimens due to its nondestructive analyzing ability. FTIR provides information as spectral lines only which are related to functional groups of chemical compounds. So for confirmation, other analyzing techniques are needed.

Finger print region ($600\text{-}1450\text{ cm}^{-1}$) is most important for interviewing biological specimens. Both frequency regions (lower and higher) are important to determine functional groups and molecular structure of specimens.

Since long ago researchers have started investigating phytochemicals in medicinal plants by using FTIR [31-34]. Current work is a simple effort to assess presence of phytochemicals in Croton specie by taking Infrared spectra on Varian 640-IR using KBR palettes techniques.

Material and Methods:

Based on leaf morphology and flowering body structure (inflorescence) the plant is recognized as *Croton bonplandianum* (ban tulsī). The plant is collected from sub-urban region of city Rawalpindi where it is grown wild. To maintain its conservation, few plants were collected from wild and re-grown in herbs garden for phytochemical studies. Flowering bodies were harvested in the month of August; the elongated flowering body or heads (inflorescence) were separated from main plant as shown in fig 4-a. Flowers and buds were separated (fig 4-b) and set for drying under shade to avoid photochemical changes. After 20 days, dried samples (buds and

flowers) were grinded to powder. The powders were saved in plastic vials and were subjected to FTIR spectroscopic analysis.




Fig. 4. Fresh twigs (a), Fresh flowers and buds (b) of *Croton bonplandianum*

Results and discussion:

The FTIR spectroscopic result of flowers of *Croton bonplandianum* is shown in fig 5.




Fig. 5. FTIR spectrum of flowers of *C. bonplandianum*.

Table 1. On the basis of FTIR spectra of flowers, expected phytochemical class of compounds are tabulated as under.

Sr. #	Absorption wave #	Group Assigned	Expected phytochemical class
1	3433 (s)	R-OH, R ₂ -NH	Amino acids, Alkaloids, Alcohols etc.
2	3322 (s)	R-OH, R ₂ -NH	Amino acids, Alkaloids, Alcohols etc.
3	3256 (w)	R ₂ -NH	Cyclic amines
4	1667 (w)	-C=C-, CO-R	Aromatic, unsaturated and conjugated compounds , Alkenyl -C=O stretch
5	1581 (s)	-C=C-	Aromatic compounds, nitro oxy comp
6	1469 (s)	-C-H ₃	CH sym and asym bending
7	1156 (w)	R-COOCH ₃	Esters
8	800 (s)	-CH-	Saturated Hydrocarbon, exocyclic double bond stretches
9	715 (w)	-CH-	Five or Six membered Rings, Methylene -(CH ₂) _n - rocking, thiol comp.
10	630 and 660 (s)	organohalogen compound gp	Poly sulfur, thiol compounds

The FTIR ν_{\max} values between 3433-3256 cm^{-1} exhibited the presence of alcohols, phenols and amino acids [35]. Absorption values between 1620-1680 cm^{-1} are of -C=C- stretching related to unsaturated aromatic, carbonyls, terpenoids and α , β unsaturated compounds [36].

Absorption between 1400-1600 cm^{-1} are -C=C- stretching of multiple bond containing compounds like terpenoids [37]. Values near about 1100 cm^{-1} are C-O stretching of alcohols. Absorption bands at around 800 cm^{-1} that were attributed to the presence of compounds with exocyclic double bonds. [38-39] The FTIR absorption peaks between 600-700 cm^{-1} indicate aromatic compounds, thiols etc.

The FTIR result of inflorescence part (buds) of *C. bonplandianum* are presented in figures 6.




Fig 6. FTIR spectrum of non-flowering part of flowering twig of *C. bonplandianum*.

Table 2.: On the basis of FTIR spectra of buds (nonflowering) expected phytochemical class of compounds are tabulated as under.

Sr. #	Absorption wave #	Group Assigned	Expected phytochemical class
1	3480 (w)	R2-NH, R-OH	Amino acids, Alcohols
2	3392 (w)	R2-NH, R-OH	Amino acids, Alcohols
3	3298 (w)	R2-NH, R-OH	Amino acids, Alcohols
4	2904 (s)	-CH-	SP ³ Hydrocarbons, fatty acids
5	2823 (s)	-CH-	SP ³ Hydrocarbons, fatty acids
6	1733 (w)	-CO-, esters stretch	Carbonyl and ester containing compounds, lipids
7	1630 (s)	C=C	Conjugated systems
8	1325 (w)	-CH- bending	Hydrocarbons
9	1235 (w)	OH Aromatic ethers stretch	
10	1025 (s)	-CO-	Ether linkages
11	600-650 (condensed)	Alkyne CH bend	Compounds of CH out-of-plane bending (aromatic compounds)

The FTIR ν_{\max} values between 3480-3298 cm^{-1} exhibited the presence of compounds as discussed before in case of FTIR spectra's of flowers. In FTIR spectrum, absorption bands at 2904 and 2823 cm^{-1} attributable to CH stretching of hydrocarbons and fatty acids [40]. Absorption band at 1733 cm^{-1} that was ascribed to a carbonyl group stretch of an α , β unsaturated γ -lactone ring [38]. Absorption peak at 1630 cm^{-1} attributed to compounds containing double bonds and conjugated groups [36]. The FTIR carbonyl group (ester) stretch was observed at 1733 cm^{-1} confirming the presence of a lactone groups[41-42].

Stretching vibrations between 1200-1300 cm^{-1} depicts presence of hydrocarbons, phenols, alcohols and ethers. Values at 1025 cm^{-1} related to ether linkage due to $-\text{CO}$ stretching. Currently, finger print region (condensed region) in between 600 to 650 cm^{-1} is not simple to evaluate due to peaks condensation.

Both spectra confirm the complete drying nature of plant samples free from moisture. Absence of strong absorption band in the region of spectrum between 3400 cm^{-1} 3100 cm^{-1} indicates the absence of hydrogen bonded OH. Most of the peaks in both spectra are common and hence both part of herb is equally important from application point of view. Hydrocarbon peaks are missing in spectrum of flowers. It may be important indication for further research on *C. bonplandianum*. Esters are also indicated in spectrum of buds. To study further for Phorbols esters, important exploration in *C. bonplandianum* may also be possible.

Conclusion:

The plant species belong to genus *Croton* are known to contain important phytochemicals which belong predominantly to family of cyclic terpenoids, chiefly diterpenoid types like Clerodanes and Phorbols. So current work may be useful for future researchers to use this plant in drug development for human consumption for the treatment of various diseases. Clear FTIR spectra of both flowers and buds revealed that samples of fine powder of delicate plant parts can be tested precisely using this technique if they are well shade dried and moisture free. Moreover, *C. bonplandianum* specie endemic to North-East Punjab has potential ecological role as source of food for variety of ecologically economical insects such as honey bees, larvae of Lepidoptera, especially under hot and droughty climatic situations.

References:

- [1] B. W. van Ee, R. Riina, and P. E. Berry, "A revised infrageneric classification and molecular phylogeny of New World Croton (Euphorbiaceae)," *Taxon*, vol. 60, no. 3, pp. 791–823, **2011**.
- [2] K. P. Tiwari, R. N. Choudhary, and G. D. Pandey, "3-methoxy-4, 6-dihydroxymorphinandien-7-one, an alkaloid from Croton bonplandianum," *Phytochemistry*, vol. 20, no. 4, pp. 863–864, **1981**.
- [3] R. R. Upadhyay and E. Hecker, "A new cryptic irritant and cocarcinogen from seeds of Croton sparciflorus," *Phytochemistry*, vol. 15, no. 6, pp. 1070–1072, **1976**.
- [4] Y. Cai, F. J. Evans, M. F. Roberts, J. D. Phillipson, M. H. Zenk, and Y. Y. Gleba, "Polyphenolic compounds from Croton lechleri," *Phytochemistry*, vol. 30, no. 6, pp. 2033–2040, **1991**.
- [5] N. R. Farnsworth, R. N. Blomster, W. M. Messmer, J. C. King, G. J. Persinos, and J. D. Wilkes, "A phytochemical and biological review of the genus Croton.," *Lloydia*, vol. 32, no. 1, p. 1, **1969**.
- [6] A. C. F. Amaral and R. A. Barnes, "A tetrahydropyroberberine alkaloid from Croton hemiargyreus," *Phytochemistry*, vol. 47, no. 7, pp. 1445–1447, **1998**.
- [7] D. J. Milanowski, R. E. K. Winter, M. P. F. Elvin-Lewis, and W. H. Lewis, "Geographic Distribution of Three Alkaloid Chemotypes of Croton lechleri," *J. Nat. Prod.*, vol. 65, no. 6, pp. 814–819, **2002**.
- [8] W.-H. Lin, H.-Z. Fu, J. Li, G. Cheng, and R. A. Barnes, "The Alkaloids from Leaves of Croton hemiargyreus var. gymnodiscus," *J. Chinese Pharm. Sci.*, vol. 12, no. 3, pp. 117–122, **2003**.
- [9] V. T. de Araújo-Júnior *et al.*, "Muscicapines, a new class of guaiane-type sesquiterpene alkaloids from Croton muscicapa," *J. Braz. Chem. Soc.*, vol. 16, no. 3B, pp. 553–557, **2005**.
- [10] F. Bracher, W. J. Eisenreich, J. Mühlbacher, M. Dreyer, and G. Bringmann,

- “Saludimerines A and B, novel-type dimeric alkaloids with stereogenic centers and configurationally semistable biaryl axes,” *J. Org. Chem.*, vol. 69, no. 25, pp. 8602–8608, **2004**.
- [11] C. A. N. Catalán, C. S. de Heluani, C. Kotowicz, T. E. Gedris, and W. Herz, “A linear sesterterpene, two squalene derivatives and two peptide derivatives from *Croton hieronymi*,” *Phytochemistry*, vol. 64, no. 2, pp. 625–629, **2003**.
- [12] V. T. de Araújo-Júnior *et al.*, “Alkaloids and diterpenes from *Croton moritibensis*,” *Pharm. Biol.*, vol. 42, no. 1, pp. 62–67, **2004**.
- [13] S. Block *et al.*, “Diterpenes from the leaves of *Croton zambesicus*,” *Phytochemistry*, vol. 65, no. 8, pp. 1165–1171, **2004**.
- [14] S. Barrion, E. L. Keya, and T. N. Ngwira, “Country-wine making from Eembe fruit (*Berchemia discolor*) of Namibia,” **2001**.
- [15] H. D. Neuwinger, *African traditional medicine: a dictionary of plant use and applications. With supplement: search system for diseases*. Medpharm, **2000**.
- [16] J. M. Watt and M. G. BREYERBRANDWIJK, “The Medicinal and Poisonous Plants of Southern Africa being an Account of their Medicinal Uses, Chemical Composition, Pharmacological Effects and Toxicology in Man and Animal,” *Med. Poisonous Plants South. Africa being an Acc. their Med. Uses, Chem. Compos. Pharmacol. Eff. Toxicol. Man Anim.*, **1932**.
- [17] H. Beentje, J. Adamson, and D. Bhanderi, *Kenya trees, shrubs, and lianas*. National Museums of Kenya, **1994**.
- [18] Y. Frum and A. M. Viljoen, “In vitro 5-lipoxygenase and anti-oxidant activities of South African medicinal plants commonly used topically for skin diseases,” *Skin Pharmacol. Physiol.*, vol. 19, no. 6, pp. 329–335, **2006**.
- [19] E. Schmidt, M. Lotter, and W. McClelland, *Trees and shrubs of Mpumalanga and Kruger national park*. Jacana Media, 2002.
- [20] A. W. Frahm and H. Hager, [*Handbuch der pharmazeutischen Praxis*]; *Hagers Handbuch*

der pharmazeutischen Praxis. Folgebd. 5. Stoffe: L-Z. Springer, **1999**.

- [21] E. A. Prozesky, J. J. M. Meyer, and A. I. Louw, "In vitro antiplasmodial activity and cytotoxicity of ethnobotanically selected South African plants," *J. Ethnopharmacol.*, vol. 76, no. 3, pp. 239–245, **2001**.
- [22] M. K. Langat, *The Phytochemistry of Three African Croton Species*. University of Surrey (United Kingdom), **2009**.
- [23] A. Zahoor, F. Ahmad, M. H. M. Naseer, and N. Irum, "Palynological features of medicinal plants from Salt Ranges of Pakistan," **2019**.
- [24] R. Nath, S. Roy, B. De, and M. D. Choudhury, "Anticancer and antioxidant activity of croton: a review," *Int J Pharm Pharm Sci*, vol. 5, no. 2, pp. 63–70, **2013**.
- [25] N. K. Sriwastava, C. S. Shreedhara, and H. N. A. Ram, "Standardization of Ajmodadi churna, a polyherbal formulation," *Pharmacognosy Res.*, vol. 2, no. 2, p. 98, **2010**.
- [26] S. Qureshi *et al.*, "Negative aspects of the beneficial herbs: An over view," *J Herb Med Toxicol*, vol. 6, pp. 1–14, **2012**.
- [27] M. S. Islam, M. M. Rahman, M. A. Rahman, M. A. Qayum, and M. F. Alam, "In vitro evaluation of *Croton bonplandianum* Baill. as potential antitumor properties using *Agrobacterium tumefaciens*," *J. Agric. Technol.*, vol. 7, no. 3, pp. 711–719, **2011**.
- [28] J. Bhavana, M. K. Kalaivani, and A. Sumathy, "Cytotoxic and pro-apoptotic activities of leaf extract of *Croton bonplandianus* Baill. against lung cancer cell line A549," **2016**.
- [29] M. S. Islam, H. Ara, K. I. Ahmad, and M. M. Uddin, "A review on medicinal uses of different plants of Euphorbiaceae family," *Univers. J Pharm Res*, vol. 4, pp. 47–51, **2019**.
- [30] S. Divya, K. Naveen Krishna, S. Ramachandran, and M. D. Dhanaraju, "Wound healing and in vitro antioxidant activities of *Croton bonplandianum* leaf extract in rats," *Glob. J. Pharmacol.*, vol. 5, no. 3, pp. 159–163, **2011**.
- [31] I. Ahmad and F. Aqil, "In vitro efficacy of bioactive extracts of 15 medicinal plants against ES β L-producing multidrug-resistant enteric bacteria," *Microbiol. Res.*, vol. 162, no. 3, pp. 264–275, **2007**.

- [32] N. Ramamurthy and S. Kannan, "Fourier transform infrared spectroscopic analysis of a plant (*Calotropis gigantea* Linn) from an industrial village, Cuddalore dt, Tamilnadu, India," *Rom. J. Biophys.*, vol. 17, no. 4, pp. 269–276, **2007**.
- [33] M. S. A. Abbasi, M. A. Tahir, and S. Meer, "FTIR Spectroscopic Study of *Aloe vera barbadensis* Mill Buds," *Asian J. Chem. Sci.*, pp. 1–6, **2020**.
- [34] M. A. Tahir and M. S. A. Abbasi, "FTIR Spectroscopic Analysis of *Mentha spicata* L.(Garden Mint)," *Asian J. Chem. Sci.*, pp. 1–5, **2020**.
- [35] D. Somit, D. Priyankar, and C. T. Kumar, "Quantification and correlation of the bioactive phytochemicals of *Croton bonplandianum* leaves of Sub-Himalayan region of West Bengal," *Asian J Pharm Clin Res*, vol. 6, no. 3, pp. 142–147, **2013**.
- [36] W.-H. Xu, W.-Y. Liu, and Q. Liang, "Chemical constituents from *Croton* species and their biological activities," *Molecules*, vol. 23, no. 9, p. 2333, **2018**.
- [37] F. Li *et al.*, "Bioactive terpenoids from *Croton laui*," *Nat. Prod. Res.*, pp. 1–9, 2019.
- [38] D. L. Pavia, G. M. Lampman, G. S. Kriz, and J. A. Vyvyan, *Introduction to spectroscopy*. Cengage Learning, **2008**.
- [39] A. Ortega, F. J. Morales, and M. Salmon, "Kaurenic acid derivatives from *Stevia eupatoria*," *Phytochemistry*, vol. 24, no. 8, pp. 1850–1852, **1985**.
- [40] V. Vennila and R. Udayakumar, "Analysis of Bioactive Compounds in Latex of *Croton bonplandianum* (Baill.) Using GC-MS," *J. Adv. Med. Pharm. Sci.*, pp. 1–7, **2016**.
- [41] T. Jasson, F. Nchu, and C. P. Laubscher, "NUTRIENT AVAILABILITY IN THE SOIL MEDIUM AND THE WET AND DRY WEIGHTS OF *HYPOXIS HEMEROCALLIDEA* AND *SIPHONCHILUS AETHIOPICUS* EXPOSED TO VARIOUS MUSHROOM COMPOST TEA EXTRACTS.," *Eff. Compost TEA Extr. GROWTH, Nutr. VALUE, SOIL Qual. HYPOXIS HEMEROCALLIDEA SIPHONCHILUS AETHIOPICUS.*, p. 65, **2017**.
- [42] W. D. Kristanto, "Study of characteristics and antioxidant activity of modified curcumin and its utilization in edible film." Universitas Pelita Harapan, **2017**.